

EQUIVARIANT TWISTED INVERSE WITHOUT EQUIVARIANT COMPACTIFICATION

MITSUYASU HASHIMOTO

Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya
464-8602 JAPAN

Abstract

An equivariant version of the twisted inverse pseudofunctor is defined, and equivariant versions of some important properties, including the Grothendieck duality of proper morphisms and flat base change are proved. As an application, a generalized version of Watanabe's theorem on the Gorenstein property of the ring of invariants is proved.

Contents

1	Introduction	3
2	Commutativity of diagrams	6
3	Sheaves on ringed sites	12
4	Derived category and derived functors of sheaves	18
5	Sheaves over a diagram of S -schemes	29
6	The inductions, the direct image, and the inverse image	32
7	Operations on sheaves via the structure data	34
8	Quasi-coherent sheaves over a diagram of schemes	42

9	Derived functors	48
10	The right adjoint of the derived direct image functor	55
11	The Composition of two pseudofunctors	58
12	Morphism of diagrams	63
13	Bökstedt–Neeman resolutions and hyperExt sheaves	65
14	Commutativity of twisted inverse with restrictions	69
15	Open immersion base change	77
16	The existence of compactification and composition data	79
17	Flat base change	81
18	Preservation of Quasi-coherent cohomology	84
19	Compatibility with derived direct image	86
20	Compatibility with derived right inductions	86
21	Equivariant Grothendieck’s duality	88
22	Morphisms of finite flat dimension	88
23	Finite morphisms of fiber type	92
24	Regular embeddings and Smooth morphisms of fiber type	95
25	Simplicial objects	101
26	Descent theory	104
27	Local noetherian property	110
28	Groupoid of schemes	113
29	Comparison of local Ext sheaves	117

30 Group schemes flat of finite type	118
31 Compatibility with derived G-invariance	120
32 Equivariant dualizing complexes and canonical modules	121
33 A generalization of Watanabe's theorem	126

1 Introduction

Let S be a separated noetherian scheme, G a flat separated S -group scheme of finite type, X and Y S -schemes separated of finite type with G -actions, and $f : X \rightarrow Y$ a G -morphism.

The purpose of these notes is to construct an equivariant version of the twisted inverse functor $f^!$.

One of the main motivation of the work is an application to the invariant theory. As an example, we give a very short proof of a generalized version of Watanabe's theorem on the Gorenstein property of invariant subrings [40]. Also, there might be some meaning in formulating the equivariant duality theorem, which has the Serre duality for the representations of reductive groups (see [20]) as a special case, in a reasonably general form for records.

In the case where G is trivial, $f^!$ is defined as follows. For a scheme Z , we denote the category of $\mathcal{O}_Z$ -modules by $\text{Mod}(Z)$. By definition, a plump subcategory of an abelian category is a non-empty full subcategory which is closed under kernels, cokernels, and extensions [23, (1.9.1)]. We denote the plump subcategory of $\text{Mod}(Z)$ consisting of quasi-coherent $\mathcal{O}_Z$ -modules by $\text{Qco}(Z)$.

By Nagata's compactification theorem, there exists some factorization

$$X \xrightarrow{i} \bar{X} \xrightarrow{p} Y$$

such that p is proper and i an open immersion. We call such a factorization a *compactification*. We define $f^! : D_{\text{Qco}(Y)}^+(\text{Mod}(Y)) \rightarrow D_{\text{Qco}(X)}^+(\text{Mod}(X))$ to be the composite $i^*p^\times$, where $p^\times : D_{\text{Qco}(Y)}^+(\text{Mod}(Y)) \rightarrow D_{\text{Qco}(\bar{X})}^+(\text{Mod}(\bar{X}))$ is the right adjoint of Rp_* , and i^* is the restriction. The definition of $f^!$ is independent of choice of compactification.

In order to consider a non-trivial G , we need to replace $\text{Qco}(X)$ and $\text{Mod}(X)$ by some appropriate categories which respect G -actions. The cate-

gory $\mathrm{Qco}(G, X)$ which corresponds to $\mathrm{Qco}(X)$ is fairly well-known. It is the category of G -linearized quasi-coherent $\mathcal{O}_X$ -modules defined by Mumford [27]. The category $\mathrm{Qco}(G, X)$ is equivalent to the category of quasi-coherent sheaves over the diagram of schemes

$$B_G^M(X) := \left(G \times_S G \times_S X \begin{array}{c} \xrightarrow{1_G \times a} \\ \xrightarrow{\mu \times 1_X} \\ \xrightarrow{p_{23}} \end{array} G \times_S X \begin{array}{c} \xrightarrow{a} \\ \xrightarrow{p_2} \end{array} X \right),$$

where $a : G \times X \rightarrow X$ is the action, $\mu : G \times G \rightarrow G$ the product, and p_{23} and p_2 are appropriate projections. Thus it is natural to embed the category $\mathrm{Qco}(G, X)$ into the category of all $\mathcal{O}_{B_G^M(X)}$ -modules $\mathrm{Mod}(B_G^M(X))$, and $\mathrm{Mod}(B_G^M(X))$ is a good substitute of $\mathrm{Mod}(X)$. As G is flat, $\mathrm{Qco}(G, X)$ is a plump subcategory of $\mathrm{Mod}(B_G^M(X))$, and we may consider $D_{\mathrm{Qco}(G, X)}(\mathrm{Mod}(B_G^M(X)))$. However, our construction utilize an intermediate category $\mathrm{LQco}(G, X)$ (the category of locally quasi-coherent sheaves), and is not an obvious interpretation of the non-equivariant case.

Note that there is a natural restriction functor $\mathrm{Mod}(B_G^M(X)) \rightarrow \mathrm{Mod}(X)$, which sends $\mathrm{Qco}(G, X)$ to $\mathrm{Qco}(X)$. This functor is regarded as the forgetful functor, forgetting the G -action. The equivariant duality theorem which we are going to establish must be compatible with this restriction functor, otherwise the theory would be something different from the usual scheme theory and probably useless.

Most of the discussion in these notes treats more general diagrams of schemes. This makes the discussion easier, as some of the important properties are proved by induction on the number of objects in the diagram. Our main construction and theorems are only for the class of finite diagrams of schemes of certain type, which contains the diagrams of the form $B_G^M(X)$.

In sections 2–4, we review general facts on homological algebra. The construction of $f^!$ is divided into five steps. The first is to analyze the functoriality of sheaves over diagrams of schemes. Sections 5–8 are devoted to this step. The second is the derived version of the first step. This will be done in sections 9, 10, and 13. Note that not only the categories of all module sheaves $\mathrm{Mod}(X_\bullet)$ and the category of quasi-coherent sheaves $\mathrm{Qco}(X_\bullet)$, but also the category of locally quasi-coherent sheaves $\mathrm{LQco}(X_\bullet)$ also plays an important role in our construction.

The third is to prove the existence of the right adjoint $p_\bullet^\times$ of $R(p_\bullet)_*$ for (componentwise) proper morphism $p_\bullet$ of diagrams of schemes. This is not

so difficult, and is done in section 12. We use Neeman's existence theorem on the right adjoint of unbounded right derived functors. Not only to utilize Neeman's theorem, but to calculate composite of various left and right derived functors, it is convenient to utilize unbounded derived functors. A short survey on unbounded derived functors is given in section 4.

The fourth step is to prove various commutativity related to the well-definedness of the twisted inverse pseudofunctors, sections 11, 14, and 15. Among them, the compatibility with restrictions (Proposition 14.14) is the key to our construction. Given a G -morphism $f : X \rightarrow Y$ between G -schemes separated of finite type over S , the associated morphism $B_G^M(f) : B_G^M(X) \rightarrow B_G^M(Y)$ is of fiber type (or cartesian), see for the definition, (5.2). If we could find a compactification

$$B_G^M(X) \xrightarrow{i_\bullet} Z_\bullet \xrightarrow{p_\bullet} B_G^M(Y)$$

such that $p_\bullet$ is proper and of fiber type, and $i_\bullet$ an image-dense open immersion, then the construction of $f^!$ and the proof of various commutative diagrams would be fairly easy. However, it seems that this is almost the same as the problem of equivariant compactifications. See [35] for equivariant compactifications. We avoid this problem, and prove the commutativity of various diagrams without assuming that $p_\bullet$ is of fiber type.

The fifth part is the existence of factorization $f_\bullet = p_\bullet i_\bullet$, where $p_\bullet$ is proper and $i_\bullet$ an image-dense open immersion. This is easily done utilizing Nagata's compactification theorem, and is done in section 16. This completes the basic construction of the equivariant twisted inverse pseudofunctor.

In sections 17–24, we prove equivariant versions of most of the known results on twisted inverse including the equivariant Grothendieck duality and flat base change, except that the equivariant dualizing complexes are treated later. We also prove that the twisted inverse functor preserves quasi-coherent cohomology groups. As we already know the corresponding results on single schemes and the commutativity with restrictions, this consists in straightforward (but not easy) check of various commutativity of diagrams of functors.

So far, almost all results are valid for any diagram of separated noetherian schemes with flat arrows over a finite ordered category. In sections 25–31, we prove some results which are special on the G -actions.

In section 32, we give a definition of the equivariant dualizing complexes. As an application, we give a very short proof of a generalized version of Watanabe's theorem on the Gorenstein property of invariant subrings in section 33.

Acknowledgement: The author is grateful to Professor Ryoshi Hotta and Professor Joseph Lipman for valuable advice.

2 Commutativity of diagrams constructed from a monoidal pair of pseudofunctors

(2.1) Let $(?)_*$ be a covariant symmetric monoidal pseudofunctor [23, (3.6.5)] of closed categories on a category $\mathcal{S}$. Thus, for each $X \in \mathcal{S}$,

$$X_* = (M_X, \otimes, \mathcal{O}_X, \alpha, \lambda, \gamma, [?, *], \pi)$$

is a (symmetric monoidal) closed category (see e.g., [23, (3.5.1)]), where M_X is the underlying category, $\otimes : M_X \times M_X \rightarrow M_X$ the product structure, $\mathcal{O}_X \in M_X$ the unit object, $\alpha : (a \otimes b) \otimes c \cong a \otimes (b \otimes c)$ the associativity isomorphism, $\lambda : \mathcal{O}_X \otimes a \cong a$ the left unit isomorphism, $\gamma : a \otimes b \cong b \otimes a$ the twisting (symmetry) isomorphism, $[?, -] : M_X^{\text{op}} \times M_X \rightarrow M_X$ the internal hom, and

$$\pi : M_X(a \otimes b, c) \cong M_X(a, [b, c]) \quad (2.2)$$

the associative adjunction isomorphism of X , respectively.

In this paper, various (different) adjoint pairs appears almost everywhere. By abuse of notation, the unit (resp. the counit) of adjunction is usually simply denoted by the same symbol u (resp. ε). However, the unit map and the counit map arising from the adjunction (2.2) are denoted (by less worse abuse of notation) by (the same symbol)

$$\text{tr} : a \rightarrow [b, a \otimes b] \quad (\text{the trace map})$$

and

$$\text{ev} : [b, c] \otimes b \rightarrow c \quad (\text{the evaluation map}),$$

respectively.

(2.3) Let $f : X \rightarrow Y$ be a morphism. Then f_* is a symmetric monoidal functor. The natural map

$$f_*a \otimes f_*b \rightarrow f_*(a \otimes b)$$

is denoted by $m = m(f)$, and the map

$$\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$$

is denoted by $\eta = \eta(f)$.

(2.4) For composable two morphisms f and g in $\mathcal{S}$, the given isomorphism $(gf)_* \cong g_*f_*$ is denoted by $c = c_{f,g}$ as in [23, (3.6.5)]. Let $gf = f'g'$ be a commutative diagram in $\mathcal{S}$. The composite isomorphism

$$g_*f_* \xrightarrow{c^{-1}} (gf)_* = (f'g')_* \xrightarrow{c} f'_*g'_*$$

is also denoted by $c = c(gf = f'g')$, by abuse of notation.

(2.5) Let $f : X \rightarrow Y$ be a morphism in $\mathcal{S}$. The composite natural map

$$f_*[a, b] \xrightarrow{\text{tr}} [f_*a, f_*[a, b] \otimes f_*a] \xrightarrow{\text{via } m} [f_*a, f_*([a, b] \otimes a)] \xrightarrow{\text{via ev}} [f_*a, f_*b]$$

is denoted by H . In other words, H is right conjugate [23, (3.3.5)] to the map

$$f_*[a, b] \otimes f_*a \xrightarrow{m} f_*([a, b] \otimes a) \xrightarrow{f_*\text{ev}} f_*b.$$

(2.6) G. Lewis proved a theorem which guarantee that some diagrams involving two symmetric monoidal closed categories and one symmetric monoidal functor commute [22].

By Lewis' result, we have that the following diagrams are commutative for any morphism $f : X \rightarrow Y$ (also checked by direct computation).

$$\begin{array}{ccc} f_*[a, b] \otimes f_*a & \xrightarrow{H \otimes 1} & [f_*a, f_*b] \otimes f_*a \\ \downarrow m & & \downarrow \text{ev} \\ f_*([a, b] \otimes a) & \xrightarrow{f_*\text{ev}} & f_*b \end{array} \quad (2.7)$$

$$\begin{array}{ccc} f_*a & \xrightarrow{f_*\text{tr}} & f_*[b, a \otimes b] \\ \downarrow \text{tr} & & \downarrow H \\ [f_*b, f_*a \otimes f_*b] & \xrightarrow{m} & [f_*b, f_*(a \otimes b)] \end{array} \quad (2.8)$$

$$\begin{array}{ccccccc} f_*a & \xrightarrow{\lambda^{-1}} & \mathcal{O}_Y \otimes f_*a & \xrightarrow{\eta} & f_*\mathcal{O}_X \otimes f_*a & \xrightarrow{\text{via tr}} & f_*[a, \mathcal{O}_X \otimes a] \otimes f_*a \\ \downarrow 1 & & & & & & \downarrow \text{via } \lambda \\ f_*a & \xleftarrow{\text{ev}} & [f_*a, f_*a] \otimes f_*a & \xleftarrow{H \otimes 1} & f_*[a, a] \otimes f_*a & & \end{array} \quad (2.9)$$

2.10 Lemma. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in $\mathcal{S}$. Then the diagram

$$\begin{array}{ccccc} g_*f_*[a, b] & \xrightarrow{H} & g_*[f_*a, f_*b] & \xrightarrow{H} & [g_*f_*a, g_*f_*b] \\ \uparrow c & & & & \uparrow [c^{-1}, c] \\ (gf)_*[a, b] & \xrightarrow{H} & [(gf)_*a, (gf)_*b] & & \end{array}$$

is commutative.

Proof. Consider the diagram

$$\begin{array}{ccccccc}
g_* f_*[a, b] \otimes g_* f_* a & \xrightarrow{m} & g_*(f_*[a, b] \otimes f_* a) & \xrightarrow{m} & g_* f_*([a, b] \otimes a) & \xrightarrow{\text{via ev}} & g_* f_* b \\
\uparrow c \otimes c & & \sigma & & \downarrow c^{-1} & \tau & \downarrow c^{-1} \\
(gf)_*[a, b] \otimes (gf)_* a & \xrightarrow{m} & (gf)_*([a, b] \otimes a) & \xrightarrow{\text{via ev}} & (gf)_* b.
\end{array}$$

The diagram σ is commutative by the definition of monoidal pseudofunctor, see [23, (3.6.7.2)]. The commutativity of τ is obvious. Taking the conjugate, we get the commutativity of the diagram in the lemma. $\square$

(2.11) Let us assume that there is a left adjoint $(?)^*$ of $(?)_*$ so that $((?)^*, (?)_*)$ is a monoidal adjoint pair [23, pp. 107–109].

For composable two morphisms f and g in $\mathcal{S}$, we denote the map $f^* g^* \rightarrow (gf)^*$ conjugate to $c : (gf)_* \rightarrow g_* f_*$ by $d = d_{f,g}$. Being the conjugate of an isomorphism, d is an isomorphism. For a commutative diagram $gf = f'g'$ in $\mathcal{S}$, the composite map

$$(g')^*(f')^* \xrightarrow{d} (f'g')^* = (gf)^* \xrightarrow{d^{-1}} f^* g^*$$

is also denoted by d , by abuse of notation. For a morphism $f : X \rightarrow Y$, the map $f^* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ conjugate to $\eta : \mathcal{O}_Y \rightarrow f_* \mathcal{O}_X$ is denoted by C . The composite map

$$f^*(a \otimes b) \xrightarrow{\text{via } u} f^*(f_* f^* a \otimes f_* f^* b) \xrightarrow{\text{via } m} f^* f_*(f^* a \otimes f^* b) \xrightarrow{\varepsilon} f^* a \otimes f^* b \quad (2.12)$$

is denoted by Δ .

Almost by definition, the diagrams

$$\begin{array}{ccc}
a \otimes b & \xrightarrow{u \otimes u} & f_* f^* a \otimes f_* f^* b \\
\downarrow u & & \downarrow m \\
f_* f^*(a \otimes b) & \xrightarrow{\Delta} & f_*(f^* a \otimes f^* b)
\end{array} \quad (2.13)$$

and

$$\begin{array}{ccc}
f^*(f_* a \otimes f_* b) & \xrightarrow{m} & f^* f_*(a \otimes b) \\
\downarrow \Delta & & \downarrow \varepsilon \\
f^* f_* a \otimes f^* f_* b & \xrightarrow{\varepsilon \otimes \varepsilon} & a \otimes b
\end{array} \quad (2.14)$$

are commutative.

2.15 Definition. A monoidal adjoint pair $((?)^*, (?)_*)$ is said to be *Lipman* if $\Delta : f^*(a \otimes b) \rightarrow f^* a \otimes f^* b$ and $C : f^* \mathcal{O}_Y \rightarrow \mathcal{O}_X$ are isomorphisms for any morphism $f : X \rightarrow Y$ in $\mathcal{S}$ and any $a, b \in M_Y$.

(2.16) Note that $\Delta : f^*(a \otimes b) \rightarrow f^*a \otimes f^*b$ is a natural isomorphism if and only if its right conjugate (see [23, (3.3.5)]) is an isomorphism. The right conjugate is the composite

$$f_*[f^*b, a] \xrightarrow{H} [f_*f^*b, f_*a] \xrightarrow{u} [b, f_*a].$$

Let $((?)^*, (?)_*)$ be a Lipman adjoint pair of monoidal pseudofunctors over $\mathcal{S}$. Then $(?)^*$ together with Δ^{-1} and C^{-1} form a *covariant* symmetric monoidal pseudofunctor on $\mathcal{S}^{\text{op}}$.

(2.17) For a morphism $f : X \rightarrow Y$ in $\mathcal{S}$, the composite map

$$f^*[a, b] \xrightarrow{\text{via tr}} [f^*a, f^*[a, b] \otimes f^*a] \xrightarrow{\text{via } \Delta^{-1}} [f^*a, f^*([a, b] \otimes a)] \xrightarrow{\text{via ev}} [f^*a, f^*b]$$

is denoted by P . We can apply Lewis' theorem to f^* . In particular, the following diagrams are commutative by (2.6) for a morphism $f : X \rightarrow Y$.

$$\begin{array}{ccc} f^*[a, b] \otimes f^*a & \xrightarrow{P \otimes 1} & [f^*a, f^*b] \otimes f^*a \\ \downarrow \Delta^{-1} & & \downarrow \text{ev} \\ f^*([a, b] \otimes a) & \xrightarrow{f^* \text{ev}} & f_*b \end{array} \quad (2.18)$$

$$\begin{array}{ccc} f^*a & \xrightarrow{f^* \text{tr}} & f^*[b, a \otimes b] \\ \downarrow \text{tr} & & \downarrow P \\ [f^*b, f^*a \otimes f^*b] & \xrightarrow{\Delta^{-1}} & [f^*b, f^*(a \otimes b)] \end{array} \quad (2.19)$$

$$\begin{array}{ccccccc} f^*a & \xrightarrow{\lambda^{-1}} & \mathcal{O}_X \otimes f^*a & \xrightarrow{C^{-1}} & f^*\mathcal{O}_Y \otimes f^*a & \xrightarrow{\text{via tr}} & f^*[a, \mathcal{O}_Y \otimes a] \otimes f^*a \\ \downarrow 1 & & & & & & \downarrow \text{via } \lambda \\ f^*a & \xleftarrow{\text{ev}} & [f^*a, f^*a] \otimes f^*a & \xleftarrow{P \otimes 1} & f^*[a, a] \otimes f^*a & & \end{array} \quad (2.20)$$

2.21 Lemma. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in $\mathcal{S}$. Then the diagram

$$\begin{array}{ccccc} f^*g^*[a, b] & \xrightarrow{P} & f^*[g^*a, g^*b] & \xrightarrow{P} & [f^*g^*a, f^*g^*b] \\ \uparrow d^{-1} & & & & \uparrow [d, d^{-1}] \\ (gf)^*[a, b] & \xrightarrow{P} & & & [(gf)^*a, (gf)^*b] \end{array}$$

is commutative.

Proof. Follows instantly by Lemma 2.10. $\square$

2.22 Lemma. *The following diagrams are commutative.*

$$\begin{array}{ccc}
[a, b] & \xrightarrow{u} & [a, f_* f^* b] \\
\downarrow u & & \uparrow u \\
f_* f^* [a, b] & \xrightarrow{HP} & [f_* f^* a, f_* f^* b] \\
\\
f^* f_* [a, b] & \xrightarrow{PH} & [f^* f_* a, f^* f_* b] \\
\downarrow \varepsilon & & \downarrow \varepsilon \\
[a, b] & \xrightarrow{\varepsilon} & [f^* f_* a, b]
\end{array}$$

Proof. We prove the commutativity of the first diagram. Taking the adjoint map, it suffices to prove the commutativity of

$$\begin{array}{ccc}
[a, b] \otimes a & \xrightarrow{\text{ev}} & b \\
\downarrow u \otimes u & & \downarrow u \\
f_* f^* [a, b] \otimes f_* f^* a & \xrightarrow{HP \otimes 1} & [f_* f^* a, f_* f^* b] \otimes f_* f^* a \xrightarrow{\text{ev}} f_* f^* b.
\end{array}$$

This is obvious by the naturality of u and the commutativity of (2.7).

The commutativity of the second diagram is similar (use (2.18)). $\square$

(2.23) Let X be an object of $\mathcal{S}$. We denote the composite isomorphism

$$M_X(a, b) \xrightarrow{\text{via } \lambda} M_X(\mathcal{O}_X \otimes a, b) \xrightarrow{\pi} M_X(\mathcal{O}_X, [a, b])$$

by h_X .

2.24 Lemma. *Let $f : X \rightarrow Y$ be a morphism in $\mathcal{S}$. Then the composite map*

$$\begin{aligned}
M_X(a, b) &\xrightarrow{h_X} M_X(\mathcal{O}_X, [a, b]) \xrightarrow{C} M_X(f^* \mathcal{O}_Y, [a, b]) \cong M_Y(\mathcal{O}_Y, f_* [a, b]) \\
&\xrightarrow{H} M_Y(\mathcal{O}_Y, [f_* a, f_* b]) \xrightarrow{h_Y^{-1}} M_Y(f_* a, f_* b)
\end{aligned}$$

agrees with the map given by $\varphi \mapsto f_* \varphi$. The composite map

$$\begin{aligned}
M_Y(a', b') &\xrightarrow{h_Y} M_Y(\mathcal{O}_Y, [a', b']) \xrightarrow{f^*} M_X(f^* \mathcal{O}_Y, f^* [a', b']) \\
&\xrightarrow{M_X(C^{-1}, P)} M_X(\mathcal{O}_X, [f^* a', f^* b']) \xrightarrow{h_X^{-1}} M_X(f^* a', f^* b')
\end{aligned}$$

agrees with f^* .

Proof. We prove the first assertion. All the maps are natural on a . By Yoneda's lemma, we may assume that $a = b$ and it suffices to show that the identity map 1_b is mapped to 1_{f_*b} by the map. It is straightforward to check that 1_b goes to the composite map

$$\begin{aligned} f_*a &\xrightarrow{\lambda^{-1}} \mathcal{O}_Y \otimes f_*a \xrightarrow{u} f_*f^*\mathcal{O}_Y \otimes f_*a \xrightarrow{C} f_*\mathcal{O}_X \otimes f_*a \\ &\xrightarrow{\text{via tr}} f_*[a, \mathcal{O}_X \otimes a] \otimes f_*a \xrightarrow{\lambda} f_*[a, a] \otimes f_*a \xrightarrow{\text{ev}} f_*a. \end{aligned}$$

By the commutativity of (2.9), we are done.

The second assertion is proved similarly, utilizing the commutativity of (2.20). $\square$

Let $((?)^*, (?)_*)$ be a monoidal pair over $\mathcal{S}$ which may not be Lipman. Let $\sigma = (fg' = gf')$ be a commutative diagram in $\mathcal{S}$.

2.25 Lemma. *The following composite map agree:*

$$\begin{aligned} 1 \quad & g^*f_* \xrightarrow{u} g^*f_*g'_*(g')^* \xrightarrow{c} g^*g_*f'_*(g')^* \xrightarrow{\varepsilon} f'_*(g')^*; \\ 2 \quad & g^*f_* \xrightarrow{u} f'_*(f')^*g^*f_* \xrightarrow{d} f'_*(g')^*f^*f_* \xrightarrow{\varepsilon} f'_*(g')^*. \end{aligned}$$

For the proof and more information, see [23, (3.7.2)]. We denote the composite map in the lemma by $\theta(\sigma)$ or θ .

2.26 Lemma. *Let $\theta = (fg' = gf')$ be a commutative diagram in $\mathcal{S}$. Then the diagram*

$$\begin{array}{ccc} g^*f_*[a, b] & \xrightarrow{\theta} & f'_*(g')^*[a, b] \\ \downarrow PH & & \downarrow HP \\ [g^*f_*a, g^*f_*b] & \xrightarrow{\theta} [g^*f_*a, f'_*(g')^*b] & \xleftarrow{\theta} [f'_*(g')^*a, f'_*(g')^*b] \end{array}$$

is commutative.

Proof. Follows from Lemma 2.22. $\square$

2.27 Lemma. *Let $\theta = (fg' = gf')$ be a commutative diagram in $\mathcal{S}$. Then the diagram*

$$\begin{array}{ccccc} f^*(g_*a \otimes g_*b) & \xrightarrow{m} & f^*g_*(a \otimes b) & \xrightarrow{\theta} & g'_*(f')^*(a \otimes b) \\ \downarrow \Delta & & & & \downarrow \Delta \\ f^*g_*a \otimes f^*g_*b & \xrightarrow{\theta \otimes \theta} & g'_*(f')^*a \otimes g'_*(f')^*b & \xrightarrow{m} & g'_*((f')^*a \otimes (f')^*b) \end{array}$$

is commutative.

Proof. Utilize (2.13) and (2.14). $\square$

3 Sheaves on ringed sites

(3.1) We fix a universe $\mathcal{U}$, and the categories of sets, abelian groups, rings, etc., denoted by $\underline{\text{Set}}$, $\underline{\text{Ab}}$, $\underline{\text{Rng}}$, etc., are those consisting of sets, abelian groups, rings, etc. inside $\mathcal{U}$. In the sequel, a $\mathcal{U}$ -small set (i.e., a set which is bijective to an element of $\mathcal{U}$) is referred simply as a small set.

(3.2) For categories I and $\mathcal{C}$, we denote the functor category $\text{Func}(I^{\text{op}}, \mathcal{C})$ by $\mathcal{P}(I, \mathcal{C})$. An object of $\mathcal{P}(I, \mathcal{C})$ is sometimes referred as a presheaf over I with values in $\mathcal{C}$. We fix a universe $\mathcal{V}$ which contains $\mathcal{U}$ as an element. If $\mathcal{C}$ is a $\mathcal{V}$ -category and I is $\mathcal{V}$ -svelte (i.e., equivalent to a $\mathcal{V}$ -small category), then $\mathcal{P}(I, \mathcal{C})$ is a $\mathcal{V}$ -category. The category $\mathcal{P}(I, \underline{\text{Ab}})$ is denoted by $\text{PA}(I)$.

In this article, a site (i.e., a category with a Grothendieck topology, in the sense of [37]) is required to be a $\mathcal{V}$ -small $\mathcal{U}$ -site defined by a pretopology. If $\mathbb{X}$ is a site, then the category of sheaves on $\mathbb{X}$ with values in $\mathcal{C}$ is denoted by $\mathcal{S}(\mathbb{X}, \mathcal{C})$. If $\mathcal{C}$ is a $\mathcal{U}$ -category, then $\mathcal{S}(\mathbb{X}, \mathcal{C})$ is also a $\mathcal{U}$ -category, by our requirement. The category $\mathcal{S}(\mathbb{X}, \underline{\text{Ab}})$ is denoted by $\text{AB}(\mathbb{X})$.

Let $\mathbb{X}$ be a site. The inclusion $\text{AB}(\mathbb{X}) \rightarrow \text{PA}(\mathbb{X})$ is denoted by $q(\mathbb{X}, \text{AB})$. We denote the sheafification functor $\text{PA}(\mathbb{X}) \rightarrow \text{AB}(\mathbb{X})$ by $a(\mathbb{X}, \text{AB})$.

(3.3) Let $\mathbb{X} = (\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. Namely, let $\mathbb{X}$ be a site and $\mathcal{O}_{\mathbb{X}}$ a sheaf of commutative rings on $\mathbb{X}$. We denote the category of presheaves (resp. sheaves) of $\mathcal{O}_{\mathbb{X}}$ -modules by $\text{PM}(\mathbb{X})$ (resp. $\text{Mod}(\mathbb{X})$). The inclusion $\text{Mod}(\mathbb{X}) \rightarrow \text{PM}(\mathbb{X})$ is denoted by $q(\mathbb{X}, \text{Mod})$. The sheafification $\text{PM}(\mathbb{X}) \rightarrow \text{Mod}(\mathbb{X})$ is denoted by $a(\mathbb{X}, \text{Mod})$. The forgetful functor $\text{Mod}(\mathbb{X}) \rightarrow \text{AB}(\mathbb{X})$ is denoted by $F(\mathbb{X})$. The categories $\text{AB}(\mathbb{X})$ and $\text{Mod}(\mathbb{X})$ are Grothendieck in the sense that they have small sets of generators and satisfy the (AB5) condition in [10]. They also satisfy (AB3*).

(3.4) Let $f : \mathbb{Y} \rightarrow \mathbb{X}$ be a functor. Then the pull-back $\text{PA}(\mathbb{X}) \rightarrow \text{PA}(\mathbb{Y})$ is denoted by $f_{\text{PA}}^{\#}$. Note that $f_{\text{PA}}^{\#}(\mathcal{F}) := \mathcal{F} \circ f^{\text{op}}$. In general, the pull-back $\mathcal{P}(\mathbb{X}, \mathcal{C}) \rightarrow \mathcal{P}(\mathbb{Y}, \mathcal{C})$ is defined in a similar way, and is denoted by $f^{\#}$. If f is continuous (i.e., $f_{\underline{\text{Set}}}^{\#}$ carries sheaves to sheaves), then $f_{\text{AB}}^{\#} : \text{AB}(\mathbb{X}) \rightarrow \text{AB}(\mathbb{Y})$ is defined to be the restriction of $f_{\text{PA}}^{\#}$. Throughout these notes, we require that a continuous functor $f : \mathbb{Y} \rightarrow \mathbb{X}$ satisfies the following condition. For $y \in \mathbb{Y}$, a covering $(y_i \rightarrow y)_{i \in I}$, and any $i, j \in I$, the morphisms $f(y_i \times_y y_j) \rightarrow f(y_i)$ and $f(y_i \times_y y_j) \rightarrow f(y_j)$ makes $f(y_i \times_y y_j)$ the fiber product $f(y_i) \times_{f(y)} f(y_j)$.

Thanks to the re-definition of sites and continuous functors, we have the following.

3.5 Lemma. *Let $f : \mathbb{Y} \rightarrow \mathbb{X}$ be a functor between sites. Then f is continuous if and only if the following holds.*

If $(\varphi_i : y_i \rightarrow y)_{i \in I}$ is a covering, then $(f\varphi_i : fy_i \rightarrow fy)_{i \in I}$ is a covering.

For the proof, see [13, (III.1.6)].

The left adjoint of $f_{\text{PA}}^\#$, which exists by Kan's lemma (see e.g., [2, Theorem I.2.1]), is denoted by $f_\#^{\text{PA}}$.

For $x \in \mathbb{X}$, we define the category I_x^f as follows. An object of I_x^f is a pair (y, ϕ) with $y \in \mathbb{Y}$ and $\phi \in \mathbb{X}(x, f(y))$. A morphism $h : (y, \phi) \rightarrow (y', \phi')$ is a morphism $h \in \mathbb{Y}(y, y')$ such that $f(h) \circ \phi = \phi'$. Note that $\Gamma(x, f_\#^{\text{PA}}(\mathcal{F})) = \varinjlim \Gamma(y, \mathcal{F})$, where the colimit is taken over $(I_x^f)^{\text{op}}$.

The left adjoint of $f_{\mathcal{C}}^\# : \mathcal{P}(\mathbb{X}, \mathcal{C}) \rightarrow \mathcal{P}(\mathbb{X}, \mathcal{C})$ is constructed similarly, provided $\mathcal{C}$ is a $\mathcal{U}$ -category with arbitrary small colimits. The left adjoint is denoted by $f_\#^{\mathcal{C}}$ or simply by $f_\#$.

Similarly, $f_\#^{\text{AB}} : \text{AB}(\mathbb{X}) \rightarrow \text{AB}(\mathbb{Y})$ and its left adjoint $f_{\text{AB}}^\#$ is defined. Note that we have $q(\mathbb{Y}, \text{AB}) \circ f_{\text{AB}}^\# = f_{\text{PA}}^\# \circ q(\mathbb{X}, \text{AB})$, and $f_\#^{\text{AB}} = a(\mathbb{X}, \text{AB}) \circ f_\#^{\text{PA}} \circ q(\mathbb{Y}, \text{AB})$.

3.6 Lemma. *If $(I_x^f)^{\text{op}}$ is pseudofiltered (see e.g., [13, 26]) for each $x \in \mathbb{X}$, then $f_\#^{\text{PA}}$ is exact.*

Proof. This is a consequence of [13, Proposition 2.8]. □

We say that $f : \mathbb{Y} \rightarrow \mathbb{X}$ is *admissible* if f is continuous and the functor $f_\#^{\text{PA}}$ is exact.

(3.7) If $\mathbb{Y}$ has finite limits and f preserves finite limits, then $f_\#^{\text{PA}}$ is exact by the lemma. It follows that a continuous map induces an admissible continuous functor between the corresponding sites.

(3.8) The right adjoint functor of $f_{\text{PA}}^\#$, which we denote by f_{b}^{PA} also exists, as $\underline{\text{Ab}}^{\text{op}}$ has arbitrary small colimits (i.e., $\underline{\text{Ab}}$ has small limits). The functor f_{b}^{PA} is the left adjoint functor

$$\text{Func}(\mathbb{Y}^{\text{op}}, \underline{\text{Ab}}) \xrightarrow{\text{op}} \text{Func}(\mathbb{Y}, \underline{\text{Ab}}^{\text{op}}) \rightarrow \text{Func}(\mathbb{X}, \underline{\text{Ab}}^{\text{op}}) \xrightarrow{\text{op}} \text{Func}(\mathbb{X}^{\text{op}}, \underline{\text{Ab}})$$

of the functor

$$(f^{\text{op}})^\# : \text{Func}(\mathbb{X}, \underline{\text{Ab}}^{\text{op}}) \rightarrow \text{Func}(\mathbb{Y}, \underline{\text{Ab}}^{\text{op}}),$$

where $f^{\text{op}} = f$ is the opposite of f , namely, f viewed as a functor $\mathbb{Y}^{\text{op}} \rightarrow \mathbb{X}^{\text{op}}$.

(3.9) For $\mathcal{M}, \mathcal{N} \in \text{PM}(\mathbb{X})$, the presheaf tensor product is denoted by $\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p \mathcal{N}$. It is defined by

$$\Gamma(x, \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p \mathcal{N}) := \Gamma(x, \mathcal{M}) \otimes_{\Gamma(x, \mathcal{O}_{\mathbb{X}})} \Gamma(x, \mathcal{N})$$

for $x \in \mathbb{X}$. The sheaf tensor product $a(q\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p q\mathcal{N})$ of $\mathcal{M}, \mathcal{N} \in \text{Mod}(\mathbb{X})$ is denoted by $\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}} \mathcal{N}$.

(3.10) We say that $f : (\mathbb{Y}, \mathcal{O}_{\mathbb{Y}}) \rightarrow (\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ is a continuous ringed functor if $f : \mathbb{Y} \rightarrow \mathbb{X}$ is a continuous functor, and a morphism of sheaves of commutative rings $\mathcal{O}_{\mathbb{Y}} \rightarrow f_{\text{AB}}^{\#}(\mathcal{O}_{\mathbb{X}})$ is given. The pull-back $\text{PM}(\mathbb{X}) \rightarrow \text{PM}(\mathbb{Y})$ is denoted by $f_{\text{PM}}^{\#}$, and its left adjoint is denoted by $f_{\#}^{\text{PM}}$. The left adjoint $f_{\#}^{\text{PM}}$ is defined by

$$\Gamma(x, f_{\#}^{\text{PM}} \mathcal{M}) := \varinjlim \Gamma(x, \mathcal{O}_{\mathbb{X}}) \otimes_{\Gamma(y, \mathcal{O}_{\mathbb{Y}})} \Gamma(y, \mathcal{M})$$

for $x \in \mathbb{X}$ and $\mathcal{M} \in \text{PM}(\mathbb{Y})$, where the colimit is taken over the category $(I_x^f)^{\text{op}}$. Similarly, $f_{\text{Mod}}^{\#} : \text{Mod}(\mathbb{X}) \rightarrow \text{Mod}(\mathbb{Y})$ and its left adjoint $f_{\#}^{\text{Mod}} = af_{\text{PM}}^{\#}q$ is defined. If $(I_x^f)^{\text{op}}$ is filtered for any $x \in \mathbb{X}$, then $f_{\#}^{\text{PA}} \mathcal{O}_{\mathbb{Y}}$ has a structure of a presheaf of rings in a natural way, and there is a canonical isomorphism $f_{\#}^{\text{PM}} \mathcal{M} \cong \mathcal{O}_{\mathbb{X}} \otimes_{f_{\#}^{\text{PA}} \mathcal{O}_{\mathbb{Y}}}^p f_{\#}^{\text{PA}} \mathcal{M}$. The right adjoint of $f_{\text{PM}}^{\#}$, which exists as in (3.8), is denoted by f_b^{PM} .

(3.11) Let $\mathbb{X}$ be a ringed site, and $x \in \mathbb{X}$. The category $\mathbb{X}/x$ is a site with the same topology as that of $\mathbb{X}$. The canonical functor $\Phi_x : \mathbb{X}/x \rightarrow \mathbb{X}$ is a continuous ringed functor, and yields the pull-backs $(\Phi_x)_{\text{AB}}^{\#}$ and $(\Phi_x)_{\text{PA}}^{\#}$, which we denote by $(?)|_x^{\text{AB}}$ and $(?)|_x^{\text{PA}}$, respectively. Their left adjoints are denoted by L_x^{AB} and L_x^{PA} , respectively. Note that Φ_x is admissible, see [26, p.78].

(3.12) Note that $\mathbb{X}/x$ is a ringed site with the structure sheaf $\mathcal{O}_{\mathbb{X}}|_x$. Thus, $(?)|_x^{\text{Mod}}$ and $(?)|_x^{\text{PM}}$ are defined in an obvious way, and their left adjoints L_x^{Mod} and L_x^{PM} are also defined. Note that L_x^{Mod} and L_x^{PM} are exact.

(3.13) For a morphism $\phi : x \rightarrow y$, we have an obvious admissible ringed continuous functor $\Phi_{\phi} : \mathbb{X}/x \rightarrow \mathbb{X}/y$. The corresponding pull-back is denoted by $\phi_{\natural}^*$, and its left adjoint is denoted by $\phi_{\star}^{\natural}$, where $\natural$ is AB, PA, Mod or PM.

For $\mathcal{M}, \mathcal{N} \in \natural(\mathbb{X})$, we define $\underline{\text{Hom}}_{\natural(\mathbb{X})}(\mathcal{M}, \mathcal{N})$ to be the object of $\natural(\mathbb{X})$ given by

$$\Gamma(x, \underline{\text{Hom}}_{\natural(\mathbb{X})}(\mathcal{M}, \mathcal{N})) := \text{Hom}_{\natural(\mathbb{X}/x)}(\mathcal{M}|_x^{\natural}, \mathcal{N}|_x^{\natural}),$$

where $\mathfrak{h} = \text{PA}, \text{AB}, \text{PM}, \text{ or } \text{Mod}$. For $\phi : x \rightarrow y$, the restriction map

$$\text{Hom}_{\mathfrak{h}(\mathbb{X}/y)}(\mathcal{M}|_y^{\mathfrak{h}}, \mathcal{N}|_y^{\mathfrak{h}}) \rightarrow \text{Hom}_{\mathfrak{h}(\mathbb{X}/x)}(\mathcal{M}|_x^{\mathfrak{h}}, \mathcal{N}|_x^{\mathfrak{h}})$$

is given by $\phi_{\mathfrak{h}}^*$.

The following is a slight generalization of [23, Example 3.5.2].

3.14 Lemma. *Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. Then we have the following.*

- 1 *The category $\text{PM}(\mathbb{X})$ is a closed symmetric monoidal category (see [24, p.180]) with $\otimes_{\mathcal{O}_{\mathbb{X}}}^p$ the multiplication, $q\mathcal{O}_{\mathbb{X}}$ the unit object, $\underline{\text{Hom}}_{\text{PM}(\mathbb{X})}(?, ?)$ the internal hom, etc., etc.*
- 2 *The category $\text{Mod}(\mathbb{X})$ is a closed symmetric monoidal category with $\otimes_{\mathcal{O}_{\mathbb{X}}}$ the multiplication, $\mathcal{O}_{\mathbb{X}}$ the unit object, $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}(?, ?)$ the internal hom, etc., etc.*
- 3 *The inclusion $q : \text{Mod}(\mathbb{X}) \rightarrow \text{PM}(\mathbb{X})$ and the natural transformations*

$$q\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p q\mathcal{N} \xrightarrow{u} qa(q\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p q\mathcal{N}) = q(\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}} \mathcal{N})$$

and

$$q\mathcal{O}_{\mathbb{X}} \xrightarrow{\text{id}} q\mathcal{O}_{\mathbb{X}}$$

form a symmetric monoidal functor.

- 4 *For a ringed continuous functor $f : (\mathbb{Y}, \mathcal{O}_{\mathbb{Y}}) \rightarrow (\mathbb{X}, \mathcal{O}_{\mathbb{X}})$, the functor $f_{\text{PM}}^{\#} : \text{PM}(\mathbb{X}) \rightarrow \text{PM}(\mathbb{Y})$, the natural map*

$$f_{\text{PM}}^{\#}\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{Y}}}^p f_{\text{PM}}^{\#}\mathcal{N} \rightarrow f_{\text{PM}}^{\#}(\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p \mathcal{N}) \quad (3.15)$$

which induces the canonical map

$$\Gamma(fy, \mathcal{M}) \otimes_{\Gamma(y, \mathcal{O}_{\mathbb{Y}})} \Gamma(fy, \mathcal{N}) \rightarrow \Gamma(fy, \mathcal{M}) \otimes_{\Gamma(fy, \mathcal{O}_{\mathbb{X}})} \Gamma(fy, \mathcal{N})$$

for each $y \in \mathbb{Y}$, and the map

$$q\mathcal{O}_{\mathbb{Y}} \rightarrow qf_{\text{Mod}}^{\#}\mathcal{O}_{\mathbb{X}} \cong f_{\text{PM}}^{\#}q\mathcal{O}_{\mathbb{X}}$$

form a symmetric monoidal functor.

5 Under the same assumption as in 4, the functor $f_{\text{Mod}}^\# : \text{Mod}(\mathbb{X}) \rightarrow \text{Mod}(\mathbb{Y})$, the natural map

$$\begin{aligned} f_{\text{Mod}}^\# \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{Y}}} f_{\text{Mod}}^\# \mathcal{N} &= a(q f_{\text{Mod}}^\# \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{Y}}}^p q f_{\text{Mod}}^\# \mathcal{N}) \cong a(f_{\text{PM}}^\# q \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{Y}}}^p f_{\text{PM}}^\# q \mathcal{N}) \\ &\xrightarrow{\text{via (3.15)}} a f_{\text{PM}}^\# (q \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p q \mathcal{N}) \xrightarrow{\text{via } \rho} f_{\text{Mod}}^\# a(q \mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}}^p q \mathcal{N}) = f_{\text{Mod}}^\# (\mathcal{M} \otimes_{\mathcal{O}_{\mathbb{X}}} \mathcal{N}), \end{aligned}$$

and the map

$$\mathcal{O}_{\mathbb{Y}} \rightarrow f_{\text{Mod}}^\# \mathcal{O}_{\mathbb{X}}$$

form a symmetric monoidal functor, where $\rho : a f_{\text{PM}}^\# \rightarrow f_{\text{Mod}}^\# a$ is the composite map

$$a f_{\text{PM}}^\# \xrightarrow{\text{via } u} a f_{\text{PM}}^\# q a \cong a q f_{\text{Mod}}^\# a \xrightarrow{\text{via } \varepsilon} f_{\text{Mod}}^\# a.$$

6 Let $\mathcal{S}$ denote the category of ringed sites (with svelte underlying categories). Then $((?)_{\#}^{\text{PM}}, (?)_{\text{PM}}^\#)$ and $((?)_{\#}^{\text{Mod}}, (?)_{\text{PM}}^\#)$ are adjoint pairs of monoidal pseudo-functors on $\mathcal{S}^{\text{op}}$, see [23, (3.6.7)].

(3.16) If there is no confusion, the $\natural$ attached to the functors of sheaves defined above are omitted. For example, $f^\#$ stands for $f_{\natural}^\#$. Note that $f_{\natural}^\#(\mathcal{F})$ viewed as a presheaf of abelian groups is independent of $\natural$.

(3.17) Let $f : \mathbb{Y} \rightarrow \mathbb{X}$ be a ringed continuous functor between ringed sites. The following is a restricted version of the results on cocontinuous functors in [38]. We give a proof for convenience of readers.

3.18 Lemma. Assume that for any $y \in \mathbb{Y}$ and any covering $(x_\lambda \rightarrow fy)_{\lambda \in \Lambda}$ of fy , there is a covering $(y_\mu \rightarrow y)_{\mu \in M}$ of y such that there is a map $\phi : M \rightarrow \Lambda$ such that $fy_\mu \rightarrow fy$ factors through $x_{\phi(\mu)} \rightarrow fy$ for each μ . Then the pull-back $f^\#$ is compatible with the sheafification in the sense that the canonical natural transformation

$$\rho : a(\mathbb{Y}, \text{Mod}) f_{\text{PM}}^\# \rightarrow f_{\text{Mod}}^\# a(\mathbb{X}, \text{Mod})$$

is a natural isomorphism. If this is the case, $f_{\text{Mod}}^\#$ has the right adjoint $f_{\natural}^{\text{Mod}}$, and in particular, it preserves arbitrary limits and arbitrary colimits.

Proof. Let $\mathcal{M} \in \text{PM}(\mathbb{X})$ and $y \in \mathbb{Y}$. Then the canonical map

$$\begin{aligned} \check{H}^0(y, f_{\text{PM}}^\# \mathcal{M}) &= \varinjlim \check{H}^0((y_\mu \rightarrow y), f_{\text{PM}}^\# \mathcal{M}) \\ &= \varinjlim \check{H}^0((fy_\mu \rightarrow fy), \mathcal{M}) \rightarrow \check{H}^0(fy, \mathcal{M}) \end{aligned}$$

is an isomorphism, because the coverings of the form $(fy_\mu \rightarrow fy)$ is cofinal in the set of all coverings of fy by assumption. Thus the canonical map

$$\check{H}^0(f_{\text{PM}}^\# \mathcal{M}) \rightarrow f_{\text{PM}}^\# \check{H}^0(\mathcal{M})$$

is an isomorphism. The first assertion follows immediately from the construction of the sheafification, see [2, (II.1)].

Next we show that $f_{\text{Mod}}^\#$ has a right adjoint. To prove this, it suffices to show that $f_b^{\text{PM}}(\mathcal{M})$ is a sheaf if so is $\mathcal{M}$ for $\mathcal{M} \in \text{PM}(\mathbb{X})$. Let $u : \text{Id}_{\text{PM}(\mathbb{Y})} \rightarrow f_{\text{PM}}^\# f_b^{\text{PM}}$ be the unit of adjunction, $\varepsilon : f_b^{\text{PM}} f_{\text{PM}}^\# \rightarrow \text{Id}_{\text{PM}(\mathbb{X})}$ the counit of adjunction, $v(\mathbb{X}) : q(\mathbb{X}, \text{Mod})a(\mathbb{X}, \text{Mod}) \rightarrow \text{Id}_{\text{PM}(\mathbb{X})}$ the unit of adjunction, and $v(\mathbb{Y}) : q(\mathbb{Y}, \text{Mod})a(\mathbb{Y}, \text{Mod}) \rightarrow \text{Id}_{\text{PM}(\mathbb{Y})}$ the unit of adjunction. Then the diagram of functors

$$\begin{array}{ccccccc} f_b^{\text{PM}} & \xrightarrow{uf_b^{\text{PM}}} & f_b^{\text{PM}} f_{\text{PM}}^\# f_b^{\text{PM}} & \xrightarrow{\text{id}} & f_b^{\text{PM}} f_{\text{PM}}^\# f_b^{\text{PM}} & \xrightarrow{f_b^{\text{PM}} \varepsilon} & f_b^{\text{PM}} \\ \downarrow v(\mathbb{X}) f_b^{\text{PM}} & & \downarrow f_b^{\text{PM}} f_{\text{PM}}^\# v(\mathbb{X}) f_b^{\text{PM}} & & \downarrow f_b^{\text{PM}} v(\mathbb{Y}) f_{\text{PM}}^\# f_b^{\text{PM}} & & \downarrow f_b^{\text{PM}} v(\mathbb{Y}) \\ qa f_b^{\text{PM}} & \xrightarrow{uqa f_b^{\text{PM}}} & f_b^{\text{PM}} f_{\text{PM}}^\# qa f_b^{\text{PM}} & \xrightarrow{\cong} & f_b^{\text{PM}} qa f_{\text{PM}}^\# f_b^{\text{PM}} & \xrightarrow{f_b^{\text{PM}} qa \varepsilon} & f_b^{\text{PM}} qa \end{array}$$

is commutative, where $\cong$ is the *inverse* of the canonical map caused by $qa f_{\text{PM}}^\# \rightarrow f_{\text{PM}}^\# qa$, which exists by the first part. As $(f_{\text{PM}}^\#, f_b^{\text{PM}})$ is an adjoint pair, the composite of the first row of the diagram is the identity. As $v(\mathbb{Y})(\mathcal{M})$ is an isomorphism, the right-most vertical arrow evaluated at $\mathcal{M}$ is an isomorphism. Hence, $v(\mathbb{X}) f_b^{\text{PM}}(\mathcal{M})$, which is the left-most vertical arrow evaluated at $\mathcal{M}$, is a split monomorphism. As it is a direct summand of a sheaf, $f_b^{\text{PM}}(\mathcal{M})$ is a sheaf, as desired.

As it is a right adjoint of $f_{\text{Mod}}^{\text{Mod}}$, the functor $f_{\text{Mod}}^\#$ preserves arbitrary limits. As it is a left adjoint of f_b^{Mod} , the functor $f_{\text{Mod}}^\#$ preserves arbitrary colimits. $\square$

(3.19) Let $\mathbb{X}$ be a ringed site, and $x \in \mathbb{X}$. It is easy to see that $\Phi_x : \mathbb{X}/x \rightarrow \mathbb{X}$ satisfies the condition in Lemma 3.18. So $(?)|_x^{\text{Mod}}$ preserves arbitrary limits and colimits. In particular, $(?)|_x^{\text{Mod}}$ is exact. Similarly, for a morphism $\phi : x \rightarrow y$ in $\mathbb{X}$, ϕ_{Mod}^* preserves arbitrary limits and colimits.

4 Derived category and derived functors of sheaves on ringed sites

We utilize the notation and terminology on triangulated categories in [39]. However, We usually write the suspension (translation) functor of a triangulated category by Σ or $(?)[1]$.

Let $\mathcal{T}$ be a triangulated category.

4.1 Lemma. *Let*

$$(a_\lambda \xrightarrow{f_\lambda} b_\lambda \xrightarrow{g_\lambda} c_\lambda \xrightarrow{h_\lambda} \Sigma a_\lambda)$$

be a small family of distinguished triangles in $\mathcal{T}$. Assume that the coproducts $\bigoplus a_\lambda$, $\bigoplus b_\lambda$, and $\bigoplus c_\lambda$ exist. Then the triangle

$$\bigoplus a_\lambda \xrightarrow{\bigoplus f_\lambda} \bigoplus b_\lambda \xrightarrow{\bigoplus g_\lambda} \bigoplus c_\lambda \xrightarrow{H \circ \bigoplus h_\lambda} \Sigma(\bigoplus a_\lambda)$$

is distinguished, where $H : \bigoplus \Sigma a_\lambda \rightarrow \Sigma(\bigoplus a_\lambda)$ is the canonical isomorphism. Similarly, a product of distinguished triangles is a distinguished triangle.

We refer the reader to [31, Proposition 1.2.1] for the proof of the second assertion. The proof for the first assertion is similar [31, Remark 1.2.2].

(4.2) Let $\mathcal{A}$ be an abelian category. The category of bounded below (resp. bounded above, bounded) complexes in $\mathcal{A}$ is denoted by $C^+(\mathcal{A})$ (resp. $C^-(\mathcal{A})$, $C^b(\mathcal{A})$). The corresponding homotopy category and the derived category are denoted by $K^?(\mathcal{A})$ and $D^?(\mathcal{A})$, where $?$ is either $+$, $-$ or b . We denote the homotopy category of complexes in $\mathcal{A}$ with bounded below (resp. bounded above, bounded) cohomology groups by $\bar{K}^?(\mathcal{A})$. The corresponding derived category is denoted by $\bar{D}^?(\mathcal{A})$.

For a plump subcategory $\mathcal{A}'$ of $\mathcal{A}$, we denote by $K_{\mathcal{A}'}^?(\mathcal{A})$ (resp. $\bar{K}_{\mathcal{A}'}^?(\mathcal{A})$) the full subcategory of $K^?(\mathcal{A})$ (resp. $\bar{K}^?(\mathcal{A})$) consisting of complexes whose cohomology groups are objects of $\mathcal{A}'$. The localization of $K_{\mathcal{A}'}^?(\mathcal{A})$ by the épaisse subcategory of exact complexes is denoted by $D_{\mathcal{A}'}^?(\mathcal{A})$. The category $\bar{D}_{\mathcal{A}'}^?(\mathcal{A})$ is defined similarly. Note that the canonical functor $D_{\mathcal{A}'}^?(\mathcal{A}) \rightarrow D(\mathcal{A})$ is fully faithful, and hence $D_{\mathcal{A}'}^?(\mathcal{A})$ is identified with the full subcategory of $D(\mathcal{A})$ consisting of bounded below (resp. bounded above, bounded) complexes whose cohomology groups are in $\mathcal{A}'$. Note also that the canonical functor $D_{\mathcal{A}'}^?(\mathcal{A}) \rightarrow \bar{D}_{\mathcal{A}'}^?(\mathcal{A})$ is an equivalence.

(4.3) Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories, and $F : K^?(\mathcal{A}) \rightarrow K^*(\mathcal{B})$ a triangulated functor. Let $\mathcal{C}$ be a triangulated subcategory of $K^?(\mathcal{A})$ such that

1 If $c \in \mathcal{C}$ is exact, then Fc is exact.

2 For any $a \in K^?(\mathcal{A})$, there exists some quasi-isomorphism $a \rightarrow c$.

The condition **2** implies that the canonical functor

$$i(\mathcal{C}) : \mathcal{C}/(\mathcal{E} \cap \mathcal{C}) \rightarrow K^?(\mathcal{A})/\mathcal{E} = D^?(\mathcal{A})$$

is an equivalence, where $\mathcal{E}$ denotes the épaisse subcategory of exact complexes in $K^?(\mathcal{A})$, see [39, (2.3)]. We fix a quasi-inverse $p(\mathcal{C}) : D^?(\mathcal{A}) \rightarrow \mathcal{C}/(\mathcal{E} \cap \mathcal{C})$. On the other hand, the composite

$$\mathcal{C} \hookrightarrow K^?(\mathcal{A}) \xrightarrow{F} K^*(\mathcal{B}) \xrightarrow{Q} D^*(\mathcal{B}) \quad (4.4)$$

is factorized as

$$\mathcal{C} \xrightarrow{Q_{\mathcal{C}}} \mathcal{C}/(\mathcal{E} \cap \mathcal{C}) \xrightarrow{\Phi} D^*(\mathcal{B}) \quad (4.5)$$

up to a unique natural isomorphism, by the universality of localization and the condition **1** above. Under the setting above, we have the following [16, (I.5.1)].

4.6 Lemma. *The composite functor*

$$RF : D^?(\mathcal{A}) \xrightarrow{p(\mathcal{C})} \mathcal{C}/(\mathcal{E} \cap \mathcal{C}) \xrightarrow{\Phi} D^*(\mathcal{B})$$

is a right derived functor of F .

We denote the map $QF \rightarrow (RF)Q$ corresponding to $\text{id} : RF \rightarrow RF$ by Ξ or $\Xi(F)$.

(4.7) Here we are going to review Spaltenstein's work on unbounded derived categories [34].

A chain complex I of $\mathcal{A}$ is called K -injective if for any exact sequence E of $\mathcal{A}$, the complex of abelian groups $\text{Hom}_{\mathcal{A}}^{\bullet}(E, I)$ is also exact.

A chain map $f : C \rightarrow I$ is called a K -injective resolution of C , if I is K -injective and f is a quasi-isomorphism.

A special case of the following was proved in [34] (the same proof works for the general case without much modification). See also [1].

4.8 Lemma. *If $\mathcal{A}$ is Grothendieck, then for any chain complex $\mathbb{F} \in K(\mathcal{A})$ there is a K -injective resolution $\mathbb{F} \rightarrow \mathbb{I}$ of $\mathbb{F}$ such that each term of $\mathbb{I}$ being an injective object of $\mathcal{A}$.*

Let us denote the homotopy category of chain complexes of $\mathcal{A}$ by $K(\mathcal{A})$. A chain complex I is K -injective if and only if $K(\mathcal{A})(E, I) = 0$ for any exact sequence E . It is easy to see that the K -injective complexes form an épaisse subcategory $I(\mathcal{A})$ of $K(\mathcal{A})$.

(4.9) Let $F : K(\mathcal{A}) \rightarrow K(\mathcal{B})$ be a triangulated functor, and assume that $\mathcal{A}$ is Grothendieck. Let $\mathcal{I}$ be the full subcategory of K -injective complexes of $K(\mathcal{A})$. It is easy to see that $\mathcal{I}$ is triangulated, and $\mathcal{I} \cap \mathcal{E} = 0$. By (4.6), the composite

$$D(\mathcal{A}) \xrightarrow{p(\mathcal{I})} \mathcal{I} \xrightarrow{\Phi} D(\mathcal{B})$$

is a right derived functor RF of F . Note that to fix $p(\mathcal{I})$ is nothing but to fix a functorial K -injective resolution $\mathbb{F} \rightarrow pQ\mathbb{F} = \mathbb{I}_{\mathbb{F}}$.

(4.10) Let $F : K(\mathcal{A}) \rightarrow K(\mathcal{B})$ be a triangulated functor, and assume that there is a right derived functor of F . For $\mathbb{F} \in K(\mathcal{A})$, we say that $\mathbb{F}$ is (right) F -acyclic if the canonical map $QF\mathbb{F} \rightarrow (RF)Q\mathbb{F}$ is an isomorphism.

4.11 Lemma. *Let $\mathcal{A}$ and $\mathcal{B}$ be abelian categories, and $F : \mathcal{A} \rightarrow \mathcal{B}$ be an exact functor with the right adjoint G . Assume that $\mathcal{B}$ is Grothendieck. Then $KG : K(\mathcal{B}) \rightarrow K(\mathcal{A})$ preserves K -injective complexes. Moreover, $RG : D(\mathcal{B}) \rightarrow D(\mathcal{A})$ is the right adjoint of $RF = F$.*

Proof. Let $\mathbb{M} \in K(\mathcal{A})$, and $\mathbb{I}$ a K -injective complex of $K(\mathcal{B})$. Then

$$\begin{aligned} \mathrm{Hom}_{K(\mathcal{A})}(\mathbb{M}, (KG)\mathbb{I}) &\cong H^0(\mathrm{Hom}_{\mathcal{A}}^{\bullet}(\mathbb{M}, G\mathbb{I})) \\ &\cong H^0(\mathrm{Hom}_{\mathcal{B}}^{\bullet}(F\mathbb{M}, \mathbb{I})) = \mathrm{Hom}_{K(\mathcal{B})}(F\mathbb{M}, \mathbb{I}). \end{aligned}$$

If $\mathbb{M}$ is exact, then the last complex is zero. This shows $(KG)\mathbb{I}$ is K -injective.

Now let $\mathbb{M} \in D(\mathcal{A})$ and $\mathbb{N} \in D(\mathcal{B})$ be arbitrary. Then by the first part, we have a functorial isomorphism

$$\begin{aligned} \mathrm{Hom}_{D(\mathcal{A})}(\mathbb{M}, (RG)\mathbb{N}) &\cong \mathrm{Hom}_{K(\mathcal{A})}(\mathbb{M}, (KG)\mathbb{I}_{\mathbb{N}}) \\ &\cong \mathrm{Hom}_{K(\mathcal{B})}(F\mathbb{M}, \mathbb{I}_{\mathbb{N}}) \cong \mathrm{Hom}_{D(\mathcal{B})}(F\mathbb{M}, \mathbb{N}). \end{aligned}$$

This proves the last assertion. $\square$

4.12 Remark. Note that for an abelian category $\mathcal{A}$, we have $\text{ob}(C(\mathcal{A})) = \text{ob}(K(\mathcal{A})) = \text{ob}(D(\mathcal{A}))$. Thus, an object of one of the three categories is sometimes viewed as an object of another.

(4.13) Let $\mathcal{A}$ be a closed symmetric monoidal abelian category which satisfies the (AB3) and (AB3*) conditions. Let $\otimes$ be the multiplication and $[?, ?]$ be the internal hom. For a fixed $b \in \mathcal{A}$, $(? \otimes b, [b, ?])$ is an adjoint pair. In particular, $? \otimes b$ preserves colimits, and $[b, ?]$ preserves limits. By symmetry, $a \otimes ?$ also preserves colimits. As we have an isomorphism

$$\mathcal{A}(a, [b, c]) \cong \mathcal{A}(a \otimes b, c) \cong \mathcal{A}(b \otimes a, c) \cong \mathcal{A}(b, [a, c]) \cong \mathcal{A}^{\text{op}}([a, c], b),$$

we have $[?, c] : \mathcal{A}^{\text{op}} \rightarrow \mathcal{A}$ is right adjoint to $[?, c] : \mathcal{A} \rightarrow \mathcal{A}^{\text{op}}$. This shows that $[?, c]$ changes colimits to limits.

As in [15], we define the tensor product $\mathbb{F} \otimes^\bullet \mathbb{G}$ of $\mathbb{F}, \mathbb{G} \in C(\mathcal{A})$ by

$$(\mathbb{F} \otimes^\bullet \mathbb{G})^n := \bigoplus_{p+q=n} \mathbb{F}^p \otimes \mathbb{G}^q$$

and

$$d^n = d_{\mathbb{F}} \otimes 1 + (-1)^n 1 \otimes d_{\mathbb{G}}.$$

We have $\mathbb{F} \otimes^\bullet \mathbb{G} \in C(\mathcal{A})$. Similarly, $[\mathbb{F}, \mathbb{G}]^\bullet \in C(\mathcal{A})$ is defined by

$$[\mathbb{F}, \mathbb{G}]^n := \prod_{p \in \mathbb{Z}} [\mathbb{F}^p, \mathbb{G}^{n+p}]$$

and

$$d^n := [d_{\mathbb{F}}, 1] + (-1)^{n+1} [1, d_{\mathbb{G}}].$$

It is straightforward to prove the following.

4.14 Lemma. *Let $\mathcal{A}$ be as above. Then the category of chain complexes $C(\mathcal{A})$ is closed symmetric monoidal with $\otimes^\bullet$ the multiplication and $[?, ?]^\bullet$ the internal hom. The bi-triangulated functors*

$$\otimes^\bullet : K(\mathcal{A}) \times K(\mathcal{A}) \rightarrow K(\mathcal{A})$$

and

$$[?, ?]^\bullet : K(\mathcal{A})^{\text{op}} \times K(\mathcal{A}) \rightarrow K(\mathcal{A}),$$

are induced, and $K(\mathcal{A})$ is a closed symmetric monoidal triangulated category (see [23, (3.5), (3.6)]).

(4.15) Let $\mathcal{A}$ be an abelian category, and $\mathfrak{P}$ a full subcategory of $C(\mathcal{A})$. An inverse system $(\mathbb{F}_i)_{i \in I}$ in $C(\mathcal{A})$ is said to be $\mathfrak{P}$ -special if the following conditions are satisfied.

i I is well-ordered.

ii If $i \in I$ has no predecessor, then the canonical map $I_i \rightarrow \varprojlim I_{j < i}$ is an isomorphism (in particular, $I_{i_0} = 0$ if i_0 is the minimum element of I).

iii If $i \in I$ has a predecessor $i - 1$, then the natural chain map $I_i \rightarrow I_{i-1}$ is an epimorphism, the kernel C_i is isomorphic to some object of $\mathfrak{P}$, and the exact sequence

$$0 \rightarrow C_i \rightarrow I_i \rightarrow I_{i-1} \rightarrow 0$$

is semi-split.

Similarly, $\mathfrak{P}$ -special direct systems are also defined, see [34].

The full subcategory of $C(\mathcal{A})$ consisting of inverse (resp. direct) limits of $\mathfrak{P}$ -special inverse (resp. direct) systems is denoted by $\underline{\mathfrak{P}}$ (resp. $\overline{\mathfrak{P}}$).

(4.16) Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. Various definitions and results on unbounded complexes of sheaves over a ringed space by Spaltenstein [34] is generalized to those for ringed sites. However, note that we can not utilize the notion related to closed subsets, points, or stalks of sheaves.

(4.17) We say that a complex $\mathbb{F} \in C(\text{Mod}(\mathbb{X}))$ is K -flat if $\mathbb{G} \otimes^{\bullet} \mathbb{F}$ is exact whenever $\mathbb{G}$ is an exact complex in $\text{Mod}(\mathbb{X})$. We say that $\mathbb{A} \in C(\text{Mod}(\mathbb{X}))$ is *weakly K -injective* if $\mathbb{A}$ is $\text{Hom}_{\text{Mod}(\mathbb{X})}^{\bullet}(\mathbb{F}, ?)$ -acyclic for any K -flat complex $\mathbb{F}$.

(4.18) Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. For $x \in \mathbb{X}$, we define $\mathcal{O}_x^p$ to be $L_x^{\text{PM}}(\mathcal{O}_{\mathbb{X}}|_x)$, and $\mathcal{O}_x := L_x^{\text{Mod}}(\mathcal{O}_{\mathbb{X}}|_x) = a\mathcal{O}_x^p$. For $x_1, \dots, x_r \in \mathbb{X}$, we define

$$\mathcal{O}_{x_1, \dots, x_r} := \mathcal{O}_{x_1} \otimes \cdots \otimes \mathcal{O}_{x_r}.$$

We denote by $\mathfrak{P}_0 = \mathfrak{P}_0(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ (resp. $\tilde{\mathfrak{P}}_0 = \tilde{\mathfrak{P}}_0(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$) the full subcategory of $C(\text{Mod}(\mathbb{X}))$ consisting of complexes of the form $\mathcal{O}_x[n]$ with $x \in \mathbb{X}$ (resp. $\mathcal{O}_{x_1, \dots, x_r}[n]$ with $x_1, \dots, x_r \in \mathbb{X}$) and $n \in \mathbb{Z}$. We define $\mathfrak{P} = \mathfrak{P}(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ to be $\underline{\mathfrak{P}}_0$ and $\tilde{\mathfrak{P}} := \tilde{\mathfrak{P}}_0$. We also define $\mathfrak{Q}$ (resp. $\tilde{\mathfrak{Q}}$) to be the full subcategory of $C(\text{Mod}(\mathbb{X}))$ consisting of bounded above complexes whose terms are direct

sums of copies of $\mathcal{O}_x$ (resp. $\mathcal{O}_{x_1, \dots, x_r}$). Note that $\tilde{\mathfrak{P}}_0$, $\tilde{\mathfrak{P}}$, and $\tilde{\mathfrak{Q}}$ are closed under tensor products.

A complex in $\tilde{\mathfrak{P}}$ is said to be *strongly K -flat*. We say that $\mathbb{A} \in C(\text{Mod}(\mathbb{X}))$ is K -limp (resp. strongly K -limp) if $\mathbb{A}$ is $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{F}, ?)$ -acyclic for any $\mathbb{F} \in \tilde{\mathfrak{P}}$ (resp. $\mathbb{F} \in \tilde{\mathfrak{P}}$).

4.19 Lemma. *Let $f : (\mathbb{Y}, \mathcal{O}_{\mathbb{Y}}) \rightarrow (\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed continuous functor. Then we have an isomorphism*

$$f_{\#}^{\text{Mod}}(\mathcal{O}_{y_1, \dots, y_r}) \cong \mathcal{O}_{f y_1, \dots, f y_r}$$

for $y_1, \dots, y_r \in \mathbb{Y}$. In particular, if $\mathbb{F} \in \tilde{\mathfrak{P}}(\mathbb{Y})$, then $f_{\#}\mathbb{F} \in \tilde{\mathfrak{P}}(\mathbb{X})$. If $\mathbb{F}$ is strongly K -flat, then so is $f_{\#}\mathbb{F}$.

Proof. Clearly, we may assume that $r = 1$. For $y \in \mathbb{Y}$, we denote the canonical continuous ringed functor

$$(\mathbb{Y}/y, \mathcal{O}_{\mathbb{Y}|y}) \rightarrow (\mathbb{X}/fy, \mathcal{O}_{\mathbb{X}|fy})$$

by f/y . We have $\Phi_{fy} \circ (f/y) = f \circ \Phi_y$. Hence,

$$f_{\#}\mathcal{O}_y = f_{\#}L_y(\mathcal{O}_{\mathbb{Y}|y}) \cong L_{fy}(f/y)_{\#}(\mathcal{O}_{\mathbb{Y}|y}) \cong L_{fy}(\mathcal{O}_{\mathbb{X}|fy}) = \mathcal{O}_{fy}.$$

The last two assertions are obvious now. □

4.20 Lemma. *Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site, and $\mathbb{F}, \mathbb{G} \in C(\text{Mod}(\mathbb{X}))$. Then the following hold:*

- 1 $\mathbb{F}$ is K -flat if and only if $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{F}, \mathbb{I})$ is K -injective for any K -injective complex $\mathbb{I}$.
- 2 If $\mathbb{F}$ is K -flat exact, then $\mathbb{G} \otimes_{\mathcal{O}_{\mathbb{X}}}^\bullet \mathbb{F}$ is exact.
- 3 The inductive limit of a pseudo-filtered inductive system of K -flat complexes is again K -flat.
- 4 The tensor product of two K -flat complexes is again K -flat.

See [34] for the proof.

4.21 Proposition. *Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site, and $x \in \mathbb{X}$. Then $\mathcal{O}_x$ is K -flat.*

Proof. It suffices to show that for any exact complex $\mathcal{E}$, $\mathcal{E} \otimes \mathcal{O}_x$ is exact. To verify this, it suffices to show that for any K -injective complex $\mathcal{I}$, the complex $\mathrm{Hom}_{\mathcal{O}_x}^\bullet(\mathcal{E} \otimes \mathcal{O}_x, \mathcal{I})$ is exact. Indeed, then if we consider the K -injective resolution $\mathcal{E} \otimes \mathcal{O}_x \rightarrow \mathcal{I}$, it must be null-homotopic and thus $\mathcal{E} \otimes \mathcal{O}_x$ must be exact.

Note that we have

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_x}^\bullet(\mathcal{E} \otimes \mathcal{O}_x, \mathcal{I}) &\cong \mathrm{Hom}_{\mathcal{O}_x}^\bullet(\mathcal{O}_x, \underline{\mathrm{Hom}}_{\mathcal{O}_x}^\bullet(\mathcal{E}, \mathcal{I})) \cong \\ &\mathrm{Hom}_{\mathrm{Mod}(\mathbb{X}/x)}^\bullet(\mathcal{E}|_x, \mathcal{I}|_x) \cong \mathrm{Hom}_{\mathcal{O}_x}^\bullet(L_x(\mathcal{E}|_x), \mathcal{I}). \end{aligned}$$

As $(?)|_x$ and L_x are exact (3.19), (3.12), the last complex is exact, and we are done. $\square$

4.22 Corollary. *A strongly K -flat complex is K -flat.*

Proof. Follows from the proposition and Lemma 4.20. $\square$

(4.23) Let $\mathcal{A}$ be an abelian category. The category of complexes of objects of $\mathcal{A}$ is denoted by $C(\mathcal{A})$. For an object

$$\mathbb{F} : \dots \rightarrow F^n \xrightarrow{d^n} F^{n+1} \xrightarrow{d^{n+1}} \dots$$

in $C(\mathcal{A})$, we denote the truncated complex

$$0 \rightarrow \mathrm{Im} \, d^{n-1} \rightarrow F^n \xrightarrow{d^n} F^{n+1} \xrightarrow{d^{n+1}} \dots$$

by $\tau^{\geq n} \mathbb{F}$. Similarly, the truncated complex

$$\dots \rightarrow F^{n-2} \xrightarrow{d^{n-2}} F^{n-1} \xrightarrow{d^{n-1}} F^n \rightarrow \mathrm{Im} \, d^n \rightarrow 0$$

is denoted by $\tau^{\leq n} \mathbb{F}$.

4.24 Lemma. *Let $(\mathbb{X}, \mathcal{O}_\mathbb{X})$ be a ringed site and $\mathbb{F} \in C(\mathrm{AB}(\mathbb{X}))$.*

- 1 *We have $\mathfrak{Q} \subset \mathfrak{P} \subset \tilde{\mathfrak{P}}$ and $\mathfrak{Q} \subset \tilde{\mathfrak{Q}} \subset \tilde{\mathfrak{P}}$.*
- 2 *A K -injective complex is weakly K -injective, a weakly K -injective complex is strongly K -limp, and a strongly K -limp complex is K -limp.*
- 3 *A tensor product of two strongly K -flat complexes is again strongly K -flat.*

4 For any $\mathbb{H} \in C(\text{Mod}(\mathbb{X}))$, there is a Ω -special direct system $\mathbb{F}_n$ and a direct system of chain maps $(f_n : \mathbb{F}_n \rightarrow \tau^{\leq n} \mathbb{H})$ such that f_n is a quasi-isomorphism for each $n \in \mathbb{N}$, and $(\mathbb{F}_n)_l = 0$ for $l \geq n + 2$. We have $\varinjlim \mathbb{F}_n \rightarrow \mathbb{H}$ is a quasi-isomorphism, and $\varinjlim \mathbb{F}_n \in \mathfrak{P}$. Moreover, $Q\mathbb{H}$ is the homotopy colimit of the inductive system $(\tau^{\leq n} Q\mathbb{H})$ in the category $D(\text{Mod}(\mathbb{X}))$.

5 The following are equivalent.

- i** $\mathbb{F}$ is K -limp.
- ii** $\mathbb{F}$ is K -limp as a complex of abelian sheaves.
- iii** $\mathbb{F}$ is $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathcal{O}_x, ?)$ -acyclic for $x \in \mathbb{X}$.
- iv** $\mathbb{F}$ is $\Gamma(x, ?)$ -acyclic for $x \in \mathbb{X}$.
- v** If $\mathbb{G} \in \mathfrak{P}$ and $\mathbb{G}$ is exact, then $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is exact.

6 The following are equivalent.

- i** $\mathbb{F}$ is strongly K -limp.
- ii** $\mathbb{F}$ is strongly K -limp as a complex of abelian sheaves.
- iii** $\mathbb{F}$ is $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathcal{O}_{x_1, \dots, x_r}, ?)$ -acyclic for $x_1, \dots, x_r \in \mathbb{X}$.
- iv** If $\mathbb{G} \in \tilde{\mathfrak{P}}$ and $\mathbb{G}$ is exact, then $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is exact.
- v** For any $\mathbb{G} \in \tilde{\mathfrak{P}}$, $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is strongly K -limp.

7 The following are equivalent.

- i** $\mathbb{F}$ is weakly K -injective.
- ii** If $\mathbb{G}$ is K -flat exact, then $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is exact.

Proof. **1** and **3** are trivial. **2** follows from the definition and Corollary 4.22. The proof of **4** and **5** are left to the reader, see [34, (5.6), (5.16), (5.17), (5.21)].

6 Applying Lemma 4.19 to the canonical functor

$$(\mathbb{X}, a\mathbb{Z}) \rightarrow (\mathbb{X}, \mathcal{O}_{\mathbb{X}}),$$

we immediately have that **i**, **ii**, and **iii** are equivalent.

It is also trivial that these conditions imply **iv**. We prove **iv** $\Rightarrow$ **iii**. Note that by **1** and **4**, we have that $\mathbb{F}$ is K -limp. Let $\xi : \mathbb{F} \rightarrow \mathbb{I}$ be a K -injective resolution, and let $\mathbb{J}$ be the mapping cone of ξ . Take a $\mathfrak{P}$ -resolution $\eta : \mathbb{H} \rightarrow \mathcal{O}_{x_1, \dots, x_r}$, and let $\mathbb{G}$ be the mapping cone of η . By assumption, $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is exact. As $\mathbb{I}$ is K -injective, we have $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{I})$ is also exact. It follows that $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{J})$ is exact. As $\mathbb{J}$ is K -limp exact and $\mathbb{H} \in \mathfrak{P}$, we have $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{H}, \mathbb{J})$ is exact. It follows that $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathcal{O}_{x_1, \dots, x_r}, \mathbb{J})$ is exact. This is what we want to prove.

We prove **i** $\Rightarrow$ **v**. Let $\mathbb{H}$ be an exact complex in $\tilde{\mathfrak{P}}$. Then $\mathbb{H} \otimes^\bullet \mathbb{G}$ is exact by Lemma 4.20 and belongs to $\tilde{\mathfrak{P}}$, since $\tilde{\mathfrak{P}}$ is closed under tensor products. Hence,

$$\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{H}, \underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})) \cong \text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{H} \otimes^\bullet \mathbb{G}, \mathbb{F})$$

is exact. By **iv** $\Rightarrow$ **i**, $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{F})$ is strongly K -limp, as desired.

The implication **v** $\Rightarrow$ **i** is trivial, letting $\mathbb{G} = \mathcal{O}_{\mathbb{X}}$.

7 is proved similarly to **6**. □

4.25 Corollary. *Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site and $\mathbb{F}, \mathbb{G} \in C(\text{Mod}(\mathbb{X}))$.*

- 1** *If $\mathbb{F}$ is strongly K -limp and $\mathbb{G} \in \tilde{\mathfrak{P}}$, then $\mathbb{F}$ is $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, ?)$ -acyclic.*
- 2** *If $\mathbb{F}$ is weakly K -injective and $\mathbb{G}$ is K -flat, then $\mathbb{F}$ is $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, ?)$ -acyclic.*

Proof. **1** Let $\mathbb{F} \rightarrow \mathbb{I}$ be the K -injective resolution, and $\mathbb{J}$ the mapping cone. Let $\rho : \mathbb{H} \rightarrow \underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{J})$ be a $\mathfrak{P}$ -resolution. As $\mathbb{H} \otimes^\bullet \mathbb{G}$ lies in $\tilde{\mathfrak{P}}$ and $\mathbb{J}$ is strongly K -limp exact,

$$\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{H}, \underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{J})) \cong \text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{H} \otimes^\bullet \mathbb{G}, \mathbb{J})$$

is exact. So ρ must be null-homotopic, and hence $\underline{\text{Hom}}_{\text{Mod}(\mathbb{X})}^\bullet(\mathbb{G}, \mathbb{J})$ is exact. This is what we wanted to prove.

2 is similar, and we omit it. □

(4.26) Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. For $\mathbb{G} \in C(\text{Mod}(\mathbb{X}))$, it is easy to see that $\mathbb{G} \otimes_{\mathcal{O}_{\mathbb{X}}}^\bullet ?$ induces a functor from $K(\text{Mod}(\mathbb{X}))$ to itself. By the lemma and [15, Theorem I.5.1], the derived functor $L(\mathbb{G} \otimes_{\mathcal{O}_{\mathbb{X}}}^\bullet ?)$ is induced, and it is calculated using any K -flat resolution of $?$. If we fix $?$, then $L(\mathbb{G} \otimes_{\mathcal{O}_{\mathbb{X}}}^\bullet ?)$ is a functor on $\mathbb{G}$, and it induces a bifunctor

$$* \otimes_{\mathcal{O}_{\mathbb{X}}}^{\bullet, L} ? : D(\text{Mod}(\mathbb{X})) \times D(\text{Mod}(\mathbb{X})) \rightarrow D(\text{Mod}(\mathbb{X})).$$

$\mathbb{G} \otimes_{\mathcal{O}_{\mathbb{X}}}^{\bullet, L} \mathbb{F}$ is calculated using any K -flat resolution of $\mathbb{F}$ or any K -flat resolution of $\mathbb{G}$. Note that $\otimes_{\mathcal{O}_{\mathbb{X}}}^{\bullet, L}$ is a $\triangle$ -functor as in [23, (2.5.7)].

We define the hyperTor functor as follows:

$$\underline{\mathrm{Tor}}_i^{\mathcal{O}_{\mathbb{X}}}(\mathbb{F}, \mathbb{G}) := H^{-i}(\mathbb{F} \otimes_{\mathcal{O}_{\mathbb{X}}}^{\bullet, L} \mathbb{G}).$$

(4.27) Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. For $\mathbb{F} \in C(\mathrm{Mod}(\mathbb{X}))$, the functor $\underline{\mathrm{Hom}}_{\mathcal{O}_{\mathbb{X}}}^{\bullet}(\mathbb{F}, ?)$ induces a functor from $K(\mathrm{Mod}(\mathbb{X}))$ to itself. As $\mathrm{Mod}(\mathbb{X})$ is Grothendieck, we can take K -injective resolutions, and hence the right derived functor $R\underline{\mathrm{Hom}}_{\mathcal{O}_{\mathbb{X}}}^{\bullet}(\mathbb{F}, ?)$ is induced. Thus a bifunctor

$$R\underline{\mathrm{Hom}}_{\mathcal{O}_{\mathbb{X}}}(*, ?) : D(\mathrm{Mod}(\mathbb{X}))^{\mathrm{op}} \times D(\mathrm{Mod}(\mathbb{X})) \rightarrow D(\mathrm{Mod}(\mathbb{X}))$$

is induced. For $\mathbb{F}, \mathbb{G} \in D(\mathrm{Mod}(\mathbb{X}))$, we define the hyperExt sheaf of $\mathbb{F}$ and $\mathbb{G}$ by

$$\underline{\mathrm{Ext}}_i^{\mathcal{O}_{\mathbb{X}}}(\mathbb{F}, \mathbb{G}) := H^i(R\underline{\mathrm{Hom}}_{\mathcal{O}_{\mathbb{X}}}^{\bullet}(\mathbb{F}, \mathbb{G})).$$

Similarly, the functor $\mathrm{Hom}_{\mathcal{O}_{\mathbb{X}}}^{\bullet}(*, ?)$ induces

$$R\mathrm{Hom}_{\mathcal{O}_{\mathbb{X}}}(*, ?) : D(\mathrm{Mod}(\mathbb{X}))^{\mathrm{op}} \times D(\mathrm{Mod}(\mathbb{X})) \rightarrow D(\underline{\mathrm{Ab}}).$$

Almost by definition, we have

$$H^i(R\mathrm{Hom}_{\mathcal{O}_{\mathbb{X}}}(\mathbb{F}, \mathbb{G})) \cong \mathrm{Hom}_{D(\mathrm{Mod}(\mathbb{X}))}(\mathbb{F}, \mathbb{G}[i]).$$

Sometimes we denote these groups by $\mathrm{Ext}_{\mathcal{O}_{\mathbb{X}}}^i(\mathbb{F}, \mathbb{G})$.

4.28 Lemma. *Let $(\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be a ringed site. Then $D(\mathrm{Mod}(\mathbb{X}))$ is a closed symmetric monoidal triangulated category with $\otimes_{\mathcal{O}_{\mathbb{X}}}^{\bullet, L}$ its product and $R\underline{\mathrm{Hom}}_{\mathcal{O}_{\mathbb{X}}}^{\bullet}(*, ?)$ its internal hom.*

Proof. This is straightforward.

4.29 Lemma. *Let $f : (\mathbb{Y}, \mathcal{O}_{\mathbb{Y}}) \rightarrow (\mathbb{X}, \mathcal{O}_{\mathbb{X}})$ be an admissible ringed continuous functor. Then the following hold:*

- 1 *If $\mathbb{I} \in C(\mathrm{AB}(\mathbb{X}))$ is a K -injective (resp. strongly K -limp, K -limp) complex of sheaves of abelian groups, then so is $f_{\mathrm{AB}}^{\#} \mathbb{I}$.*
- 2 *If $\mathbb{F} \in C(\mathrm{Mod}(\mathbb{Y}))$ is strongly K -flat and exact, then $f_{\#}^{\mathrm{Mod}} \mathbb{F}$ is strongly K -flat and exact.*
- 3 *If $\mathbb{I} \in C(\mathrm{AB}(\mathbb{X}))$ is K -limp and exact, then so is $f_{\mathrm{AB}}^{\#} \mathbb{I}$.*

Proof. As $f_{\text{AB}}^\#$ has an exact left adjoint $f_\#^{\text{AB}}$, the assertion for K -injectivity in **1** is obvious.

We prove the assertion for the K -limp property in **1**. Let $\mathbb{P} \in \mathfrak{P}(\mathbb{Y})$ be an exact complex. As $f_\#^{\text{AB}}$ is exact, $f_\#^{\text{AB}}\mathbb{P}$ is exact and a complex in $\mathfrak{P}(\mathbb{X})$ by Lemma 4.19. Hence,

$$\text{Hom}_{\text{AB}(\mathbb{Y})}^\bullet(\mathbb{P}, f_{\text{AB}}^\# \mathbb{I}) \cong \text{Hom}_{\text{AB}(\mathbb{X})}^\bullet(f_\#^{\text{AB}} \mathbb{P}, \mathbb{I})$$

is exact. This shows $f_{\text{AB}}^\# \mathbb{I}$ is K -limp. The proof for the strongly K -limp property is similar.

We prove **2**. We already know that $f_\# \mathbb{F}$ is strongly K -flat by Lemma 4.19. We prove that $\text{Hom}_{\text{Mod}(\mathbb{X})}^\bullet(f_\#^{\text{Mod}} \mathbb{F}, \mathbb{I})$ is exact, where $\eta : f_\#^{\text{Mod}} \mathbb{F} \rightarrow \mathbb{I}$ is a K -injective resolution. Then η must be null-homotopic, and we have $f_\#^{\text{Mod}} \mathbb{F}$ is exact and the proof is complete. Clearly, $\mathbb{I}$ is strongly K -limp, and hence so is $f_{\text{Mod}}^\# \mathbb{I}$ by **1** and Lemma 4.24, **6**. The assertion follows immediately by adjunction.

We prove **3**. Let $\xi : \mathbb{F} \rightarrow f_{\text{AB}}^\# \mathbb{I}$ be a $\mathfrak{P}(\text{AB})$ -resolution of $f_{\text{AB}}^\# \mathbb{I}$. It suffices to show $\text{Hom}_{\text{AB}(\mathbb{Y})}^\bullet(\mathbb{F}, f_{\text{AB}}^\# \mathbb{I})$ is exact, which is obvious since $f_\#^{\text{AB}} \mathbb{F} \in \mathfrak{P}(\text{AB})$. $\square$

(4.30) By the lemma, there is a derived functor $Lf_\#^{\text{Mod}} : D(\text{Mod}(\mathbb{Y})) \rightarrow D(\text{Mod}(\mathbb{X}))$ of $f_\#^{\text{Mod}}$ for an admissible ringed continuous functor f . It is calculated via strongly K -flat resolutions. By the same lemma, we also know that a K -limp complex is $f^\#$ -acyclic, if f is admissible.

Now as in [34, section 6] and [23], the following is proved.

4.31 Lemma. *Let $\mathcal{S}$ be the category of ringed sites and admissible ringed continuous functors. Then $(L(?)_\#^{\text{Mod}}, R(?)_\#^{\text{Mod}})$ is a monoidal adjoint pair of Δ -pseudofunctors on $\mathcal{S}^{\text{op}}$.*

The proof is basically the same as that in [23, Chapter 1–3], and left to the reader. We only remark how the structure maps (i.e., natural transformations) which are necessary to prove the assertion are defined. For example, the unit of adjunction $u : \text{Id} \rightarrow (Rf^\#)(Lf_\#)$ is the unique map which makes the diagram

$$\begin{array}{ccc} Q & \xrightarrow{uQ} & (Rf^\#)(Lf_\#)Q \\ \downarrow Qu & & \downarrow (Rf^\#)\Xi \\ Qf_\#f_\# & \xrightarrow{\Xi f_\#} & (Rf^\#)Qf_\# \end{array}$$

commutative. See [23, Chapter 3] for more.

4.32 Remark. The author does not know if $f_{\#}^{\text{Mod}}$ preserves K -flatness of complexes or exactness of K -flat complexes. This is true if $\mathbb{Y}$ has finite limits and f preserves finite limits. This condition is satisfied by morphisms of ringed spaces. The author does not know if $\mathfrak{P}$ is closed under tensor products. On the other hand, strongly K -flat resolutions can be used to calculate both the derived tensor products and $Lf_{\#}$. The class of strongly K -flat resolutions are closed under tensor products and preserved by $f_{\#}$, and this is enough to prove Lemma 4.31.

5 Sheaves over a diagram of S -schemes

(5.1) Let S be a scheme, and I a small category. We call an object of $\mathcal{P}(I^{\text{op}}, \text{Sch}/S)$ an I -diagram of S -schemes, where Sch/S denotes the category of S -schemes. So an object of $\mathcal{P}(I, \text{Sch}/S)$ is referred as an I^{op} -diagram of S -schemes. Let $X_{\bullet} \in \mathcal{P}(I, \text{Sch}/S)$. We denote $X_{\bullet}(i)$ by X_i for $i \in I$, and $X_{\bullet}(\phi)$ by X_{ϕ} for $\phi \in \text{Mor}(I)$. Let $\mathbb{P}$ be a property of schemes (e.g., quasi-compact, locally noetherian, regular). We say that $X_{\bullet}$ satisfies $\mathbb{P}$ if X_i satisfies $\mathbb{P}$ for any $i \in I$. Let $\mathbb{Q}$ be a property of morphisms of schemes (e.g., quasi-compact, locally of finite type, smooth). We say that $X_{\bullet}$ is $\mathbb{Q}$ over S if the structure map $X_i \rightarrow S$ satisfies $\mathbb{Q}$ for any $i \in I$. We say that $X_{\bullet}$ has $\mathbb{Q}$ arrows if X_{ϕ} satisfies $\mathbb{Q}$ for any $\phi \in \text{Mor}(I)$.

(5.2) Let $f_{\bullet} : X_{\bullet} \rightarrow Y_{\bullet}$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. For $i \in I$, we denote $f_{\bullet}(i) : X_i \rightarrow Y_i$ by f_i . For a property $\mathbb{Q}$ of morphisms of schemes, we say that $f_{\bullet}$ satisfies $\mathbb{Q}$ if so does f_i for any $i \in I$. We say that $f_{\bullet}$ is of *fiber type* if the canonical map $(f_j, X_{\phi}) : X_j \rightarrow Y_j \times_{Y_i} X_i$ is an isomorphism for any morphism $\phi : i \rightarrow j$ of I .

(5.3) Let S , I and $X_{\bullet}$ be as above. We define the *Zariski site* of $X_{\bullet}$, denoted by $\text{Zar}(X_{\bullet})$, as follows. An object of $\text{Zar}(X_{\bullet})$ is a pair (i, U) such that $i \in I$ and U is an open subset of X_i . A morphism $(\phi, h) : (j, V) \rightarrow (i, U)$ is a pair (ϕ, h) such that $\phi \in I(i, j)$ and $h : V \rightarrow U$ is the restriction of X_{ϕ} . For a given morphism $\phi : i \rightarrow j$, U , and V , such an h exists if and only if $V \subset X_{\phi}^{-1}(U)$, and it is unique. We denote this h by $h(\phi; U, V)$. The composition of morphisms is defined in an obvious way. Thus $\text{Zar}(X_{\bullet})$ is a small category. For $(i, U) \in \text{Zar}(X_{\bullet})$, a covering of (i, U) is a family of morphisms of the form

$$((\text{id}_i, h(\text{id}_i; U, U_{\lambda})) : (i, U_{\lambda}) \rightarrow (i, U))_{\lambda \in \Lambda}$$

such that $\bigcup_{\lambda \in \Lambda} U_\lambda = U$. This defines a pretopology of $\text{Zar}(X_\bullet)$, and $\text{Zar}(X_\bullet)$ is a site. As we will consider only the Zariski topology, a presheaf or sheaf on $\text{Zar}(X_\bullet)$ will be sometimes referred as a presheaf or sheaf on $X_\bullet$, if there is no danger of confusion. Thus $\mathcal{P}(X_\bullet, \mathcal{C})$ and $\mathcal{S}(X_\bullet, \mathcal{C})$ mean $\mathcal{P}(\text{Zar}(X_\bullet), \mathcal{C})$ and $\mathcal{S}(\text{Zar}(X_\bullet), \mathcal{C})$, respectively.

(5.4) Let S and I be as above, and let $\sigma : J \hookrightarrow I$ be a subcategory of I . Then we have an obvious restriction functor $\sigma^\# : \mathcal{P}(I, \text{Sch}/S) \rightarrow \mathcal{P}(J, \text{Sch}/S)$, which we denote by $(?)|_J$. If $\text{ob}(J)$ is finite and $I(j, i)$ is finite for each $j \in J$ and $i \in I$, then $(?)|_J$ has a right adjoint functor $\text{cok}_J^I = (?)^{\text{op}} \sigma_\#^{\text{op}} (?)^{\text{op}}$, because Sch/S has finite limits. See [7, pp. 9–12].

(5.5) Let $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Then we have an obvious continuous functor $Q(X_\bullet, J) : \text{Zar}((X_\bullet)|_J) \hookrightarrow \text{Zar}(X_\bullet)$. Note that $Q(X_\bullet, J)$ may not be admissible. The restriction functors $Q(X_\bullet, i)_{\text{AB}}^\#$ and $Q(X_\bullet, J)_{\text{PA}}^\#$ are denoted by $(?)_J^{\text{AB}}$ and $(?)_J^{\text{PA}}$, respectively. For $i \in I$, we consider that i is the subcategory of I whose object set is $\{i\}$ with $\text{Hom}_i(i, i) = \{\text{id}\}$. The restriction $(?)_i^\natural$ is defined.

(5.6) Let $\mathcal{F} \in \text{PA}(X_\bullet)$ and $i \in I$. Then $\mathcal{F}_i \in \text{PA}(X_i)$, and thus we have a family of sheaves $(\mathcal{F}_i)_{i \in I}$. Moreover, for $(i, U) \in \text{Zar}(X_\bullet)$ and $\phi : i \rightarrow j$, we have the restriction map

$$\Gamma(U, \mathcal{F}_i) = \Gamma((i, U), \mathcal{F}) \xrightarrow{\text{res}} \Gamma((j, X_\phi^{-1}(U)), \mathcal{F}) = \Gamma(X_\phi^{-1}(U), \mathcal{F}_j) = \Gamma(U, (X_\phi)_* \mathcal{F}_j),$$

which induces

$$\beta_\phi(\mathcal{F}) \in \text{Hom}_{\text{PA}(X_j)}(\mathcal{F}_i, (X_\phi)_* \mathcal{F}_j). \quad (5.7)$$

The corresponding map in $\text{Hom}_{\text{PA}(X_j)}((X_\phi)_*^{\text{PA}}(\mathcal{F}_i), \mathcal{F}_j)$ is denoted by $\alpha_\phi^{\text{PA}}(\mathcal{F})$. If $\mathcal{F}$ is a sheaf, then (5.7) yields

$$\alpha_\phi^{\text{AB}}(\mathcal{F}) \in \text{Hom}_{\text{AB}(X_j)}((X_\phi)_*^{\text{AB}}(\mathcal{F}_i), \mathcal{F}_j).$$

It is straightforward to check the following.

5.8 Lemma ([7]). *Let $\natural$ be either AB or PA. The following hold:*

- 1 For any $i \in I$, we have $\alpha_{\text{id}_i}^\natural : (X_{\text{id}_i})_*^\natural(\mathcal{F}_i) \rightarrow \mathcal{F}_i$ is the canonical identification.

2 If $\phi \in I(i, j)$ and $\psi \in I(j, k)$, then the composite map

$$(X_{\psi\phi})_{\natural}^*(\mathcal{F}_i) \cong (X_{\psi})_{\natural}^*(X_{\phi})_{\natural}^*(\mathcal{F}_i) \xrightarrow{(X_{\psi})_{\natural}^* \alpha_{\phi}^{\natural}} (X_{\psi})_{\natural}^*(\mathcal{F}_j) \xrightarrow{\alpha_{\psi}^{\natural}} \mathcal{F}_k \quad (5.9)$$

agrees with $\alpha_{\psi\phi}^{\natural}$.

3 Conversely, a family $((\mathcal{G}_i)_{i \in I}, (\alpha_{\phi})_{\phi \in \text{Mor}(I)})$ such that $\mathcal{G}_i \in \natural(X_i)$, $\alpha_{\phi} \in \text{Hom}_{\natural(X_j)}((X_{\phi})_{\natural}^*(\mathcal{G}_i), \mathcal{G}_j)$ for $\phi \in I(i, j)$, and that the conditions corresponding to **1,2** are satisfied yields $\mathcal{G} \in \natural(X_{\bullet})$, and this correspondence gives an equivalence.

Similarly, a family $((\mathcal{G}_i)_{i \in \text{ob}(I)}, (\beta_{\phi})_{\phi \in \text{Mor}(I)})$ with

$$\mathcal{G}_i \in \natural(X_i) \text{ and } \beta_{\phi} \in \text{Hom}_{\natural(X_i)}(\mathcal{G}_i, (X_{\phi})_{\natural}^* \mathcal{G}_j)$$

satisfying the conditions

1' For $i \in \text{ob}(I)$, $\beta_{\text{id}_i} : \mathcal{G}_i \rightarrow (\text{id}_i)_* \mathcal{G}_i$ is the canonical identification;

2' For $\phi \in I(i, j)$ and $\psi \in I(j, k)$, the composite

$$\mathcal{F}_i \xrightarrow{\beta_{\phi}} (X_{\phi})_*(\mathcal{F}_j) \xrightarrow{(X_{\phi})_* \beta_{\psi}} (X_{\phi})_*(X_{\psi})_*(\mathcal{F}_k) \cong (X_{\psi\phi})_*(\mathcal{F}_k)$$

agrees with $\beta_{\psi\phi}$

are in one to one correspondence with $\mathcal{G} \in \natural(X_{\bullet})$.

(5.10) Let $\mathcal{F} \in \text{AB}(X_{\bullet})$. We say that $\mathcal{F}$ is an *equivariant* abelian sheaf if $\alpha_{\phi}^{\text{AB}}$ are isomorphisms for all $\phi \in \text{Mor}(I)$. For $\mathcal{F} \in \text{PA}(X_{\bullet})$, we say that $\mathcal{F}$ is an equivariant abelian presheaf if $\alpha_{\phi}^{\text{PA}}$ are isomorphisms for all $\phi \in \text{Mor}(I)$. An equivariant sheaf may not be an equivariant presheaf. However, an equivariant presheaf which is a sheaf is an equivariant sheaf. We denote the category of equivariant sheaves and presheaves by $\text{EqAB}(X_{\bullet})$ and $\text{EqPA}(X_{\bullet})$, respectively. As $(X_{\phi})_{\natural}^*$ is exact for $\natural = \text{AB}, \text{PA}$ and any ϕ , we have that $\text{EqAB}(X_{\bullet})$ is plump in $\text{AB}(X_{\bullet})$, and $\text{EqPA}(X_{\bullet})$ is plump in $\text{PA}(X_{\bullet})$.

(5.11) Let $X_{\bullet} \in \mathcal{P}(I, \text{Sch}/S)$. The datum

$$((\mathcal{O}_{X_i})_{i \in I}, (\beta_{\phi} : \mathcal{O}_{X_i} \rightarrow (X_{\phi})_* \mathcal{O}_{X_j})_{\phi \in \text{Mor}(I)})$$

gives a sheaf of commutative rings on $X_{\bullet}$, which we denote by $\mathcal{O}_{X_{\bullet}}$, and thus $\text{Zar}(X_{\bullet})$ is a ringed site. The categories $\text{PM}(\text{Zar}(X_{\bullet}))$ and $\text{Mod}(\text{Zar}(X_{\bullet}))$ are denoted by $\text{PM}(X_{\bullet})$ and $\text{Mod}(X_{\bullet})$, respectively.

(5.12) For $J \subset I$, we have $\mathcal{O}_{X_\bullet|_J} = (\mathcal{O}_{X_\bullet})_J$ by definition. The continuous functor

$$Q(X_\bullet, J) : (\text{Zar}(X_\bullet|_J), \mathcal{O}_{X_\bullet|_J}) \rightarrow (\text{Zar}(X_\bullet), \mathcal{O}_{X_\bullet})$$

is actually a ringed continuous functor.

The corresponding restriction $Q(X_\bullet, J)_\natural^\#$ is denoted by $(?)_J^\natural$ for $\natural = \text{PM}, \text{Mod}$. For subcategories $J_1 \subset J \subset I$ of I , we denote the restriction $(?)_{J_1}^\natural : \natural(X_\bullet|_J) \rightarrow \natural(X_\bullet|_{J_1})$ is denoted by $(?)_{J_1, J}^\natural$, to emphasize J .

(5.13) Let $\natural$ be PM or Mod. Note that $\mathcal{M} \in \natural(X_\bullet)$ is nothing but a family

$$\text{Dat}(\mathcal{M}) := ((\mathcal{M}_i)_{i \in I}, (\alpha_\phi^\natural)_{\phi \in \text{Mor}(I)})$$

such that $\mathcal{M}_i \in \natural(X_i)$, $\alpha_\phi^\natural : (X_\phi)_\natural^*(\mathcal{M}_i) \rightarrow \mathcal{M}_j$ is a morphism of $\natural(X_\bullet)$ for any $\phi : i \rightarrow j$, and the conditions corresponding to **1,2** in Lemma 5.8 are satisfied.

We say that $\mathcal{M} \in \natural(X_\bullet)$ is *equivariant* if $\alpha_\phi^\natural$ is an isomorphism for any $\phi \in \text{Mor}(I)$. Note that equivariance depends on $\natural$, and is not preserved by the forgetful functors in general. We denote the full subcategory of $\text{Mod}(X_\bullet)$ consisting of equivariant objects by $\text{EM}(X_\bullet)$.

6 The left and right inductions and the direct and inverse images

Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$.

(6.1) Let J be a subcategory of I . The left adjoint $Q(X_\bullet, J)_\#^\natural$ of $(?)_J^\natural$ (see (5.12)) is denoted by $L_J^\natural$ for $\natural = \text{PA}, \text{AB}, \text{PM}, \text{Mod}$. The right adjoint $Q(X_\bullet, J)_\flat^\natural$ of $(?)_J^\natural$, which exists by Lemma 3.18, is denoted by $R_J^\natural$ for $\natural = \text{PA}, \text{AB}, \text{PM}, \text{Mod}$. We call $L_J^\natural$ and $R_J^\natural$ the left and right induction functor, respectively.

Let $J_1 \subset J \subset I$ be subcategories of I . The left and right adjoints of $(?)_{J_1, J}^\natural$ is denoted by $L_{J, J_1}^\natural$ and $R_{J, J_1}^\natural$, respectively. As $(?)_J^\natural$ has both a left adjoint and a right adjoint, we have

6.2 Lemma. *The functor $(?)_J^\natural$ preserves arbitrary limits and colimits (hence is exact) for $\natural = \text{PA}, \text{AB}, \text{PM}, \text{Mod}$.*

The functor $\natural(X_\bullet) \rightarrow \prod_{i \in I} \natural(X_i)$ given by $\mathcal{F} \mapsto (\mathcal{F}_i)_{i \in I}$ is faithful for $\natural = \text{PA}, \text{AB}, \text{PM}, \text{Mod}$.

(6.3) Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. This induces an obvious morphism

$$f_\bullet^{-1} : (\text{Zar}(Y_\bullet), \mathcal{O}_{Y_\bullet}) \rightarrow (\text{Zar}(X_\bullet), \mathcal{O}_{X_\bullet})$$

of ringed sites. We have $\text{id}^{-1} = \text{id}$, and $(g_\bullet \circ f_\bullet)^{-1} = f_\bullet^{-1} \circ g_\bullet^{-1}$ for $g_\bullet : Y_\bullet \rightarrow Z_\bullet$.

We define the direct image $(f_\bullet)_*^{\mathfrak{h}}$ to be $(f_\bullet^{-1})_\#^{\mathfrak{h}}$, and the inverse image $(f_\bullet)^*_{\mathfrak{h}}$ to be $(f_\bullet^{-1})_{\#}^{\mathfrak{h}}$ for $\mathfrak{h} = \text{Mod}, \text{PM}, \text{AB}, \text{PA}$.

6.4 Lemma. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$, and $K \subset J \subset I$. Then we have

$$1 \quad Q(X_\bullet, J) \circ Q(X_\bullet|_J, K) = Q(X_\bullet, K)$$

$$2 \quad f_\bullet^{-1} \circ Q(Y_\bullet, J) = Q(X_\bullet, J) \circ (f_\bullet|_J)^{-1}.$$

(6.5) Let us fix I and S . By Lemma 3.14, we have various natural maps between functors on sheaves arising from the closed structures and the monoidal pairs, involving various J -diagrams of schemes, where J varies subcategories of I . In the sequel, many of the natural maps are referred as ‘the canonical maps’ or ‘the canonical isomorphisms’ without any explicit definitions. Many of them are defined in [23], and various commutativity theorems are proved there.

6.6 Example. Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y_\bullet \rightarrow Z_\bullet$ are morphisms in $\mathcal{P}(I, \text{Sch}/S)$. Let $K \subset J \subset I$ be subcategories, and $\mathfrak{h}$ denote PM, Mod, PA, or AB.

1 There is a natural isomorphism

$$c_{I,J,K}^{\mathfrak{h}} : (?)_{K,I}^{\mathfrak{h}} \cong (?)_{K,J}^{\mathfrak{h}} \circ (?)_{J,I}^{\mathfrak{h}}.$$

Taking the conjugate,

$$d_{I,J,K}^{\mathfrak{h}} : L_{I,J}^{\mathfrak{h}} \circ L_{J,K}^{\mathfrak{h}} \cong L_{I,K}^{\mathfrak{h}}$$

is induced.

2 There is a natural isomorphism

$$c_{J,f_\bullet}^{\mathfrak{h}} : (?)_J^{\mathfrak{h}} \circ (f_\bullet)_*^{\mathfrak{h}} \cong (f_\bullet|_J)_*^{\mathfrak{h}} \circ (?)_J^{\mathfrak{h}}$$

and its conjugate

$$d_{J,f_\bullet}^{\mathfrak{h}} : L_J^{\mathfrak{h}} \circ (f_\bullet|_J)^*_{\mathfrak{h}} \cong (f_\bullet)^*_{\mathfrak{h}} \circ L_J^{\mathfrak{h}}.$$

3 We have

$$(c_{K,f_\bullet|_J}^\natural(?)_J^\natural) \circ ((?)_{K,J}^\natural c_{J,f}^\natural) = ((f_\bullet|_K)_*^\natural c_{I,J,K}^\natural) \circ c_{K,f_\bullet}^\natural \circ ((c_{I,J,K}^\natural)^{-1}(f_\bullet)_*^\natural).$$

4 We have

$$((g_\bullet|_J)_*^\natural c_{J,f_\bullet}^\natural) \circ (c_{J,g_\bullet}^\natural (f_\bullet)_*^\natural) = (c_{f_\bullet|_J,g_\bullet|_J}^\natural(?)_J^\natural) \circ c_{J,g_\bullet \circ f_\bullet}^\natural \circ ((?)_J^\natural (c_{f_\bullet,g_\bullet}^\natural)^{-1}),$$

where $c_{f_\bullet,g_\bullet}^\natural : (g_\bullet \circ f_\bullet)_*^\natural \cong (g_\bullet)_*^\natural \circ (f_\bullet)_*^\natural$ is the canonical isomorphism, and similarly for $c_{f_\bullet|_J,g_\bullet|_J}^\natural$.

5 The canonical map

$$H : \mathcal{M}_J \otimes_{\mathcal{O}_{X_\bullet|_J}} \mathcal{N}_J \rightarrow (\mathcal{M} \otimes_{\mathcal{O}_{X_\bullet}} \mathcal{N})_J$$

is an isomorphism, as can be seen easily. The canonical map

$$\Delta : L_J(\mathcal{M} \otimes_{\mathcal{O}_{X_\bullet|_J}} \mathcal{N}) \cong (L_J \mathcal{M}) \otimes_{\mathcal{O}_{X_\bullet}} (L_J \mathcal{N}).$$

is defined, which may not be an isomorphism.

7 Operations on sheaves via the structure data

Let I be a small category, S a scheme, and $\mathcal{P} := \mathcal{P}(I, \text{Sch}/S)$. To study sheaves on objects of $\mathcal{P}$, it is convenient to utilize the structure data of them, and then utilize the usual sheaf theory on schemes.

(7.1) Let $X_\bullet \in \mathcal{P}$. Let $\natural$ be any of PA, AB, PM, Mod, and $\mathcal{M}, \mathcal{N} \in \natural(X_\bullet)$. An element (φ_i) in $\prod \text{Hom}_{\natural(X_i)}(\mathcal{M}_i, \mathcal{N}_i)$ is given by some $\varphi \in \text{Hom}_{\natural(X_\bullet)}(\mathcal{M}, \mathcal{N})$ (by the canonical faithful functor $\natural(X_\bullet) \rightarrow \prod \natural(X_i)$), if and only if

$$\varphi_j \circ \alpha_\phi(\mathcal{M}) = \alpha_\phi(\mathcal{N}) \circ (X_\phi)_\natural^*(\varphi_i) \quad (7.2)$$

holds (or equivalently, $\beta_\phi(\mathcal{N}) \circ \varphi_i = (X_\phi)_* \varphi_j \circ \beta_\phi(\mathcal{M})$ holds) for any $(\phi : i \rightarrow j) \in \text{Mor}(I)$.

We say that a family of morphisms $(\varphi_i)_{i \in I}$ between structure data

$$\varphi_i : \mathcal{M}_i \rightarrow \mathcal{N}_i$$

is a morphism of structure data if φ_i is a morphism in $\natural(X_i)$ for each i , and (7.2) is satisfied for any ϕ . Thus the category of structure data of sheaves, presheaves, modules, and premodules on $X_\bullet$, denoted by $\mathfrak{D}_\natural(X_\bullet)$ is defined, and the equivalence $\text{Dat}_\natural : \natural(X_\bullet) \cong \mathfrak{D}_\natural(X_\bullet)$ is given. This is the precise meaning of Lemma 5.8.

(7.3) Let $X_\bullet \in \mathcal{P}$ and $\mathcal{M}, \mathcal{N} \in \text{Mod}(X_\bullet)$. As in Example 6.6, **5**, we have an isomorphism

$$m_i : \mathcal{M}_i \otimes_{\mathcal{O}_{X_i}} \mathcal{N}_i \cong (\mathcal{M} \otimes_{\mathcal{O}_{X_\bullet}} \mathcal{N})_i.$$

This is trivial for presheaves, and utilize the fact the sheafification is compatible with $(?)_i$ for sheaves. Under this identification, the structure map α_ϕ of $\mathcal{M} \otimes \mathcal{N}$ can be completely described via those of $\mathcal{M}$ and $\mathcal{N}$. Namely, for $\phi \in I(i, j)$, $\alpha_\phi(\mathcal{M} \otimes \mathcal{N})$ agrees with the composite map

$$X_\phi^*(\mathcal{M} \otimes \mathcal{N})_i \xrightarrow{\text{via } m_i^{-1}} X_\phi^*(\mathcal{M}_i \otimes \mathcal{N}_i) \cong X_\phi^* \mathcal{M}_i \otimes X_\phi^* \mathcal{N}_i \xrightarrow{\alpha_\phi \otimes \alpha_\phi} \mathcal{M}_j \otimes \mathcal{N}_j \xrightarrow{m_j} (\mathcal{N} \otimes \mathcal{N})_j.$$

This can be checked directly at the presheaf level, and then sheafify them.

(7.4) Let $X_\bullet \in \mathcal{P}$, and J a subcategory of I . The left adjoint functor $L_J^\natural = Q(X_\bullet, J)_\#^\natural$ of $(?)_J^\natural$ is given by the structure data as follows explicitly. For $\mathcal{M} \in \mathfrak{h}(X_\bullet|_J)$ and $i \in I$, we have

7.5 Lemma. *There is an isomorphism*

$$\lambda_{J,i} : (L_J^\natural(\mathcal{M}))_i^\natural \cong \varinjlim (X_\phi)_\natural^*(\mathcal{M}_j),$$

where the colimit is taken over the subcategory $(I_i^{(J^{\text{op}} \rightarrow I^{\text{op}})})^{\text{op}}$ of I/i whose objects are $(\phi : j \rightarrow i) \in I/i$ with $j \in \text{ob}(J)$ and morphisms are morphisms φ of I/i such that $\varphi \in \text{Mor}(J)$.

In particular, we have an isomorphism

$$\lambda_{j,i} : (L_J^\natural(\mathcal{M}))_i^\natural \cong \bigoplus_{\phi \in I(j,i)} (X_\phi)_\natural^*(\mathcal{M}_j). \quad (7.6)$$

(7.7) Let $\psi : i \rightarrow i'$ be a morphism. The structure map

$$\alpha_\psi : (X_\psi)_\natural^*((L_J^\natural(\mathcal{M}))_i^\natural) \rightarrow (L_J^\natural(\mathcal{M}))_{i'}^\natural$$

is induced by

$$(X_\psi)_\natural^*((X_\phi)_\natural^*(\mathcal{M}_j)) \cong (X_{\psi\phi})_\natural^*(\mathcal{M}_j).$$

(7.8) The counit map $\varepsilon : L_J(?)_J \rightarrow \text{Id}$ is given as a morphism of structure data as follows.

$$\varepsilon_i : (?)_i L_J(?)_J \rightarrow (?)_i$$

agrees with

$$(?)_i L_J(?)_J \xrightarrow{\lambda_{J,i}} \varinjlim X_\phi^*(?)_j(?)_J \cong \varinjlim X_\phi^*(?)_j \xrightarrow{\alpha} (?)_i,$$

where α is induced by $\alpha_\phi : X_\phi^*(?)_j \rightarrow (?)_i$.

(7.9) The unit map $u : \text{Id} \rightarrow (?)_J L_J$ is also described, as follows.

$$u_j : (?)_j \rightarrow (?)_j (?)_J L_J$$

agrees with

$$(?)_j \cong \text{id}_j^*(?)_j \rightarrow \varinjlim X_\phi^*(?)_k \xrightarrow{\lambda_{J,j}^{-1}} (?)_j L_J \cong (?)_j (?)_J L_J,$$

where the colimit is taken over $(I_j^{(J^{\text{op}} \subset I^{\text{op}})})^{\text{op}}$.

(7.10) Let $X_\bullet \in \mathcal{P}$, and J a subcategory of I . The right adjoint functor $R_J^\natural$ of $(?)_J^\natural$ is given as follows explicitly. For $\mathcal{M} \in \natural(X_\bullet|_J)$ and $i \in I$, we have

$$\rho^{J,i} : (R_J^\natural(\mathcal{M}))_i^\natural = \varprojlim (X_\phi)_*^\natural(\mathcal{M}_j),$$

where the limit is taken over $I_i^{(J \rightarrow I)}$, see (3.4) for the notation. The description of α , u , and ε for the right induction are left to the reader.

7.11 Lemma. *Let $X_\bullet \in \mathcal{P}$, and J a full subcategory of I . Then we have the following.*

- 1** *The counit of adjunction $\varepsilon : (?)_J^\natural \circ R_J^\natural \rightarrow \text{Id}$ is an isomorphism.*
- 2** *The unit of adjunction $u : \text{Id} \rightarrow (?)_J^\natural \circ L_J^\natural$ is an isomorphism.*

Proof. **1** For $i \in J$, the restriction

$$\varepsilon_i : (?)_i^\natural (?)_J^\natural R_J^\natural \mathcal{M} = \varprojlim (X_\phi)_*^\natural(\mathcal{M}_j) \rightarrow \mathcal{M}_i = (?)_i \mathcal{M}$$

is induced by β_ϕ 's, where the limit is taken over $(\phi : i \rightarrow j) \in I_i^{(J \rightarrow I)}$. As J is a full subcategory, we have $I_i^{(J \rightarrow I)}$ equals i/J , and hence it has the initial object id_i . So the limit is equal to $\mathcal{M}_i$, and ε_i is the identity map. Since ε_i is an isomorphism for each $i \in J$, we have that ε is an isomorphism.

The proof of **2** is similar, and we omit it. $\square$

Let $\mathcal{C}$ be a small category. A connected component of $\mathcal{C}$ is a full subcategory of $\mathcal{C}$ whose object set is one of the equivalence classes of $\text{ob}(\mathcal{C})$ with respect to the transitive symmetric closure of the relation $\sim$ given by

$$c \sim c' \iff \mathcal{C}(c, c') \neq \emptyset.$$

7.12 Definition. We say that a subcategory J of I is admissible if

- 1 For $i \in I$, the category $(I_i^{(J \subset I)})^{\text{op}}$ is pseudofiltered.
- 2 For $j \in J$, we have id_j is the initial object of one of the connected components of $I_j^{(J \subset I)}$.

Note that for $j \in I$, the subcategory $j = (\{j\}, \{\text{id}_j\})$ of I is admissible.

In Lemma 7.5, the colimit is pseudo-filtered and hence it preserves exactness, if **1** is satisfied. As in the proof of Lemma 7.11, $(?)_j$ is a direct summand of $(?)_j \circ L_J$ for $j \in J$ so that L_J is faithful, if **2** is satisfied. We have the following.

7.13 Lemma. *Let $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$, and $K \subset J \subset I$ be admissible subcategories of I . Then $L_{J,K}^{\text{PA}}$ is faithful and exact. The morphism of sites $Q(X_\bullet|_J, K)$ is admissible. If, moreover, X_ϕ is flat for any $\phi \in I(k, j)$ with $j \in J$ and $k \in K$, then $L_{J,K}^{\natural}$ is faithful and exact for $\natural = \text{Mod}, \text{PM}$.*

(7.14) As in Example 6.6, **2**, we have an isomorphism

$$c_{i,f_\bullet} : (?)_i \circ (f_\bullet)_* \cong (f_i)_* \circ (?)_i. \quad (7.15)$$

The translation α_ϕ is described, as follows. For $\phi \in I(i, j)$,

$$\alpha_\phi(f_\bullet)_* : (Y_\phi)^*(?)_i(f_\bullet)_* \rightarrow (?)_j(f_\bullet)_*$$

agrees with

$$\begin{aligned} (Y_\phi)^*(?)_i(f_\bullet)_* &\xrightarrow{c_{i,f_\bullet}} (Y_\phi)^*(f_i)_*(?)_i \xrightarrow{\text{via } \theta} (f_j)_*(X_\phi)^*(?)_i \\ &\xrightarrow{(f_j)_*\alpha_\phi} (f_j)_*(?)_j \xrightarrow{c_{j,f_\bullet}^{-1}} (?)_j(f_\bullet)_*, \end{aligned} \quad (7.16)$$

where θ is the map defined and discussed in [23, (3.7.2)]. One of the definitions of θ is the composite

$$\begin{aligned} \theta : (Y_\phi)^*(f_i)_* &\xrightarrow{\text{via } u} (Y_\phi)^*(f_i)_*(X_\phi)_*(X_\phi)^* \cong \\ &(Y_\phi)^*(Y_\phi)_*(f_j)_*(X_\phi)^* \xrightarrow{\text{via } \varepsilon} (f_j)_*(X_\phi)^*. \end{aligned} \quad (7.17)$$

7.18 Proposition. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$, J a subcategory of I , and $i \in I$. Then the composite map*

$$(?)_i L_J(f_\bullet|_J)_* \xrightarrow{\text{via } \theta} (?)_i(f_\bullet)_* L_J \xrightarrow{\text{via } c_{i,f_\bullet}} (f_i)_*(?)_i L_J$$

agrees with the composite map

$$\begin{aligned} (?)_i L_J(f_\bullet|_J)_* &\xrightarrow{\text{via } \lambda_{J,i}} \varinjlim Y_\phi^*(?)_j (f_\bullet|_J)_* \xrightarrow{\text{via } c_{j,f_\bullet|_J}} \varinjlim Y_\phi^*(f_j)_*(?)_j \\ &\xrightarrow{\text{via } \theta} \varinjlim (f_i)_* X_\phi^*(?)_j \rightarrow (f_i)_* \varinjlim X_\phi^*(?)_j \xrightarrow{\text{via } \lambda_{J,i}^{-1}} (f_i)_*(?)_i L_J. \end{aligned}$$

Proof. Note that θ in the first composite map is the composite

$$\theta = \theta(J, f_\bullet) : L_J(f_\bullet|_J)_* \xrightarrow{\text{via } u} L_J(f_\bullet|_J)_*(?)_J L_J \xrightarrow{\cong} L_J(?)_J (f_\bullet)_* L_J \xrightarrow{\varepsilon} (f_\bullet)_* L_J.$$

The $\cong$ at the middle is the identity as a morphism of structure data. The description of u and ε are already given, and the proof is reduced to the iterative use of (7.7), (7.8), (7.9), and (7.14). The detailed argument is left to a patient reader. The reason why the second map involves θ is (7.14). $\square$

Similarly, we have the following.

7.19 Proposition. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$, J a subcategory of I , and $i \in I$. Then the composite map*

$$(f_i)^*(?)_i L_J \xrightarrow{\text{via } \theta(f_\bullet, i)} (?)_i (f_\bullet)_* L_J \xrightarrow{\text{via } d_{f_\bullet, J}} (?)_i L_J (f_\bullet|_J)^*$$

agrees with the composite map

$$\begin{aligned} (f_i)^*(?)_i L_J &\xrightarrow{\text{via } \lambda_{J,i}} (f_i)^* \varinjlim Y_\phi^*(?)_j \cong \varinjlim X_\phi^*(f_j)^*(?)_j \\ &\xrightarrow{\text{via } \theta(f_\bullet|_J, j)} \varinjlim X_\phi^*(?)_j (f_\bullet|_J)^* \xrightarrow{\text{via } \lambda_{J,i}^{-1}} (?)_i L_J (f_\bullet|_J)^*. \end{aligned}$$

The proof is left to the reader. The proof of Proposition 7.18 and Proposition 7.19 are formal, and the propositions are valid for $\mathfrak{h} = \text{PM}, \text{Mod}, \text{PA}$, and AB .

Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$, and $J \subset I$ a subcategory. The inverse image $(f_\bullet)_\mathfrak{h}^*$ is compatible with the restriction $(?)_J$.

7.20 Lemma. *The natural map*

$$\theta_\mathfrak{h} = \theta_\mathfrak{h}(f_\bullet, J) : ((f_\bullet)|_J)_\mathfrak{h}^* \circ (?)_J \rightarrow (?)_J \circ (f_\bullet)_\mathfrak{h}^*$$

is an isomorphism for $\mathfrak{h} = \text{PA}, \text{AB}, \text{PM}, \text{Mod}$. In particular, $f_\bullet^{-1} : \text{Zar}(Y_\bullet) \rightarrow \text{Zar}(X_\bullet)$ is an admissible continuous functor.

Proof. Note that θ is the composite map

$$(f_\bullet|_J)^*(?)_J \xrightarrow{\text{via } u} (f_\bullet|_J)^*(?)_J (f_\bullet)_* f_\bullet^* \xrightarrow{\text{via } c} (f_\bullet|_J)^*(f_\bullet|_J)_*(?)_J f_\bullet^* \xrightarrow{\text{via } \varepsilon} (?)_J f_\bullet^*,$$

see [23, (3.7.2)].

We construct a natural map

$$\eta : ((f_\bullet|_J)_\natural^* \circ (?)_J \rightarrow (?)_J \circ (f_\bullet)_\natural^*,$$

for the case where $\natural = \text{PM}$.

Let $\mathcal{M} \in \text{PM}(Y_\bullet)$, and $(j, U) \in \text{Zar}(X_\bullet|_J)$. We have

$$\Gamma((j, U), (f_\bullet|_J)^* \mathcal{M}_J) = \varinjlim \Gamma((j, U), \mathcal{O}_{X_\bullet}) \otimes_{\Gamma((j', V), \mathcal{O}_{Y_\bullet})} \Gamma((j', V), \mathcal{M}),$$

where the colimit is taken over $(j', V) \in (I_{(j, U)}^{(f_\bullet|_J)^{-1}})^{\text{op}}$. On the other hand, we have

$$\Gamma((j, U), (?)_J f_\bullet^* \mathcal{M}) = \varinjlim \Gamma((j, U), \mathcal{O}_{X_\bullet}) \otimes_{\Gamma((i, V), \mathcal{O}_{Y_\bullet})} \Gamma((i, V), \mathcal{M}),$$

where the colimit is taken over $(i, V) \in (I_{(j, U)}^{f_\bullet^{-1}})^{\text{op}}$. There is an obvious map from the first to the second.

To verify that this is an isomorphism, it suffices to show that the category $(I_{(j, U)}^{f_\bullet^{-1}})^{\text{op}}$ is final in the category $(I_{(j, U)}^{f_\bullet^{-1}})^{\text{op}}$. In fact, any $(\phi, h) : (j, U) \rightarrow (i, f_i^{-1}(V))$ with $(i, V) \in \text{Zar}(Y_\bullet)$ factors through

$$(\text{id}_j, h) : (j, U) \rightarrow (j, f_j^{-1}(Y_\phi)^{-1}(V)).$$

Hence, $\eta_\natural$ is an isomorphism for $\natural = \text{PM}$. The construction for the case where $\natural = \text{PA}$ is similar. As $(?)_J$ is compatible with the sheafification by Lemma 3.18, an isomorphism η for $\natural = \text{Mod}, \text{AB}$ is also defined.

In any case, it is straightforward to check that the diagram

$$\begin{array}{ccc} (?)_J & \xrightarrow{\text{id}} & (?)_J \\ \downarrow \text{via } u & & \downarrow \text{via } u \\ (f_\bullet|_J)_*(f_\bullet|_J)^*(?)_J & \xrightarrow{\text{via } \eta} & (f_\bullet|_J)_*(?)_J f_\bullet^* \xrightarrow{\text{via } c^{-1}} (?)_J (f_\bullet)_* f_\bullet^* \end{array} \quad (7.21)$$

is commutative. By the definition of θ and the identity $(\varepsilon(f_\bullet|_J)^*) \circ ((f_\bullet|_J)^* u) = \text{id}$, it is easy to check that $\eta^{-1} \theta = \text{id}$. This shows $\theta = \eta$, and we are done. $\square$

7.22 Lemma. *Let the notation be as above, and $\mathcal{M}, \mathcal{N} \in \mathfrak{h}(Y_\bullet)$. Then the diagram*

$$\begin{array}{ccccc}
(f_\bullet|_J)^*(\mathcal{M}_J \otimes \mathcal{N}_J) & \xrightarrow{m} & ((f_\bullet|_J)^*(\mathcal{M} \otimes \mathcal{N}))_J & \xrightarrow{\theta} & ((f_\bullet)^*(\mathcal{M} \otimes \mathcal{N}))_J \\
\downarrow \Delta & & & & \downarrow (?)_J \Delta \\
(f_\bullet|_J)^*\mathcal{M}_J \otimes (f_\bullet|_J)^*\mathcal{N}_J & \xrightarrow{\theta \otimes \theta} & ((f_\bullet)^*\mathcal{M})_J \otimes ((f_\bullet)^*\mathcal{N})_J & \xrightarrow{m} & ((f_\bullet)^*\mathcal{M} \otimes (f_\bullet)^*\mathcal{N})_J \\
& & & & (7.23)
\end{array}$$

is commutative.

Proof. This is an immediate consequence of Lemma 2.27. $\square$

7.24 Corollary. *The adjoint pair $((?)_{\text{Mod}}^*, (?)_*^{\text{Mod}})$ over the category $\mathcal{P}(I, \text{Sch}/S)$ is Lipman.*

Proof. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism of $\mathcal{P}(I, \text{Sch}/S)$. The morphism $\eta : f^*\mathcal{O}_{Y_\bullet} \rightarrow \mathcal{O}_{X_\bullet}$ is an isomorphism, like any such morphism arising from a ringed continuous functor. Let us consider $\mathcal{M}, \mathcal{N} \in \mathfrak{h}(Y_\bullet)$. To verify that Δ is an isomorphism, it suffices to show that

$$(?)_i \Delta : (f_\bullet^*(\mathcal{M} \otimes \mathcal{N}))_i \rightarrow (f_\bullet^*\mathcal{M} \otimes f_\bullet^*\mathcal{N})_i$$

is an isomorphism for any $i \in \text{ob}(I)$. Now consider the diagram (7.23) for $J = i$. Horizontal maps in the diagram are isomorphisms by (7.3) and (7.20). The left Δ is an isomorphism, since f_i is a morphism of single schemes. By Lemma 7.22, $(?)_i \Delta$ is also an isomorphism. $\square$

(7.25) The description of the translation map α_ϕ is as follows. For $\phi \in I(i, j)$,

$$\alpha_\phi : X_\phi^*(?)_i f_\bullet^* \rightarrow (?)_j f_\bullet^*$$

is the composite

$$X_\phi^*(?)_i f_\bullet^* \xrightarrow{X_\phi^* \theta^{-1}} X_\phi^* f_i^* (?)_i \cong f_j^* Y_\phi^* (?)_i \xrightarrow{f_j^* \alpha_\phi} f_j^* (?)_j \xrightarrow{\theta} (?)_j f_\bullet^*.$$

(7.26) Let $X_\bullet \in \mathcal{P}$, and $\mathcal{M}, \mathcal{N} \in \mathfrak{h}(X_\bullet)$. Although there is a canonical map

$$(?)_i : \underline{\text{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_i \rightarrow \underline{\text{Hom}}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i)$$

arising from the closed structure for $i \in I$, this may not be an isomorphism. However, we have the following.

7.27 Lemma. *Let $i \in I$. If $\mathcal{M}$ is equivariant, then the canonical map*

$$H_i : \underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_i \rightarrow \underline{\mathrm{Hom}}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i)$$

is an isomorphism.

Proof. It suffices to prove that

$$H_i : \mathrm{Hom}_{\mathfrak{h}(\mathrm{Zar}(X_\bullet)/(i, U))}(\mathcal{M}|_{(i, U)}, \mathcal{N}|_{(i, U)}) \rightarrow \mathrm{Hom}_{\mathfrak{h}(U)}(\mathcal{M}_i|_U, \mathcal{N}_i|_U)$$

is an isomorphism for any Zariski open set U in X_i .

To give an element of $\varphi \in \mathrm{Hom}_{\mathfrak{h}(\mathrm{Zar}(X_\bullet)/(i, U))}(\mathcal{M}|_{(i, U)}, \mathcal{N}|_{(i, U)})$ is the same as to give a family $(\varphi_\phi)_{\phi: i \rightarrow j}$ with

$$\varphi_\phi \in \mathrm{Hom}_{\mathfrak{h}(X_\phi^{-1}(U))}(\mathcal{M}_j|_{X_\phi^{-1}(U)}, \mathcal{N}_j|_{X_\phi^{-1}(U)})$$

such that

$$\varphi_\phi \circ (\alpha_\phi(\mathcal{M}))|_{X_\phi^{-1}(U)} = (\alpha_\phi(\mathcal{N}))|_{X_\phi^{-1}(U)} \circ ((X_\phi)|_{X_\phi^{-1}(U)})_{\mathfrak{h}}^*(\varphi_{\mathrm{id}_i}).$$

As $\alpha_\phi(\mathcal{M})$ is an isomorphism for any $\phi : i \rightarrow j$, we have that such a (φ_ϕ) is uniquely determined by φ_{id_i} . Hence H_i is bijective, as desired. $\square$

7.28 Lemma. *Let J be a subcategory of I . If $\mathcal{M}$ is equivariant, then the canonical map*

$$H_J : \underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_J \rightarrow \underline{\mathrm{Hom}}_{\mathfrak{h}(X_J)}(\mathcal{M}_J, \mathcal{N}_J)$$

is an isomorphism.

Proof. It suffices to show that

$$(H_J)_i : (\underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_J)_i \rightarrow \underline{\mathrm{Hom}}_{\mathfrak{h}(X_J)}(\mathcal{M}_J, \mathcal{N}_J)_i$$

is an isomorphism for each $i \in J$. By Lemma 2.10, the composite map

$$\begin{aligned} \underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_i &\cong (\underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})_J)_i \\ &\xrightarrow{(H_J)_i} \underline{\mathrm{Hom}}_{\mathfrak{h}(X_J)}(\mathcal{M}_J, \mathcal{N}_J)_i \xrightarrow{H_i} \underline{\mathrm{Hom}}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i) \end{aligned}$$

agrees with H_i . As $\mathcal{M}_J$ is also equivariant, we have that the two H_i are isomorphisms by Lemma 7.27, and hence $(H_J)_i$ is an isomorphism for any $i \in J$. $\square$

(7.29) By the lemma, the sheaf $\underline{\mathrm{Hom}}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N})$ is given by the collection

$$(\underline{\mathrm{Hom}}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i))_{i \in I}$$

provided $\mathcal{M}$ is equivariant. The structure map is the canonical composite map

$$\begin{aligned} \alpha_\phi : (X_\phi)_\#^* \underline{\mathrm{Hom}}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i) &\xrightarrow{P} \underline{\mathrm{Hom}}_{\mathfrak{h}(X_j)}((X_\phi)_\#^* \mathcal{M}_i, (X_\phi)_\#^* \mathcal{N}_i) \\ &\xrightarrow{\underline{\mathrm{Hom}}_{\mathfrak{h}(X_j)}(\alpha_\phi^{-1}, \alpha_\phi)} \underline{\mathrm{Hom}}_{\mathfrak{h}(X_j)}(\mathcal{M}_j, \mathcal{N}_j). \end{aligned}$$

Similarly, the following is also easy to prove.

7.30 Lemma. *Let $i \in I$ be an initial object of I . Then the following hold:*

1 *If $\mathcal{M} \in \mathfrak{h}(X_\bullet)$ is equivariant, then*

$$(\cdot)_i : \mathrm{Hom}_{\mathfrak{h}(X_\bullet)}(\mathcal{M}, \mathcal{N}) \rightarrow \mathrm{Hom}_{\mathfrak{h}(X_i)}(\mathcal{M}_i, \mathcal{N}_i)$$

is an isomorphism.

2 *$(\cdot)_i : \mathrm{EM}(X_\bullet) \rightarrow \mathrm{Mod}(X_i)$ is an equivalence, whose quasi-inverse is L_i .*

The fact that $L_i(\mathcal{M})$ is equivariant for $\mathcal{M} \in \mathrm{Mod}(X_i)$ is checked directly from the definition.

8 Quasi-coherent sheaves over a diagram of schemes

Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \mathrm{Sch}/S)$.

(8.1) Let $\mathcal{M} \in \mathrm{Mod}(X_\bullet)$. We say that $\mathcal{M}$ is locally quasi-coherent (resp. locally coherent) if $\mathcal{M}_i$ is quasi-coherent (resp. coherent) for any $i \in I$. We say that $\mathcal{M}$ is *quasi-coherent* if for any $(i, U) \in \mathrm{Zar}(X_\bullet)$ with $U = \mathrm{Spec} A$ being affine, there exists an exact sequence in $\mathrm{Mod}(\mathrm{Zar}(X_\bullet)/(i, U))$ of the form

$$(O_{X_\bullet}|_{(i, U)})^{(T)} \rightarrow (O_{X_\bullet}|_{(i, U)})^{(\Sigma)} \rightarrow \mathcal{M}|_{(i, U)} \rightarrow 0, \quad (8.2)$$

where T and Σ are arbitrary small sets.

8.3 Lemma. *Let $\mathcal{M} \in \mathrm{Mod}(X_\bullet)$. Then the following are equivalent.*

1 $\mathcal{M}$ is quasi-coherent.

2 $\mathcal{M}$ is locally quasi-coherent and equivariant.

3 For any morphism $(\phi, h) : (j, V) \rightarrow (i, U)$ in $\text{Zar}(X_\bullet)$ such that $V = \text{Spec } B$ and $U = \text{Spec } A$ are affine, the canonical map $B \otimes_A \Gamma((i, U), \mathcal{M}) \rightarrow \Gamma((j, V), \mathcal{M})$ is an isomorphism.

Proof. **1** $\Rightarrow$ **2** Let $i \in I$ and U an affine open subset of X_i . Then there is an exact sequence of the form (8.2). Applying the restriction functor $\text{Mod}(\text{Zar}(X_\bullet)/(i, U)) \rightarrow \text{Mod}(U)$, we get an exact sequence

$$\mathcal{O}_U^{(T)} \rightarrow \mathcal{O}_U^{(\Sigma)} \rightarrow (\mathcal{M}_i)|_U \rightarrow 0,$$

which shows that $\mathcal{M}_i$ is quasi-coherent for any $i \in I$. We prove that $\alpha_\phi(\mathcal{M})$ is an isomorphism for any $\phi : i \rightarrow j$, to show that $\mathcal{M}$ is equivariant. Take an affine open covering (U_λ) of X_i , and we prove that $\alpha_\phi(\mathcal{M})$ is an isomorphism over $X_\phi^{-1}(U_\lambda)$ for each λ . But this is obvious by the existence of an exact sequence of the form (8.2) and the five lemma.

2 $\Rightarrow$ **3** Set $W := X_\phi^{-1}(U)$, and let $\iota : V \hookrightarrow W$ be the inclusion map. Obviously, we have $h = (X_\phi)|_W \circ \iota$. As $\mathcal{M}$ is equivariant, we have that the canonical map

$$\alpha_\phi|_W(\mathcal{M}) : (X_\phi)|_{W_{\text{Mod}}}^*(\mathcal{M}_i)|_U \rightarrow (\mathcal{M}_j)|_W$$

is an isomorphism. Applying ι_{Mod}^* to the isomorphism, we have that $h_{\text{Mod}}^*((\mathcal{M}_i)|_U) \cong (\mathcal{M}_j)|_V$. The assertion follows from the assumption that $\mathcal{M}_i$ is quasi-coherent.

3 $\Rightarrow$ **1** Let $(i, U) \in \text{Zar}(X_\bullet)$ with $U = \text{Spec } A$ affine. There is a presentation of the form

$$A^{(T)} \rightarrow A^{(\Sigma)} \rightarrow \Gamma((i, U), \mathcal{M}) \rightarrow 0.$$

It suffices to prove that the induced sequence (8.2) is exact. To verify this, it suffices to prove that the sequence is exact after taking the section at $((\phi, h) : (j, V) \rightarrow (i, U)) \in \text{Zar}(X_\bullet)/(i, U)$ with $V = \text{Spec } B$ being affine. We have a commutative diagram

$$\begin{array}{ccccccc} B \otimes_A A^{(T)} & \rightarrow & B \otimes_A A^{(\Sigma)} & \rightarrow & B \otimes \Gamma((i, U), \mathcal{M}) & \rightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \\ B^{(T)} & \rightarrow & B^{(\Sigma)} & \rightarrow & \Gamma((j, V), \mathcal{M}) & \rightarrow & 0 \end{array}$$

whose first row is exact and vertical arrows are isomorphisms. Hence, the second row is also exact, and (8.2) is exact. $\square$

8.4 Definition. We say that $\mathcal{M} \in \text{Mod}(X_\bullet)$ is *coherent* if it is equivariant and $\mathcal{M}_i$ is coherent for any $i \in I$. We denote the full subcategory of $\text{Mod}(X_\bullet)$ consisting of coherent objects by $\text{Coh}(X_\bullet)$.

(8.5) Let $J \subset I$ be a subcategory. We say that J is *big* in I if for any $(\psi : j \rightarrow k) \in \text{Mor}(I)$, there exists some $(\phi : i \rightarrow j) \in \text{Mor}(J)$ such that $\psi \circ \phi \in \text{Mor}(J)$. Note that $\text{ob}(J) = \text{ob}(I)$ if J is big in I . Let $\mathbb{Q}$ be a property of morphisms of schemes. We say that $X_\bullet$ has $\mathbb{Q}$ J -arrows if $(X_\bullet)|_J$ has $\mathbb{Q}$ -arrows.

8.6 Lemma. Let $J \subset I$ be a subcategory, and $\mathcal{M} \in \text{Mod}(X_\bullet)$.

- 1 The full subcategory $\text{LQco}(X_\bullet)$ of $\text{Mod}(X_\bullet)$ consisting of locally quasi-coherent objects is a plump subcategory.
- 2 If $\mathcal{M}$ is equivariant (resp. locally quasi-coherent, quasi-coherent), then so is $\mathcal{M}_J^{\text{Mod}}$.
- 3 If J is big in I and $\mathcal{M}_J^{\text{Mod}}$ is equivariant (resp. locally quasi-coherent, quasi-coherent), then so is $\mathcal{M}$.
- 4 If J is big in I and $X_\bullet$ has flat J -arrows, then the full subcategory $\text{EM}(X_\bullet)$ (resp. $\text{Qco}(X_\bullet)$) of $\text{Mod}(X_\bullet)$ consisting of equivariant (resp. quasi-coherent) objects is a plump subcategory.
- 5 If J is big in I , then $(?)_J$ is faithful and exact.

Proof. 1 and 2 are trivial.

We prove 3. The assertion for the local quasi-coherence is obvious, because we have $\text{ob}(J) = \text{ob}(I)$. By Lemma 8.3, it remains to show the assertion for the equivariance. Let us assume that $\mathcal{M}_J^{\text{Mod}}$ is equivariant and $\psi : j \rightarrow k$ be a morphism in I , and take $\phi : i \rightarrow j$ such that $\phi, \psi\phi \in \text{Mor}(J)$. Then the composite map

$$(X_{\psi\phi})_{\text{Mod}}^*(\mathcal{M}_i) \cong (X_\psi)_{\text{Mod}}^*(X_\phi)_{\text{Mod}}^*(\mathcal{M}_i) \xrightarrow{(X_\psi)_{\text{Mod}}^* \alpha_\phi^{\text{Mod}}} (X_\psi)_{\text{Mod}}^*(\mathcal{M}_j) \xrightarrow{\alpha_\psi^{\text{Mod}}} \mathcal{M}_k,$$

which agrees with $\alpha_{\psi\phi}^{\text{Mod}}$, is an isomorphism by assumption. As we have α_ϕ^{Mod} is also an isomorphism, we have that α_ψ^{Mod} is an isomorphism. Thus $\mathcal{M}$ is equivariant.

4 By **1** and Lemma 8.3, it suffices to prove the assertion only for the equivariance. Let

$$\mathcal{M}_1 \rightarrow \mathcal{M}_2 \rightarrow \mathcal{M}_3 \rightarrow \mathcal{M}_4 \rightarrow \mathcal{M}_5$$

be an exact sequence in $\text{Mod}(X_\bullet)$, and assume that $\mathcal{M}_i$ is equivariant for $i = 1, 2, 4, 5$. We prove that $\mathcal{M}_3$ is equivariant. The sequence remains exact after applying the functor $(?)_J^{\text{Mod}}$. By **3**, replacing I by J and $X_\bullet$ by $(X_\bullet)|_J$, we may assume that $X_\bullet$ has flat arrows. Now the assertion follows easily from the five lemma.

The assertion **5** is obvious, because $\text{ob}(J) = \text{ob}(I)$. $\square$

8.7 Lemma. *Let $(\mathcal{M}_\lambda)$ be a diagram in $\text{Mod}(X_\bullet)$. If each $\mathcal{M}_\lambda$ is locally quasi-coherent (resp. equivariant, quasi-coherent), then so is $\varinjlim \mathcal{M}_\lambda$.*

Proof. As $(?)_i$ preserves colimits, the assertion for local quasi-coherence is trivial. Assume that each $\mathcal{M}_\lambda$ is equivariant. For $(\phi : i \rightarrow j) \in \text{Mor}(I)$, $\alpha_\phi(\mathcal{M}_\lambda)$ is an isomorphism. As $\alpha_\phi(\varinjlim \mathcal{M}_\lambda)$ is nothing but the composite

$$(X_\phi)_{\text{Mod}}^*((\varinjlim \mathcal{M}_\lambda)_i) \cong \varinjlim (X_\phi)_{\text{Mod}}^*(\mathcal{M}_\lambda)_i \xrightarrow{\varinjlim \alpha_\phi(\mathcal{M}_\lambda)} \varinjlim (\mathcal{M}_\lambda)_j \cong (\varinjlim \mathcal{M}_\lambda)_j,$$

it is an isomorphism. The rest of the assertions follow. $\square$

By Lemma 7.4, we have the following.

8.8 Lemma. *Let $J \subset I$ be a subcategory, and $\mathcal{M} \in \text{LQco}(X_\bullet|_J)$. Then we have $L_J^{\text{Mod}}(\mathcal{M}) \in \text{LQco}(X_\bullet)$.*

Similarly, we have the following.

8.9 Lemma. *Let $j \in I$. Assume that $X_\bullet$ has separated quasi-compact arrows, and that $I(i, j)$ is finite for any $i \in I$. If $\mathcal{M} \in \text{Qco}(X_j)$, then we have $R_j \mathcal{M} \in \text{LQco}(X_\bullet)$.*

The following is also proved easily, using (7.3).

8.10 Lemma. *Let $\mathcal{M}$ and $\mathcal{N}$ be locally quasi-coherent (resp. equivariant, quasi-coherent) $\mathcal{O}_{X_\bullet}$ -modules. Then $\mathcal{M} \otimes_{\mathcal{O}_{X_\bullet}} \mathcal{N}$ is also locally quasi-coherent (resp. equivariant, quasi-coherent).*

The following is a consequence of the observation in (7.29).

8.11 Lemma. *Let $\mathcal{M}$ be a coherent $\mathcal{O}_{X_\bullet}$ -module, and $\mathcal{N}$ a locally quasi-coherent $\mathcal{O}_{X_\bullet}$ -module. Then $\underline{\mathrm{Hom}}_{\mathrm{Mod}(X_\bullet)}(\mathcal{M}, \mathcal{N})$ is locally quasi-coherent. If, moreover, there is a big subcategory J of I such that $X_\bullet$ has flat J -arrows and $\mathcal{N}$ is quasi-coherent, then $\underline{\mathrm{Hom}}_{\mathrm{Mod}(X_\bullet)}(\mathcal{M}, \mathcal{N})$ is quasi-coherent.*

(8.12) Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$.

As θ in (7.17), which appears in (7.16) is not an isomorphism in general, $(f_\bullet)_*^{\natural}(\mathcal{M})$ need not be equivariant even if $\mathcal{M}$ is equivariant. However, we have

8.13 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$, and J a big subcategory of I . Then we have the following:*

- 1 $f_\bullet$ is of fiber type if and only if $(f_\bullet)|_J$ is of fiber type.
- 2 If $f_\bullet$ is quasi-compact separated and $\mathcal{M} \in \mathrm{LQco}(X_\bullet)$, then $(f_\bullet)_*(\mathcal{M}) \in \mathrm{LQco}(Y_\bullet)$.
- 3 If $Y_\bullet$ has flat J -arrows, $f_\bullet$ is of fiber type and quasi-compact separated, and $\mathcal{M} \in \mathrm{Qco}(X_\bullet)$, then we have $(f_\bullet)_*(\mathcal{M}) \in \mathrm{Qco}(Y_\bullet)$.

Proof. **1** Assume that $f_\bullet|_J$ is of fiber type, and let $\psi : j \rightarrow k$ be a morphism in I . Take $\phi : i \rightarrow j$ such that $\phi, \psi\phi \in \mathrm{Mor}(J)$. Consider the commutative diagram

$$\begin{array}{ccccc} X_k & \xrightarrow{X_\psi} & X_j & \xrightarrow{X_\phi} & X_i \\ \downarrow f_k & \text{(a)} & \downarrow f_j & \text{(b)} & \downarrow f_i \\ Y_k & \xrightarrow{Y_\psi} & Y_j & \xrightarrow{Y_\phi} & Y_i. \end{array}$$

By assumption, the square (b) and the whole rectangle ((a)+(b)) are fiber squares. Hence (a) is also a fiber square. This shows that $f_\bullet$ is of fiber type. The converse is obvious.

The assertion **2** is obvious by the isomorphism $(f_\bullet)_*(\mathcal{M})_i \cong (f_i)_*(\mathcal{M}_i)$ for $i \in I$, see [11, (9.2.2)].

We prove **3**. By Lemma 8.6, we may assume that $J = I$. Then $(f_\bullet)_*(\mathcal{M})$ is locally quasi-coherent by **2**. As $\mathcal{M}$ is equivariant and θ in (7.17) is an isomorphism by [12, (1.4.15)], we have that $(f_\bullet)_*(\mathcal{M})$ is equivariant. Hence by Lemma 8.3, $(f_\bullet)_*(\mathcal{M})$ is quasi-coherent. $\square$

(8.14) If $f_\bullet$ is quasi-compact separated of fiber type and $Y_\bullet$ has flat J -arrows for some big subcategory J of I , then $(f_\bullet)_*^{\text{Qco}} : \text{Qco}(X_\bullet) \rightarrow \text{Qco}(Y_\bullet)$ is defined as the restriction of $(f_\bullet)_*^{\text{Mod}}$.

Similarly, we have the following.

8.15 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y_\bullet \rightarrow Z_\bullet$ be morphisms in $\mathcal{P}$. Then the following hold.*

0 *An isomorphism is a morphism of fiber type.*

1 *If $f_\bullet$ and $g_\bullet$ are of fiber type, then so is $g_\bullet \circ f_\bullet$.*

2 *If $g_\bullet$ and $g_\bullet \circ f_\bullet$ are of fiber type, then so is $f_\bullet$.*

3 *If $f_\bullet$ is faithfully flat of fiber type and $g_\bullet \circ f_\bullet$ is of fiber type, then $g_\bullet$ is of fiber type.*

8.16 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y'_\bullet \rightarrow Y_\bullet$ be morphisms in $\mathcal{P}$. Let $f'_\bullet : X'_\bullet \rightarrow Y'_\bullet$ be the base change of $f_\bullet$ by $g_\bullet$.*

1 *If $f_\bullet$ is of fiber type, then so is $f'_\bullet$.*

2 *If $f'_\bullet$ is of fiber type and $g_\bullet$ is faithfully flat, then $f_\bullet$ is of fiber type.*

(8.17) Let $f : X \rightarrow Y$ be a morphism of schemes. If f is quasi-compact separated, then f_*^{Qco} is compatible with filtered inductive limits.

8.18 Lemma ([21, p.641, Proposition 6]). *Let $f : X \rightarrow Y$ be a quasi-compact separated morphism of schemes, and $(\mathcal{M}_i)$ a pseudo-filtered inductive system of $\mathcal{O}_X$ -modules. Then the canonical map*

$$\varinjlim f_* \mathcal{M}_i \rightarrow f_* \varinjlim \mathcal{M}_i$$

is an isomorphism.

By the lemma, the following follows immediately.

8.19 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. If $f_\bullet$ is quasi-compact and separated, then $(f_\bullet)_*^{\text{Mod}}$ and $(f_\bullet)_*^{\text{LQco}}$ preserve pseudo-filtered inductive limits.*

8.20 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$. Let J be an admissible subcategory of I . If $Y_\bullet$ has flat arrows and $f_\bullet$ is of fiber type and quasi-compact separated, then the canonical map*

$$\theta(J, f_\bullet) : L_J \circ (f_J)_* \rightarrow (f_\bullet)_* \circ L_J$$

is an isomorphism of functors from $\mathrm{LQco}(X_\bullet|_J)$ to $\mathrm{LQco}(Y_\bullet)$.

Proof. This is obvious by Proposition 7.18, Lemma 8.18, and [12, (1.4.15)]. $\square$

The following is obvious by Lemma 7.20 and (7.25).

8.21 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$. If $\mathcal{M} \in \mathrm{Mod}(Y_\bullet)$ is equivariant (resp. locally quasi-coherent, quasi-coherent), then so is $(f_\bullet)_*^{\mathrm{Mod}}(\mathcal{M})$. If $\mathcal{M} \in \mathrm{Mod}(Y_\bullet)$, $f_\bullet$ is faithfully flat, and $(f_\bullet)_*^{\mathrm{Mod}}(\mathcal{M})$ is equivariant, then we have $\mathcal{M}$ is equivariant.*

The restriction $(f_\bullet)^* : \mathrm{Qco}(Y_\bullet) \rightarrow \mathrm{Qco}(X_\bullet)$ is sometimes denoted by $(f_\bullet)_{\mathrm{Qco}}^*$.

9 Derived functors of functors on sheaves of modules over diagrams of schemes

Let I be a small category, and S a scheme. Set $\mathcal{P} := \mathcal{P}(I, \mathrm{Sch}/S)$, and let $X_\bullet \in \mathcal{P}$.

9.1 Proposition. *Let $X_\bullet \in \mathcal{P}$, and $\mathbb{I} \in K(\mathrm{Mod}(X_\bullet))$. We have $\mathbb{I}$ is K -limp if and only if so is $\mathbb{I}_i$ for $i \in I$.*

Proof. The only if part is proved in Lemma 4.29.

We prove the if part. Let $\mathbb{I} \rightarrow \mathbb{J}$ be a K -injective resolution, and let $\mathbb{C}$ be the mapping cone. Note that $\mathbb{C}_i$ is exact for each i .

Let $(U, i) \in \mathrm{Zar}(X_\bullet)$. We have an isomorphism

$$\Gamma((U, i), \mathbb{C}) \cong \Gamma(U, \mathbb{C}_i).$$

As $\mathbb{C}_i$ is K -limp by the only if part, these are exact for each (U, i) . It follows that $\mathbb{I}$ is K -limp. $\square$

9.2 Corollary. *Let J be a subcategory of I , and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \mathrm{Sch}/S)$. Then there is an isomorphism*

$$c(J, f_\bullet) : (?)_J R(f_\bullet)_* \cong R(f_\bullet|_J)_* (?)_J.$$

(9.3) Let X be a scheme, $x \in X$, and M an $\mathcal{O}_{X,x}$ -module. We define $\xi_x(M) \in \text{Mod}(X)$ by $\Gamma(U, \xi_x(M)) = M$ if $x \in U$, and zero otherwise. The restriction maps are defined in an obvious way. For an exact complex $\mathbb{H}$ of $\mathcal{O}_{X,x}$ -modules, $\xi_x(\mathbb{H})$ is exact not only as a complex of sheaves, but also as a complex of presheaves. For a morphism of schemes $f : X \rightarrow Y$, we have that $f_*\xi_x(M) \cong \xi_{f(x)}(M)$.

9.4 Lemma. Let $\mathbb{F} \in C(\text{Mod}(X_\bullet))$. The following are equivalent.

- 1 $\mathbb{F}$ is K -flat.
- 2 $\mathbb{F}_i$ is K -flat for $i \in \text{ob}(I)$.
- 3 $\mathbb{F}_{i,x}$ is a K -flat complex of $\mathcal{O}_{X,x}$ -modules for any $i \in \text{ob}(I)$.

Proof. **3** $\Rightarrow$ **1** Let $\mathbb{G} \in C(\text{Mod}(X_\bullet))$ be exact. We are to prove that $\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} \mathbb{G}$ is exact. For $i \in \text{ob}(I)$ and $x \in X$, we have

$$(\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} \mathbb{G})_{i,x} \cong (\mathbb{F}_i \otimes_{\mathcal{O}_{X_i}} \mathbb{G}_i)_x \cong \mathbb{F}_{i,x} \otimes_{\mathcal{O}_{X_{i,x}}} \mathbb{G}_{i,x}.$$

Since $\mathbb{G}_{i,x}$ is exact, $(\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} \mathbb{G})_{i,x}$ is exact. So $\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} \mathbb{G}$ is exact.

1 $\Rightarrow$ **3** Let $\mathbb{H} \in C(\text{Mod}(\mathcal{O}_{X_{i,x}}))$ be an exact complex, and we are to prove that $\mathbb{F}_{i,x} \otimes_{\mathcal{O}_{X_{i,x}}} \mathbb{H}$ is exact. For each $j \in \text{ob}(I)$,

$$(\mathbb{F})_j R_i \xi_x(\mathbb{H}) \cong \prod_{\phi \in I(j,i)} (X_\phi)_* \xi_x(\mathbb{H}) \cong \prod_{\phi \in I(j,i)} \xi_{X_\phi(x)}(\mathbb{H})$$

is exact, since a direct product of exact complexes of presheaves is exact. So $R_i \xi_x(\mathbb{H})$ is exact. It follows that $\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} R_i \xi_x(\mathbb{H})$ is exact. Hence

$$(\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}} R_i \xi_x(\mathbb{H}))_i \cong \mathbb{F}_i \otimes_{\mathcal{O}_{X_i}} \prod_{\phi \in I(i,i)} \xi_{X_\phi(x)} \mathbb{H}$$

is also exact. So $\mathbb{F}_i \otimes_{\mathcal{O}_{X_i}} \xi_{\text{id}_X(x)}(\mathbb{H}) = \mathbb{F}_i \otimes_{\mathcal{O}_{X_i}} \xi_x \mathbb{H}$ is exact. So

$$(\mathbb{F}_i \otimes_{\mathcal{O}_{X_i}} \xi_x \mathbb{H})_x \cong \mathbb{F}_{i,x} \otimes_{\mathcal{O}_{X_{i,x}}} (\xi_x \mathbb{H})_x \cong \mathbb{F}_{i,x} \otimes_{\mathcal{O}_{X_{i,x}}} \mathbb{H}$$

is also exact.

Applying **1** $\Leftrightarrow$ **3**, which has already been proved, to the complex $\mathbb{F}_i$ over the single scheme X_i , we get **2** $\Leftrightarrow$ **3**. $\square$

Hence by [34], we have the following.

9.5 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$. Then we have the following.*

- 1 *If $\mathbb{F} \in C(\text{Mod}(Y_\bullet))$ is K -flat, then so is $f_\bullet^* \mathbb{F}$.*
- 2 *If $\mathbb{F} \in C(\text{Mod}(Y_\bullet))$ is K -flat exact, then so is $f_\bullet^* \mathbb{F}$.*
- 3 *If $\mathbb{I} \in C(\text{Mod}(X_\bullet))$ is weakly K -injective, then so is $(f_\bullet)_* \mathbb{I}$.*

(9.6) By the lemma, the left derived functor $Lf_\bullet^*$, which we already know its existence by Lemma 7.20, can also be calculated by K -flat resolutions.

9.7 Lemma. *Let J be a subcategory of I , and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}$. Then we have the following.*

- 1 *The canonical map*

$$\theta(f_\bullet, J) : L(f_\bullet|_J)^*(?)_J \rightarrow (?)_J Lf_\bullet^*$$

is an isomorphism.

- 2 *The diagram*

$$\begin{array}{ccccc} (?)_J & \xrightarrow{\text{id}} & (?)_J \\ \downarrow \text{via } u & & \downarrow \text{via } u \\ R(f_\bullet|_J)_* L(f_\bullet|_J)^*(?)_J & \xrightarrow{\text{via } \theta} & R(f_\bullet|_J)_* (?)_J Lf_\bullet^* & \xrightarrow{\text{via } c^{-1}} & (?)_J R(f_\bullet)_* Lf_\bullet^* \end{array}$$

is commutative.

- 3 *The diagram*

$$\begin{array}{ccccc} (?)_J & \xrightarrow{\text{id}} & (?)_J \\ \uparrow \text{via } \varepsilon & & \uparrow \text{via } \varepsilon \\ L(f_\bullet|_J)^* R(f_\bullet|_J)_* (?)_J & \xrightarrow{\text{via } c^{-1}} & L(f_\bullet|_J)^* (?)_J R(f_\bullet)_* & \xrightarrow{\text{via } \theta} & (?)_J L(f_\bullet)^* R(f_\bullet)_* \end{array}$$

is commutative.

Proof. Since $(?)_J$ preserves K -flat complexes by Lemma 9.4, we have

$$L(f_\bullet|_J)^*(?)_J \cong L((f_\bullet|_J) \circ (?)_J).$$

On the other hand, it is obvious that we have $(?)_J Lf_\bullet^* \cong L((?)_J f_\bullet^*)$. By Lemma 7.20, we have a composite isomorphism

$$\theta' : L(f_\bullet|_J)^*(?)_J \cong L((f_\bullet|_J)^* \circ (?)_J) \xrightarrow{L\theta} L((?)_J \circ f_\bullet^*) \cong (?)_J L(f_\bullet)^*.$$

It is straightforward to check the diagram in **2** (θ there should be replaced by θ') is commutative utilizing (7.21) (note that η there equals θ). As in the proof of Lemma 7.20, we have $\theta = \theta'$, and **1** and **2** are proved. The proof of **3** is formal, and we omit it. $\square$

9.8 Lemma. *Let $X_\bullet \in \mathcal{P}$, and $\mathbb{F}, \mathbb{G} \in D(\text{Mod}(X_\bullet))$. Then we have the following.*

- 1** $\mathbb{F}_J \otimes_{\mathcal{O}_{X_\bullet|J}}^{\bullet, L} \mathbb{G}_J \cong (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{G})_J$ for any subcategory $J \subset I$.
- 2** If $\mathbb{F}$ and $\mathbb{G}$ has locally quasi-coherent cohomology groups, then $\underline{\text{Tor}}_i^{\mathcal{O}_{X_\bullet}}(\mathbb{F}, \mathbb{G})$ is also locally quasi-coherent for any $i \in \mathbb{Z}$.
- 3** Assume that there exists some big subcategory J of I such that $X_\bullet$ has flat J -arrows. If both $\mathbb{F}$ and $\mathbb{G}$ has equivariant (resp. quasi-coherent) cohomology groups, then $\underline{\text{Tor}}_i^{\mathcal{O}_{X_\bullet}}(\mathbb{F}, \mathbb{G})$ is also equivariant (resp. quasi-coherent).

Proof. The assertion **1** is an immediate consequence of Lemma 9.4 and Example 6.6, **5**.

2 In view of **1**, we may assume that $X = X_\bullet$ is a single scheme. As the question is local, we may assume that X is even affine.

We may assume that $\mathbb{F} = \varinjlim \mathbb{F}_n$, where $(\mathbb{F}_n)$ is the $\mathfrak{P}(X_\bullet)$ -special direct system such that each $\mathbb{F}_n$ is bounded above and has locally quasi-coherent cohomology groups as in Lemma 4.24, **4**. Similarly, we may assume that $\mathbb{G} = \varinjlim \mathbb{G}_n$. As filtered inductive limits are exact and compatible with tensor products, and the colimit of locally quasi-coherent sheaf is locally quasi-coherent, we may assume that both $\mathbb{F}$ and $\mathbb{G}$ are bounded above, flat, and has locally quasi-coherent cohomology groups. By [15, Proposition I.7.3], we may assume that both $\mathbb{F}$ and $\mathbb{G}$ are single quasi-coherent sheaves. This case is trivial.

3 In view of **1**, we may assume that $J = I$ and $X_\bullet$ has flat arrows. By **2**, it suffices to show the assertion for equivariance. Assuming that $\mathbb{F}$ and $\mathbb{G}$ are K -flat with equivariant cohomology groups, we prove that $\mathbb{F} \otimes \mathbb{G}$ has equivariant cohomology groups. This is enough.

Let $\phi : i \rightarrow j$ be a morphism of I . As X_ϕ is flat and $\mathbb{F}$ and $\mathbb{G}$ have equivariant cohomology groups, $\alpha_\phi : X_\phi^* \mathbb{F}_i \rightarrow \mathbb{F}_j$ and $\alpha_\phi : X_\phi^* \mathbb{G}_i \rightarrow \mathbb{G}_j$ are quasi-isomorphisms. The composite

$$X_\phi^*(\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^\bullet \mathbb{G})_i \cong X_\phi^* \mathbb{F}_i \otimes_{\mathcal{O}_{X_j}}^\bullet X_\phi^* \mathbb{G}_i \xrightarrow{\alpha_\phi \otimes \alpha_\phi} \mathbb{F}_j \otimes_{\mathcal{O}_{X_j}}^\bullet \mathbb{G}_j \cong (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^\bullet \mathbb{G})_j$$

is a quasi-isomorphism, since $\mathbb{F}_j$ and $X_\phi^* \mathbb{F}_i$ are K -flat. By (7.3), $\alpha_\phi(\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^\bullet \mathbb{G})$ is a quasi-isomorphism.

As $X_\bullet$ has flat arrows, this shows that $\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^\bullet \mathbb{G}$ has equivariant cohomology groups. $\square$

(9.9) Let $X_\bullet \in \mathcal{P}$, and J an admissible subcategory of I . By Lemma 7.13, the left derived functor

$$LL_J^{\text{Mod}} : D(\text{Mod}(X_\bullet|_J)) \rightarrow D(\text{Mod}(X_\bullet))$$

of L_J^{Mod} is defined, since $Q(X_\bullet, J)$ is admissible. This is also calculated using K -flat resolutions. Namely,

9.10 Lemma. *Let $X_\bullet$ and J be as above. If $\mathbb{F} \in K(\text{Mod}(X_\bullet|_J))$ is K -flat, then so is $L_J \mathbb{F}$. If $\mathbb{F}$ is K -flat exact, then so is $L_J \mathbb{F}$.*

Proof. This is trivial by (7.5). $\square$

9.11 Corollary. *Let $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$, J an admissible subcategory of I , and $\mathbb{I} \in K(\text{Mod}(X_\bullet))$. If $\mathbb{I}$ is weakly K -injective, then $\mathbb{I}_J$ is weakly K -injective.*

Proof. Let $\mathbb{F}$ be a K -flat exact complex in $K(\text{Mod}(X_\bullet|_J))$. Then,

$$\text{Hom}_{\text{Mod}(X_\bullet|_J)}^\bullet(\mathbb{F}, \mathbb{I}_J) \cong \text{Hom}_{\text{Mod}(X_\bullet)}^\bullet(L_J \mathbb{F}, \mathbb{I})$$

is exact by the lemma. By Lemma 4.24, **7**, we are done. $\square$

9.12 Lemma. *Let $X_\bullet$ and J be as above. Let $\mathbb{F} \in D_{\text{LQco}(X_\bullet|_J)}(\text{Mod}(X_\bullet|_J))$. Then, $L_J \mathbb{F} \in D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$.*

Proof. This is trivial by the standard spectral sequence argument, if $\mathbb{F}$ is bounded above. The general case follows immediately by Lemma 4.24, **4**. $\square$

(9.13) Let I be a small category, and S a scheme. Set $\mathcal{P} := \mathcal{P}(I, \text{Sch}/S)$. As we have seen, for a morphism $f_\bullet : X_\bullet \rightarrow Y_\bullet$, $f_\bullet^{-1} : \text{Zar}(Y_\bullet) \rightarrow \text{Zar}(X_\bullet)$ is an admissible ringed continuous functor by Lemma 7.20. Moreover, if J and K are admissible subcategories of I such that $J \subset K$, then $Q(X_\bullet|_J, K) : \text{Zar}(X_\bullet|_K) \rightarrow \text{Zar}(X_\bullet|_J)$ is also admissible. By Lemma 4.31 and Lemma 6.4, we have the following.

9.14 Example. Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y_\bullet \rightarrow Z_\bullet$ are morphisms in $\mathcal{P}(I, \text{Sch}/S)$. Let $K \subset J \subset I$ be admissible subcategories. Then we have the following.

1 There is a natural isomorphism

$$c_{I,J,K} : (?)_{K,I} \cong (?)_{K,J} \circ (?)_{J,I}.$$

Taking the conjugate,

$$d_{I,J,K} : LL_{I,J} \circ LL_{J,K} \cong LL_{I,K}$$

is induced.

2 There are natural isomorphism

$$c_{J,f_\bullet} : (?)_J \circ R(f_\bullet)_* \cong R(f_\bullet|_J)_* \circ (?)_J$$

and its conjugate

$$d_{J,f_\bullet} : LL_J \circ L(f_\bullet|_J)^* \cong L(f_\bullet)^* \circ LL_J.$$

3 We have

$$(c_{K,f_\bullet|_J} (?)_J) \circ ((?)_{K,J} c_{J,f_\bullet}) = (R(f_\bullet|_K)_* c_{I,J,K}) \circ c_{K,f_\bullet} \circ (c_{I,J,K}^{-1} R(f_\bullet)_*).$$

4 We have

$$(R(g_\bullet|_J)_* c_{J,f_\bullet}) \circ (c_{J,g_\bullet} R(f_\bullet)_*) = (c_{f_\bullet|_J, g_\bullet|_J} (?)_J) \circ c_{J, g_\bullet \circ f_\bullet} \circ ((?)_J c_{f_\bullet, g_\bullet}^{-1}),$$

where $c_{f_\bullet, g_\bullet} : R(g_\bullet \circ f_\bullet)_* \cong R(g_\bullet)_* \circ R(f_\bullet)_*$ is the canonical isomorphism, and similarly for $c_{f_\bullet|_J, g_\bullet|_J}$.

5 The adjoint pair $(L(?)_{\text{Mod}}^*, R(?)_*^{\text{Mod}})$ over the category $\mathcal{P}(I, \text{Sch}/S)$ is Lipman.

It is now quite formal to prove the following.

9.15 Lemma. *Let the notation be as in Example 9.14. Let $\theta = \theta(J, f_\bullet)$ denotes the natural map*

$$LL_J R(f_\bullet|_J)_* \rightarrow R(f_\bullet)_* LL_J$$

defined in [23, (3.7.2)]. Then we have the following.

1 *The diagram*

$$\begin{array}{ccccc} LL_J & \xrightarrow{\text{id}} & LL_J \\ \downarrow \text{via } u & & \downarrow \text{via } u \\ LL_J R(f_\bullet|_J)_* L(f_\bullet|_J)^* & \xrightarrow{\text{via } \theta} & R(f_\bullet)_* LL_J L(f_\bullet|_J)^* & \xrightarrow{\text{via } d} & R(f_\bullet)_* L f_\bullet^* LL_J \end{array}$$

is commutative.

2 *The diagram*

$$\begin{array}{ccccc} LL_J & \xrightarrow{\text{id}} & LL_J \\ \uparrow \text{via } \varepsilon & & \uparrow \text{via } \varepsilon \\ LL_J L(f_\bullet|_J)^* R(f_\bullet|_J)_* & \xrightarrow{\text{via } d} & L(f_\bullet)^* LL_J R(f_\bullet|_J)_* & \xrightarrow{\text{via } \theta} & L(f_\bullet)^* R(f_\bullet)_* LL_J \end{array}$$

is commutative.

3 *Let $h_\bullet : Y'_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}$. Set $X'_\bullet := Y'_\bullet \times_{Y_\bullet} X_\bullet$. Let $h'_\bullet : X'_\bullet \rightarrow X_\bullet$ be the second projection, and $f'_\bullet : X'_\bullet \rightarrow Y'_\bullet$ be the first projection. Then the diagram*

$$\begin{array}{ccccccc} LL_J L(h_\bullet|_J)^* R(f_\bullet|_J)_* & \xrightarrow{d} & L(h_\bullet)^* LL_J R(f_\bullet|_J)_* & \xrightarrow{\theta} & L(h_\bullet)^* R(f_\bullet)_* LL_J \\ \downarrow \theta & & & & \downarrow \theta \\ LL_J R(f'_\bullet|_J)_* L(h'_\bullet|_J)^* & \xrightarrow{\theta} & R(f'_\bullet)_* LL_J L(h'_\bullet|_J)^* & \xrightarrow{d} & R(f'_\bullet)_* L(h'_\bullet)^* LL_J \end{array}$$

is commutative.

Proof. Left to the reader.

10 The right adjoint of the derived direct image functor of a morphism of quasi-compact separated schemes

Let X be a quasi-compact separated scheme.

10.1 Lemma. *The following hold:*

- 1** *There is a right adjoint $\mathrm{qco}(X) : \mathrm{Mod}(X) \rightarrow \mathrm{Qco}(X)$ of the canonical inclusion $F_X : \mathrm{Qco}(X) \hookrightarrow \mathrm{Mod}(X)$.*
- 1'** *$\mathrm{qco}(X)$ preserves K -injective complexes. $R\mathrm{qco}(X) : D(\mathrm{Mod}(X)) \rightarrow D(\mathrm{Qco}(X))$ is right adjoint to $F_X : D(\mathrm{Qco}(X)) \rightarrow D(\mathrm{Mod}(X))$.*
- 2** *The functor $F_X : D(\mathrm{Qco}(X)) \rightarrow D(\mathrm{Mod}(X))$ is full and faithful, and induces an equivalence $D(\mathrm{Qco}(X)) \rightarrow D_{\mathrm{Qco}(X)}(\mathrm{Mod}(X))$.*
- 3** *The unit of adjunction $u : \mathrm{Id} \rightarrow R\mathrm{qco}(X)F_X$ is an isomorphism, and $\varepsilon : F_X R\mathrm{qco}(X) \rightarrow \mathrm{Id}$ is an isomorphism on $D_{\mathrm{Qco}(X)}(\mathrm{Mod}(X))$.*

Proof. See [19] for **1**. The assertion **1'** follows from **1** and Lemma 4.11. For **2**, see [6]. The assertion **3** follows from **2**. $\square$

By the lemma, the following follows immediately.

10.2 Corollary. *Let X be a quasi-compact separated scheme. Then $\mathrm{Qco}(X)$ has enough injectives and has direct products. For any $\mathbb{F} \in K(\mathrm{Qco}(X))$, there is a K -injective resolution $\mathbb{F} \rightarrow \mathbb{I}$ such that each term of $\mathbb{I}$ being an injective object of $\mathrm{Qco}(X)$.*

In fact, if $F_X \mathbb{F} \rightarrow \mathbb{J}$ is a K -injective resolution such that each term of $\mathbb{J}$ is injective, then the corresponding $\mathbb{F} \rightarrow \mathrm{qco} \mathbb{J}$ is still a quasi-isomorphism by the lemma, and this is a desired K -injective resolution.

(10.3) Let $f : X \rightarrow Y$ be a quasi-compact separated morphism of schemes. Then $f_*^{\mathrm{Qco}} : \mathrm{Qco}(X) \rightarrow \mathrm{Qco}(Y)$ is defined, and we have $F_Y \circ f_*^{\mathrm{Qco}} \cong f_*^{\mathrm{Mod}} \circ F_X$, where F_Y and F_X are the forgetful functors. By the corollary, there is a right derived functor Rf_*^{Qco} of f_*^{Qco} .

10.4 Lemma. *Let $f : X \rightarrow Y$ be a morphism of schemes. Assume that both X and Y are quasi-compact separated. Then, we have the following.*

1 *The canonical maps*

$$F_Y \circ Rf_*^{\text{Qco}} \cong R(F_Y \circ f_*^{\text{Qco}}) \cong R(f_*^{\text{Mod}} \circ F_X) \rightarrow Rf_*^{\text{Mod}} \circ F_X$$

are all isomorphisms.

2 *There are a left adjoint F of Rf_*^{Qco} and an isomorphism $F_X F \cong Lf_{\text{Mod}}^* F_Y$.*

3 *There is an isomorphism*

$$Rf_*^{\text{Qco}} R\text{qco} \cong R\text{qco} Rf_*^{\text{Mod}}.$$

Proof. We prove **1**. It suffices to show that, if $\mathbb{I} \in K(\text{Qco}(X))$ is K -injective, then $F_X \mathbb{I}$ is f_*^{Mod} -acyclic. By Corollary 10.2, we may assume that each term of $\mathbb{I}$ is injective. By [23, (3.9.3.5)], it suffices to show that an injective object $\mathcal{I}$ of $\text{Qco}(X)$ is f_*^{Mod} -acyclic. Let $g : Z \rightarrow X$ be an affine faithfully flat morphism such that Z is affine. Such a morphism exists, as X is quasi-compact separated. Then it is easy to see that $\mathcal{I}$ is a direct summand of $g_* \mathcal{J}$ for some injective object $\mathcal{J}$ in $\text{Qco}(Z)$. So it suffices to show that $F_X g_* \mathcal{J}$ is f_*^{Mod} -acyclic. Let $F_Z \mathcal{J} \rightarrow \mathbb{J}$ be an injective resolution. As g is affine and $\mathcal{J}$ is quasi-coherent, we have $R^i f_*^{\text{Mod}} F_Z \mathcal{J} = 0$ for $i > 0$. Hence $g_* F_Z \mathcal{J} \rightarrow g_* \mathbb{J}$ is a quasi-isomorphism, and hence is a K -limp resolution of $F_X g_* \mathcal{J} \cong g_* F_Z \mathcal{J}$. As $f \circ g$ is also affine, $f_* g_* F_Z \mathcal{J} \rightarrow f_* g_* \mathbb{J}$ is still a quasi-isomorphism, and this shows that $F_X g_* \mathcal{J}$ is f_* -acyclic.

2 Define $F : D(\text{Qco}(Y)) \rightarrow D(\text{Qco}(X))$ by $F := R\text{qco} Lf_{\text{Mod}}^* F_Y$. As we have $Lf_{\text{Mod}}^* F_Y(D(\text{Qco}(Y))) \subset D_{\text{Qco}(X)}(D(\text{Mod}(X)))$, we have

$$F_X F = F_X R\text{qco} Lf_{\text{Mod}}^* F_Y \xrightarrow{\text{via } \epsilon} Lf_{\text{Mod}}^* F_Y$$

is an isomorphism by Lemma 10.1. Hence

$$\begin{aligned} \text{Hom}_{D(\text{Qco}(X))}(F\mathbb{F}, \mathbb{G}) &\cong \text{Hom}_{D(\text{Mod}(X))}(F_X F\mathbb{F}, F_X \mathbb{G}) \cong \\ \text{Hom}_{D(\text{Mod}(X))}(Lf_{\text{Mod}}^* F_Y \mathbb{F}, F_X \mathbb{G}) &\cong \text{Hom}_{D(\text{Mod}(Y))}(F_Y \mathbb{F}, Rf_*^{\text{Mod}} F_X \mathbb{G}) \cong \\ \text{Hom}_{D(\text{Mod}(Y))}(F_Y \mathbb{F}, F_Y Rf_*^{\text{Qco}} \mathbb{G}) &\cong \text{Hom}_{D(\text{Qco}(Y))}(\mathbb{F}, Rf_*^{\text{Qco}} \mathbb{G}). \end{aligned}$$

This shows that F is left adjoint to Rf_*^{Qco} .

3 Take the conjugate of **2**. □

10.5 Remark. By the lemma, we know that there is no difference between Rf_*^{Qco} on $D(\text{Qco}(X))$ and Rf_*^{Mod} on $D_{\text{Qco}(X)}(\text{Mod}(X))$ when we consider quasi-compact separated schemes. We are to consider the latter.

(10.6) Let $\mathcal{C}$ be an additive category, and $c \in \mathcal{C}$. We say that c is a compact object, if for any small family of objects $(t_\lambda)_{\lambda \in \Lambda}$ of $\mathcal{C}$ such that the coproduct (direct sum) $\bigoplus_{\lambda \in \Lambda} t_\lambda$ exists, the canonical map

$$\bigoplus_{\lambda} \text{Hom}_{\mathcal{C}}(c, t_\lambda) \rightarrow \text{Hom}_{\mathcal{C}}(c, \bigoplus_{\lambda} t_\lambda)$$

is an isomorphism.

A triangulated category $\mathcal{T}$ is said to be compactly generated, if $\mathcal{T}$ has small coproducts, and there is a small set C of compact objects of $\mathcal{T}$ such that $\text{Hom}_{\mathcal{T}}(c, t) = 0$ for all $c \in C$ implies $t = 0$. The following was proved by A. Neeman [30].

10.7 Theorem. *Let $\mathcal{S}$ be a compactly generated triangulated category, $\mathcal{T}$ any triangulated category, and $F : \mathcal{S} \rightarrow \mathcal{T}$ a triangulated functor. Suppose that F preserves coproducts, that is to say, for any small family of objects (s_λ) of $\mathcal{S}$, the canonical maps $F(s_\lambda) \rightarrow F(\bigoplus_{\lambda} s_\lambda)$ make $F(\bigoplus_{\lambda} s_\lambda)$ the coproduct of $F(s_\lambda)$. Then F has a right adjoint $G : \mathcal{T} \rightarrow \mathcal{S}$.*

Note that a right adjoint G of a triangulated functor F is triangulated. Note also that, if both $\mathcal{S}$ and $\mathcal{T}$ have t -structures and F is way-out left, then G is way-out right.

The following was also proved by A. Neeman [30].

10.8 Theorem. *Let X be a quasi-compact separated scheme. Then $c \in D(\text{Qco}(X))$ is a compact object if and only if c is isomorphic to a perfect complex, where we say that $\mathbb{C} \in C(\text{Qco}(X))$ is perfect if $\mathbb{C}$ is bounded, and each term of $\mathbb{C}$ is locally free of finite rank. Moreover, $D(\text{Qco}(X))$ is compactly generated.*

By the theorem, $D_{\text{Qco}(X)}(\text{Mod}(X))$ is also compactly generated.

10.9 Lemma. *Let $f : X \rightarrow Y$ be a quasi-compact separated morphism of schemes. Then $Rf_*^{\text{Mod}} : D_{\text{Qco}(X)}(\text{Mod}(X)) \rightarrow D(\text{Mod}(Y))$ preserves coproducts.*

Proof. See [30] or [23]. □

(10.10) Let $f : X \rightarrow Y$ be a morphism between quasi-compact separated schemes. Then by Theorem 10.7, Theorem 10.8 and Lemma 10.9, there is a right adjoint

$$f^\times : D(\text{Mod}(Y)) \rightarrow D_{\text{Qco}(X)}(\text{Mod}(X))$$

of Rf_*^{Mod} .

By restriction, $f^\times : D_{\text{Qco}(Y)}(\text{Mod}(Y)) \rightarrow D_{\text{Qco}(X)}(\text{Mod}(X))$ is a right adjoint of $Rf_*^{\text{Mod}} : D_{\text{Qco}(X)}(\text{Mod}(X)) \rightarrow D_{\text{Qco}(Y)}(\text{Qco}(Y))$. By [23, (3.6.8.2)], $(R(?)_*, (?)^{\times})$ is an adjoint pair of Δ -pseudofunctors on the category of quasi-compact separated schemes.

11 The Composition of two pseudofunctors

11.1 Definition. We say that $\mathcal{C} = (\mathcal{A}, \mathcal{F}, \mathcal{P}, \mathcal{I}, \mathcal{D}, \mathcal{D}^+, (?)^{\#}, (?)^{\flat}, \zeta)$ is a *composition data of contravariant pseudofunctors* if the following sixteen conditions are satisfied:

- 1 $\mathcal{A}$ is a category with fiber products.
- 2 $\mathcal{P}$ and $\mathcal{I}$ are classes of morphisms of $\mathcal{A}$.
- 3 Any isomorphism in $\mathcal{A}$ is in $\mathcal{P} \cap \mathcal{I}$.
- 4 The composite of two morphisms in $\mathcal{P}$ (resp. $\mathcal{I}$) is again a morphism in $\mathcal{P}$ (resp. $\mathcal{I}$).
- 5 A base change of a morphism in $\mathcal{P}$ is again a morphism in $\mathcal{P}$.
- 6 If f is a morphism in $\mathcal{A}$, $g \in \mathcal{P}$, and $g \circ f$ is defined and a morphism in $\mathcal{P}$, then $f \in \mathcal{P}$.
- 7 Any $f \in \text{Mor}(\mathcal{A})$ admits a factorization $f = pi$ such that $p \in \mathcal{P}$ and $i \in \mathcal{I}$.

Before we state the remaining conditions, we give some definitions for convenience.

- i Let $\mathcal{C}$ be a class of morphisms in $\mathcal{A}$ containing all identity maps and closed under compositions. We define $\mathcal{A}_{\mathcal{C}}$ by $\text{ob}(\mathcal{A}_{\mathcal{C}}) := \text{ob}(\mathcal{A})$ and $\text{Mor}(\mathcal{A}_{\mathcal{C}}) := \mathcal{C}$. In particular, the subcategories $\mathcal{A}_{\mathcal{P}}$ and $\mathcal{A}_{\mathcal{I}}$ of $\mathcal{A}$ are defined.

ii We call a commutative diagram of the form $p \circ i = i' \circ p'$ with $p, p' \in \mathcal{P}$ and $i, i' \in \mathcal{I}$ a *pi-square*. We denote the set of all pi-squares by Π .

8 $\mathcal{D} = (\mathcal{D}(X))_{X \in \text{ob}(\mathcal{A})}$ is a family of categories.

9 $(?)^\#$ is a contravariant pseudofunctor on $\mathcal{A}_{\mathcal{P}}$, $(?)^\flat$ is a contravariant pseudofunctor on $\mathcal{A}_{\mathcal{I}}$, and we have $X^\# = X^\flat = \mathcal{D}(X)$ for each $X \in \text{ob}(\mathcal{A})$.

10 $\zeta = (\zeta(\sigma))_{\sigma=(pi=jq) \in \Pi}$ is a family of natural transformations

$$\zeta(\sigma) : i^\flat p^\# \rightarrow q^\# j^\flat.$$

11 If

$$\begin{array}{ccccc} U_1 & \xrightarrow{j_1} & V_1 & \xrightarrow{i_1} & X_1 \\ \downarrow p_U \sigma' & & \downarrow p_V \sigma & & \downarrow p_X \\ U & \xrightarrow{j} & V & \xrightarrow{i} & X \end{array}$$

is a commutative diagram in $\mathcal{A}$ such that $\sigma, \sigma' \in \Pi$, then the composite map

$$(i_1 j_1)^\flat p_X^\# \cong j_1^\flat i_1^\flat p_X^\# \xrightarrow{\text{via } \zeta(\sigma)} j_1^\flat p_V^\# i^\flat \xrightarrow{\text{via } \zeta(\sigma')} p_U^\# j^\flat i^\flat \cong p_U^\# (ij)^\flat$$

agrees with $\zeta(\sigma'\sigma)$, where $\sigma'\sigma$ is the square $p_X(i_1 j_1) = (ij)p_U$.

12 If

$$\begin{array}{ccccc} U_2 & \xrightarrow{i_2} & X_2 & & \\ \downarrow q_U \sigma' & & \downarrow q_X & & \\ U_1 & \xrightarrow{i_1} & X_1 & & \\ \downarrow p_U \sigma & & \downarrow p_X & & \\ U & \xrightarrow{i} & X & & \end{array}$$

is a commutative diagram in $\mathcal{A}$ such that $\sigma, \sigma' \in \Pi$, then the composite

$$i_2^\flat (p_X q_X)^\# \cong i_2^\flat q_X^\# p_X^\# \xrightarrow{\text{via } \zeta(\sigma')} q_U^\# i_1^\flat p_X^\# \xrightarrow{\text{via } \zeta(\sigma)} q_U^\# p_U^\# i^\flat \cong (p_U q_U)^\# i^\flat$$

agrees with $\zeta(((p_X q_X)i_2 = i(p_U q_U)))$.

13 $\mathcal{F}$ is a subcategory of $\mathcal{A}$, and any isomorphism in $\mathcal{A}$ between objects of $\mathcal{F}$ is in $\text{Mor}(\mathcal{F})$.

14 $\mathcal{D}^+ = (\mathcal{D}^+(X))_{X \in \text{ob}(\mathcal{F})}$ is a family of categories such that $\mathcal{D}^+(X)$ is a full subcategory of $\mathcal{D}(X)$ for each $X \in \text{ob}(\mathcal{F})$.

15 If $f : X \rightarrow Y$ is a morphism in $\mathcal{F}$, $f = p \circ i$, $p \in \mathcal{P}$ and $i \in \mathcal{I}$, then we have $i^\flat p^\sharp(\mathcal{D}^+(Y)) \subset \mathcal{D}^+(X)$.

16 If

$$\begin{array}{ccccc} V & \xrightarrow{j} & U_1 & \xrightarrow{i_1} & X_1 \\ & & \downarrow p_U \sigma & & \downarrow p_X \\ & & U & \xrightarrow{i} & X \xrightarrow{q} Y \end{array}$$

is a diagram in $\mathcal{A}$ such that $\sigma \in \Pi$, $V, U, Y \in \text{ob}(\mathcal{F})$, $p_U j \in \text{Mor}(\mathcal{F})$ and $qi \in \text{Mor}(\mathcal{F})$, then

$$j^\flat \zeta(\sigma) q^\sharp : j^\flat i_1^\flat p_X^\sharp q^\sharp \rightarrow j^\flat p_U^\sharp i^\flat q^\sharp$$

is an isomorphism between functors from $\mathcal{D}^+(Y)$ to $\mathcal{D}^+(V)$.

(11.2) Let $\mathcal{C} = (\mathcal{A}, \mathcal{F}, \mathcal{P}, \mathcal{I}, \mathcal{D}, \mathcal{D}^+, (?)^\sharp, (?)^\flat, \zeta)$ be a composition data of contravariant pseudofunctors. We call a commutative diagram of the form $f = pi$ with $p \in \mathcal{P}$, $i \in \mathcal{I}$ and $f \in \text{Mor}(\mathcal{A})$ a *compactification*. We call a commutative diagram of the form $pi = qj$ with $p, q \in \mathcal{P}$, $i, j \in \mathcal{I}$ and $pi = qj \in \text{Mor}(\mathcal{F})$ an *independence diagram*.

11.3 Lemma. *Let*

$$\begin{array}{ccccc} U & \xrightarrow{i_1} & X_1 \\ \downarrow i & \tau & \downarrow p_1 \\ X & \xrightarrow{p} & Y \end{array}$$

be an independence diagram. Then the following hold:

1 *There is a diagram of the form*

$$\begin{array}{ccccc} U & \xrightarrow{j} & Z & \xrightarrow{q_1} & X_1 \\ & & \downarrow q & & \downarrow p_1 \\ & & X & \xrightarrow{p} & Y \end{array}$$

such that $qj = i$, $q_1 j = i_1$, $pq = p_1 q_1$, $q, q_1 \in \mathcal{P}$ and $j \in \mathcal{I}$.

2 $\zeta(qj = i1_U)p^\sharp : j^\flat q^\sharp p^\sharp \rightarrow i^\flat p^\sharp$ *is an isomorphism between functors from $\mathcal{D}^+(Y)$ to $\mathcal{D}^+(U)$.*

3 $\zeta(q_1 j_1 = i_1 1_U) p_1^\#$ is also an isomorphism between functors from $\mathcal{D}^+(Y)$ to $\mathcal{D}^+(U)$.

4 The composite isomorphism

$$\rho(\tau) : i^\flat p^\# \xrightarrow{(\zeta(qj=i_1 1_U) p^\#)^{-1}} j^\flat q^\# p^\# \cong j^\flat q_1^\# p_1^\# \xrightarrow{\zeta(q_1 j_1 = i_1 1_U) p_1^\#} i_1^\flat p_1^\#$$

(between functors defined over $\mathcal{D}^+(Y)$, not over $\mathcal{D}(Y)$) depends only on τ .

5 If $\tau' = (p_1 i_1 = p_2 i_2)$ is an independence diagram, then we have

$$\rho(\tau') \circ \rho(\tau) = \rho(pi = p_2 i_2).$$

The proof is left to the reader. We call $\rho(\tau)$ the independence isomorphism of τ .

(11.4) Any $f \in \text{Mor}(\mathcal{A})$ has a compactification by assumption. We fix a family of compactifications $\mathcal{T} := (\tau(f) : (f = p(f) \circ i(f)))_{f \in \text{Mor}(\mathcal{A})}$ so that $p(1_X) = i(1_X) = 1_X$ for $X \in \text{ob}(\mathcal{A})$. We denote the target of $i(f)$, which agrees with the source of $p(f)$ by $M(f)$.

For $X \in \text{ob}(\mathcal{F})$, we define $X^! := \mathcal{D}^+(X)$. For a morphism $f : X \rightarrow Y$ in $\mathcal{F}$, we define $f^! := i(f)^\flat p(f)^\#$, which is a functor from $X^!$ to $Y^!$ by assumption.

Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in $\mathcal{F}$. Set $q := p(i(g)p(f))$ and $j := i(i(g)p(f))$. Then by **16** in Definition 11.1 and Lemma 11.3, the composite map

$$\begin{aligned} (gf)^! &= i(gf)^\flat p(gf)^\# \xrightarrow{\rho(p(gf)i(gf)=(p(g)q)(ji(f)))} (j \circ i(f))^\flat (p(g) \circ q)^\# \\ &\cong i(f)^\flat j^\flat q^\# p(g)^\# \xrightarrow{i(f)^\flat \zeta(qj=i(g)p(f))p(g)^\#} i(f)^\flat p(f)^\# i(g)^\flat p(g)^\# = f^! g^! \end{aligned}$$

is an isomorphism. We define $d_{f,g} : f^! g^! \rightarrow (gf)^!$ to be the inverse of this composite.

11.5 Lemma. *The definition of $d_{f,g}$ is independent of choice of q and j above.*

The proof of the lemma is left to the reader.

11.6 Proposition. *Let the notation be as above.*

- 1 $(?)^!$ and $(d_{f,g})$ form a contravariant pseudofunctor on $\mathcal{F}$.
- 2 For $j \in \mathcal{I} \cap \text{Mor}(\mathcal{F})$, define $\rho : j^! \rightarrow j^b$ to be $\rho(p(j)i(j)) = j\text{id}$. Then $\rho : (?)^! \rightarrow (?)^b$ is an isomorphism of pseudofunctors on $\mathcal{A}_{\mathcal{I}} \cap \mathcal{F}$.
- 3 For $q \in \mathcal{P} \cap \text{Mor}(\mathcal{F})$, define $\rho : q^! \rightarrow q^\#$ to be $\rho(p(q)i(q)) = \text{id}q$. Then $\rho : (?)^! \rightarrow (?)^\#$ is an isomorphism of pseudofunctors on $\mathcal{A}_{\mathcal{P}} \cap \mathcal{F}$.
- 4 Let us take another family of compactifications $(f = p_1(f)i_1(f))_{f \in \text{Mor}(\mathcal{F})}$, and let $(?)^*$ be the resulting pseudofunctor defined by $f^* = i_1(f)^b p_1(f)^\#$. Then $\rho : f^! \rightarrow f^*$ induces an isomorphism of pseudofunctors $(?)^! \cong (?)^*$.

The proof is left to the reader. We call $(?)^!$ the composition of $(?)^\#$ and $(?)^b$. The composition is uniquely defined up to isomorphisms of pseudofunctors on $\mathcal{F}$. The discussion above has an obvious triangulated version. Composition data of contravariant triangulated pseudofunctors are defined appropriately, and the composition of two triangulated pseudofunctors is obtained as a triangulated pseudofunctor.

(11.7) Let S be a separated noetherian scheme. Let $\mathcal{A}$ be the category whose objects are separated noetherian S -schemes and morphisms are morphisms of finite type. Set $\mathcal{F} = \mathcal{A}$. Let $\mathcal{I}$ be the class of open immersions. Let $\mathcal{P}$ be the class of proper morphisms. Set $\mathcal{D}(X) = \mathcal{D}^+(X) = D_{\text{Qco}(X)}^+(\text{Mod}(X))$ for $X \in \mathcal{A}$. Let $(?)^b := (?)^*$, the derived inverse image pseudofunctor for morphisms in $\mathcal{I}$, where $X^b := \mathcal{D}(X)$. Let $(?)^\# := (?)^\times$, the twisted inverse pseudofunctor for morphisms in $\mathcal{P}$, where $X^\# := \mathcal{D}(X)$ again. Note that the left adjoint $R(?)_*$ is way-out left for morphisms in $\mathcal{P}$ so that $(?)^\times$ is way-out right, and $(?)^\#$ is well-defined. The conditions **1–9** in Definition 11.1 hold. Note that **7** is nothing but Nagata's compactification theorem [29]. The conditions **13–15** are trivial.

Let $\sigma_0 : pi = jq$ be a pi-diagram, which is also a fiber square. Then the canonical map

$$\theta : j^*(Rp_*) \rightarrow (Rq_*)i^*$$

is an isomorphism of triangulated functors, see [23, (3.9.5)]. Hence, taking the inverse of the conjugate, we have an isomorphism

$$\xi = \xi(\sigma_0) : (Ri_*)q^\times \cong p^\times(Rj_*). \quad (11.8)$$

So we have a morphism of triangulated functors

$$\zeta_0(\sigma_0) : i^* p^\times \xrightarrow{\text{via } u} i^* p^\times (Rj_*) j^* \xrightarrow{\text{via } \xi^{-1}} i^* (Ri_*) q^\times j^* \xrightarrow{\text{via } \varepsilon} q^\times j^*,$$

which is an *isomorphism*, see [36]. The statements corresponding to **10–12** only for *fiber square pi-diagrams*, is readily proved.

In particular, for a closed open immersion $\eta : U \rightarrow X$, we have an isomorphism

$$v(\eta) := \zeta_0(\pi 1_U = 1_U \pi)^{-1} : \eta^* \rightarrow \eta^\times.$$

Let $\sigma = (pi = qj)$ be an arbitrary pi-diagram. Let j_1 be the base change of j by p , and let p_1 be the base change of p by j . Let η be the unique morphism such that $q = p_1 \eta$ and $i = j_1 \eta$. Note that η is a closed open immersion. Define $\zeta(\sigma)$ to be the composite isomorphism

$$i^* p^\times \cong \eta^* j_1^* p^\times \xrightarrow{\text{via } \zeta_0} \eta^* p_1^\times j^* \xrightarrow{\text{via } v(\eta)} \eta^\times p_1^\times j^* \cong q^\times j^*.$$

Now the proof of conditions **10–12** consists in diagram chasing arguments, while **16** is trivial, since $\zeta(\sigma)$ is always an isomorphism. Thus the twisted inverse triangulated pseudofunctor $(?)^!$ on $\mathcal{A}$ is defined to be the composite of $(?)^\times$ and $(?)^*$.

12 The right adjoint of the derived direct image functor of a morphism of diagrams

Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$.

12.1 Lemma. *Assume that $X_\bullet$ is quasi-compact separated. That is, X_i is quasi-compact separated for each $i \in I$. Let C_i be a small set of compact generators of $D(\text{Qco}(X_i))$. Then*

$$C := \{LL_i c \mid i \in I, c \in C_i\}$$

is a small set of compact generators of $D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$. In particular, the category $D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$ is compactly generated.

Proof. Let $t \in D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$ and assume that

$$\text{Hom}_{D(\text{Mod}(X_\bullet))}(LL_i c, t) \cong \text{Hom}_{D(\text{Mod}(X_i))}(c, t_i) = 0$$

for any $i \in I$ and any $c \in C_i$. Then, $t_i = 0$ for all i . This shows $t = 0$. It is easy to see that $LL_i c$ is compact, and C is small. So C is a small set of compact generators. As $D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$ has coproducts, it is compactly generated. $\square$

12.2 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a quasi-compact separated morphism in $\mathcal{P}(I, \text{Sch}/S)$. Then*

$$R(f_\bullet)_* : D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet)) \rightarrow D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$$

preserves coproducts.

Proof. Let (t_λ) be a small family of objects in $D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$, and consider the canonical map

$$\bigoplus_{\lambda} R(f_\bullet)_* t_\lambda \rightarrow R(f_\bullet)_* \left(\bigoplus_{\lambda} t_\lambda \right).$$

For each $i \in I$, apply $(?)_i$ to the map. As $(?)_i$ obviously preserves coproducts and we have a canonical isomorphism

$$(?)_i R(f_\bullet)_* \cong R(f_i)_* (?)_i,$$

the result is the canonical map

$$\bigoplus_{\lambda} R(f_i)_* (t_\lambda)_i \rightarrow R(f_i)_* \left(\bigoplus_{\lambda} (t_\lambda)_i \right),$$

which is an isomorphism by Lemma 10.9. Hence, $R(f_\bullet)_*$ preserves coproducts. $\square$

By Theorem 10.7, we have:

12.3 Theorem. *Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/S)$. If $X_\bullet$ and $Y_\bullet$ are quasi-compact separated, then*

$$R(f_\bullet)_* : D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet)) \rightarrow D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$$

has a right adjoint $f_\bullet^\times$.

12.4 Lemma. *Let S be a scheme, I a small category, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y'_\bullet \rightarrow Y_\bullet$ be morphisms in $\mathcal{P}(I, \text{Sch}/S)$. Set $X'_\bullet := Y'_\bullet \times_{Y_\bullet} X_\bullet$. Let $f'_\bullet : X'_\bullet \rightarrow Y'_\bullet$ be the first projection, and $g'_\bullet : X'_\bullet \rightarrow X_\bullet$ be the second*

projection. Assume that $f_\bullet$ is quasi-compact separated, and $g_\bullet$ is flat. Then the canonical map

$$\theta(g_\bullet, f_\bullet) : (g_\bullet)^* R(f_\bullet)_* \rightarrow R(f'_\bullet)_* (g'_\bullet)^*$$

is an isomorphism of functors from $D_{\text{LQco}}(X_\bullet)(\text{Mod}(X_\bullet))$ to $D_{\text{LQco}}(Y'_\bullet)(\text{Mod}(Y'_\bullet))$.

Proof. It suffices to show that for each $i \in I$,

$$(\?)_i \theta : (\?)_i (g_\bullet)^* R(f_\bullet)_* \rightarrow (\?)_i R(f'_\bullet)_* (g'_\bullet)^*$$

is an isomorphism. By Lemma 9.7, **2**, **3**, it is easy to verify that the diagram

$$\begin{array}{ccccc} g_i^* R(f_i)_* (\?)_i & \xrightarrow{c^{-1}} & g_i^* (\?)_i R(f_\bullet)_* & \xrightarrow{\theta} & (\?)_i g_\bullet^* R(f_\bullet)_* \\ \downarrow \theta(g_i, f_i) & & & & \downarrow (\?)_i \theta(g_\bullet, f_\bullet) \\ R(f'_i)_* (g'_i)^* (\?)_i & \xrightarrow{\theta} & R(f'_i)_* (\?)_i (g'_\bullet)^* & \xrightarrow{c^{-1}} & (\?)_i R(f'_\bullet)_* (g'_\bullet)^* \end{array} \quad (12.5)$$

is commutative. As the horizontal maps and $\theta(g_i, f_i)$ are isomorphisms by [23, (3.9.5)], $(\?)_i \theta(g_\bullet, f_\bullet)$ is also an isomorphism. $\square$

13 Bökstedt–Neeman resolutions and hyper-Ext sheaves

(13.1) Let $\mathcal{T}$ be a triangulated category with small direct products. Note that a direct product of distinguished triangles is again a distinguished triangle.

Let

$$\cdots \rightarrow t_3 \xrightarrow{s_3} t_2 \xrightarrow{s_2} t_1 \quad (13.2)$$

be a sequence of morphisms in $\mathcal{T}$. We define $d : \prod_{i \geq 1} t_i \rightarrow \prod_{i \geq 1} t_i$ by $p_i \circ d = p_i - s_{i+1} \circ p_{i+1}$, where $p_i : \prod_{i \geq 1} t_i \rightarrow t_i$ is the projection. Consider a distinguished triangle of the form

$$M \xrightarrow{m} \prod_{i \geq 1} t_i \xrightarrow{p} \prod_{i \geq 1} t_i \xrightarrow{q} \Sigma M,$$

where Σ denotes the suspension.

We call M , which is determined uniquely up to isomorphisms, the homotopy limit of (13.2) and denote it by $\text{holim } t_i$.

(13.3) Dually, homotopy colimit is defined and denoted by hocolim , if $\mathcal{T}$ has small coproducts.

(13.4) Let $\mathcal{A}$ be an abelian category which satisfies (AB3*). Let $(\mathbb{F}_\lambda)_{\lambda \in \Lambda}$ be a small family of objects in $K(\mathcal{A})$. Then for any $\mathbb{G} \in K(\mathcal{A})$, we have that

$$\begin{aligned} \text{Hom}_{K(\mathcal{A})}(\mathbb{G}, \prod_{\lambda} \mathbb{F}_\lambda) &= H^0(\text{Hom}_{\mathcal{A}}^{\bullet}(\mathbb{G}, \prod_{\lambda} \mathbb{F}_\lambda)) \cong H^0(\prod_{\lambda} \text{Hom}_{\mathcal{A}}^{\bullet}(\mathbb{G}, \mathbb{F}_\lambda)) \\ &\cong \prod_{\lambda} H^0(\text{Hom}_{\mathcal{A}}^{\bullet}(\mathbb{G}, \mathbb{F}_\lambda)) = \prod_{\lambda} \text{Hom}_{K(\mathcal{A})}(\mathbb{G}, \mathbb{F}_\lambda). \end{aligned}$$

That is, the direct product $\prod_{\lambda} \mathbb{F}_\lambda$ in $C(\mathcal{A})$ is also a direct product in $K(\mathcal{A})$.

(13.5) Let $\mathcal{A}$ be a Grothendieck abelian category which satisfies (AB3*), and (t_λ) a small family of objects of $D(\mathcal{A})$. Let $(\mathbb{F}_\lambda)$ be a family of K -injective objects of $K(\mathcal{A})$ such that $\mathbb{F}_\lambda$ represents t_λ for each λ . Then $Q(\prod_{\lambda} \mathbb{F}_\lambda)$ is a direct product of t_λ in $D(\mathcal{A})$. Hence $D(\mathcal{A})$ has small products.

13.6 Lemma. *Let I be a small category, S be a scheme, and let $X_{\bullet} \in \mathcal{P}(I, \text{Sch}/S)$. Let $\mathbb{F}$ be an object of $C(\text{Mod}(X_{\bullet}))$. Assume that $\mathbb{F}$ has locally quasi-coherent cohomology groups. Then the following hold.*

1 *Let $\mathfrak{I}$ denote the full subcategory of $C(\text{Mod}(X_{\bullet}))$ consisting of bounded below complexes of injective objects of $\text{Mod}(X_{\bullet})$ with locally quasi-coherent cohomology groups. There is an $\mathfrak{I}$ -special inverse system $(I_n)_{n \in \mathbb{N}}$ with the index set $\mathbb{N}$ and an inverse system of chain maps $(f_n : \tau^{\geq -n} \mathbb{F} \rightarrow I_n)$ such that*

- i** f_n is a quasi-isomorphism for any $n \in \mathbb{N}$.
- ii** $(I_n)_i = 0$ for $i \leq -n - 1$.

2 *If (I_n) and (f_n) are as in 1, then the following hold.*

- i** *For each $i \in \mathbb{Z}$, the canonical map $H^i(I_n) \rightarrow H^i(\varprojlim I_n)$ is an isomorphism for $n \geq \max(1, -i)$, where the projective limit is taken in the category $C(\text{Mod}(X_{\bullet}))$, and $H^i(?)$ denotes the i th cohomology sheaf of a complex of sheaves.*
- ii** $\varprojlim f_n : \mathbb{F} \rightarrow \varprojlim I_n$ is a quasi-isomorphism.
- iii** *The projective limit $\varprojlim I_n$, viewed as an object of $K(\text{Mod}(X))$, is the homotopy limit of (I_n) .*

iv $\varprojlim I_n$ is K -injective.

3 $Q\mathbb{F}$ is the projective limit of the projective system $(\tau^{\geq -n}Q\mathbb{F})$ in the category $D(\text{Mod}(X_\bullet))$.

Proof. The assertion 1 is [34, (3.7)].

We prove 2, i. Let $j \in \text{ob}(I)$ and U an affine open subset of X . Then for any $n \geq 1$, I_n^i and $H^i(I_n)$ are $\Gamma((j, U), ?)$ -acyclic for each $i \in \mathbb{Z}$. As I_n is bounded below, each $Z^i(I_n)$ and $B^i(I_n)$ are also $\Gamma((j, U), ?)$ -acyclic, and the sequence

$$0 \rightarrow \Gamma((j, U), Z^i(I_n)) \rightarrow \Gamma((j, U), I_n^i) \rightarrow \Gamma((j, U), B^{i+1}(I_n)) \rightarrow 0 \quad (13.7)$$

and

$$0 \rightarrow \Gamma((j, U), B^i(I_n)) \rightarrow \Gamma((j, U), Z^i(I_n)) \rightarrow \Gamma((j, U), H^i(I_n)) \rightarrow 0 \quad (13.8)$$

are exact for each i , as can be seen easily, where B^i and Z^i respectively denote the i th coboundary and the cocycle sheaves.

In particular, the inverse system $(\Gamma((j, U), B^i(I_n)))$ is a Mittag-Leffler inverse system of abelian groups by (13.7), since $(\Gamma((j, U), I_n^i))$ is. On the other hand, as we have $H^i(I_n) \cong H^i(\mathbb{F})$ for $n \geq \max(1, -i)$, the inverse system $(\Gamma((j, U), H^i(I_n)))$ stabilizes, and hence we have $(\Gamma((j, U), Z^i(I_n)))$ is also Mittag-Leffler.

Passing through the projective limit,

$$0 \rightarrow \Gamma((j, U), Z^i(\varprojlim I_n)) \rightarrow \Gamma((j, U), \varprojlim I_n) \rightarrow \Gamma((j, U), \varprojlim B^{i+1}(I_n)) \rightarrow 0$$

is exact. Hence, the canonical map $B^i(\varprojlim I_n) \rightarrow \varprojlim B^i(I_n)$ is an isomorphism, since (j, U) with U an affine open subset of X_j generates the topology of $\text{Zar}(X_\bullet)$.

Taking the projective limit of (13.8), we have

$$0 \rightarrow \Gamma((j, U), B^i(\varprojlim I_n)) \rightarrow \Gamma((j, U), Z^i(\varprojlim I_n)) \rightarrow \Gamma((j, U), \varprojlim H^i(I_n)) \rightarrow 0$$

is an exact sequence for any affine open subset U of X .

Hence, the canonical maps

$$\Gamma((j, U), H^i(I_n)) \cong \Gamma((j, U), \varprojlim H^i(I_n)) \leftarrow \Gamma((j, U), H^i(\varprojlim I_n))$$

are all isomorphisms for $n \geq \max(1, -i)$, and we have $H^i(I_n) \cong H^i(\varprojlim I_n)$ for $n \geq \max(1, -i)$.

The assertion **ii** is now trivial.

The assertion **iii** is now a consequence of [6, (2.3.1)] (one can work at the presheaf level where we have the (AB4*) property). The assertion **iv** is now obvious. $\square$

Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$.

13.9 Lemma. *Assume that $X_\bullet$ has flat arrows. Let J be a subcategory of I , and let $\mathbb{F} \in D_{\text{EM}(X_\bullet)}(\text{Mod}(X_\bullet))$ and $\mathbb{G} \in D(\text{Mod}(X_\bullet))$. Then the canonical map*

$$H_J : (?)_J R\text{Hom}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, \mathbb{G}) \rightarrow R\text{Hom}_{\text{Mod}(X_J)}^\bullet(\mathbb{F}_J, \mathbb{G}_J)$$

is an isomorphism.

Proof. First, we prove the lemma for the case where $\mathbb{F}$ has bounded above cohomology groups.

By Lemma 2.10, we may assume that $J = i$ for an object i of I . By the way-out lemma [15, Proposition I.7.1], we may assume that $\mathbb{F}$ is a single equivariant $\mathcal{O}_{X_\bullet}$ -module, and $\mathbb{G}$ is a single injective object of $\text{Mod}(X_\bullet)$.

As $X_\bullet$ has flat arrows, the restriction $(?)_i$ has an exact left adjoint L_i . It follows that $\mathbb{G}_i$ is an injective $\mathcal{O}_{X_i}$ -module. The assertion is now clear by Lemma 7.28.

Now consider the general case. As the functors in question on $\mathbb{F}$ changes coproducts to products and a direct sum of complexes with equivariant cohomology has equivariant cohomology, the map in question is an isomorphism if $\mathbb{F}$ is a direct sum of objects in $D_{\text{EM}(X_\bullet)}^-(X_\bullet)$. In particular, the map is an isomorphism if $\mathbb{F}$ is a homotopy colimit of objects of $D_{\text{EM}(X_\bullet)}^-(X_\bullet)$. As any object $\mathbb{F}$ of $D_{\text{EM}(X_\bullet)}(X_\bullet)$ is the homotopy colimit of $(\tau^{\leq n}\mathbb{F})$, we are done. $\square$

13.10 Lemma. *Assume that $X_\bullet$ has flat arrows. Let J be a subcategory of I , and let $\mathbb{F} \in D_{\text{EM}(X_\bullet)}(\text{Mod}(X_\bullet))$ and $\mathbb{G} \in D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$. Then the canonical map*

$$H_J : (?)_J R\text{Hom}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, \mathbb{G}) \rightarrow R\text{Hom}_{\text{Mod}(X_J)}^\bullet(\mathbb{F}_J, \mathbb{G}_J)$$

is an isomorphism.

Proof. By Lemma 13.6, $\mathbb{G}$ is the homotopy limit of objects of $D^+(\text{Mod}(X_\bullet))$. The functors in question on $\mathbb{G}$ preserve direct products and triangles. In particular, they preserve homotopy limits. The assertion follows immediately by Lemma 13.9. $\square$

13.11 Lemma. *Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$ has flat arrows and is locally noetherian. Let $\mathbb{F} \in D_{\text{Coh}(X_\bullet)}^-(\text{Mod}(X_\bullet))$ and $\mathbb{G} \in D_{\text{LQco}(X_\bullet)}^+(\text{Mod}(X_\bullet))$. Then $\underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^i(\mathbb{F}, \mathbb{G})$ is locally quasi-coherent for $i \in \mathbb{Z}$. If moreover $\mathbb{G}$ has quasi-coherent cohomology groups, then $\underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^i(\mathbb{F}, \mathbb{G})$ is quasi-coherent for $i \in \mathbb{Z}$.*

Proof. We prove the assertion for the local quasi-coherence. By Lemma 13.9, we may assume that $X_\bullet$ is a single scheme. This case is [15, Proposition II.3.3].

We prove the assertion for the quasi-coherence, assuming that $\mathbb{G}$ has quasi-coherent cohomology groups. By [15, Proposition I.7.3], we may assume that $\mathbb{F}$ is a single coherent sheaf, and $\mathbb{G}$ is an injective resolution of a single quasi-coherent sheaf.

As $X_\bullet$ has flat arrows and the restrictions are exact, it suffices to show that

$$\alpha_\phi : X_\phi^*(?)_i \underline{\text{Hom}}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, \mathbb{G}) \rightarrow (?)_j \underline{\text{Hom}}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, \mathbb{G})$$

is a quasi-isomorphism for any morphism $\phi : i \rightarrow j$ in I .

As X_ϕ is flat, $\alpha_\phi : X_\phi^*\mathbb{F}_i \rightarrow \mathbb{F}_j$ and $\alpha_\phi : X_\phi^*\mathbb{G}_i \rightarrow \mathbb{G}_j$ are quasi-isomorphisms. In particular, the latter is a K -injective resolution.

By the derived version of (7.29), it suffices to show that

$$P : X_\phi^* R \underline{\text{Hom}}_{\mathcal{O}_{X_i}}^\bullet(\mathbb{F}_i, \mathbb{G}_i) \rightarrow R \underline{\text{Hom}}_{\mathcal{O}_{X_j}}^\bullet(X_\phi^*\mathbb{F}_i, X_\phi^*\mathbb{G}_i)$$

is an isomorphism. This is [15, Proposition II.5.8]. $\square$

14 Commutativity of twisted inverse with restrictions

(14.1) Let S be a scheme, I a small category, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Let J be an admissible subcategory of I . Assume that $f_\bullet$ is quasi-compact and separated. Then there is a natural map

$$\theta(J, f_\bullet) : LL_J \circ R(f_\bullet|_J)_* \rightarrow R(f_\bullet)_* \circ LL_J \quad (14.2)$$

between functors from $D_{\text{LQco}(X_\bullet|_J)}(\text{Mod}(X_\bullet|_J))$ to $D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$, see [23, (3.7.2)].

(14.3) Let S , I and $f_\bullet$ be as in (14.1). We assume that $X_\bullet$ and $Y_\bullet$ are quasi-compact $\mathbb{Z}$ -separated, so that the right adjoint functor

$$f_\bullet^\times : D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet)) \rightarrow D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$$

of

$$R(f_\bullet)_* : D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet)) \rightarrow D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$$

exists. Let J be a subcategory of I which may not be admissible.

We define the natural transformation

$$\xi(J, f_\bullet) : (?)_J \circ f_\bullet^\times \rightarrow f_J^\times \circ (?)_J$$

to be the composite

$$(?)_J f_\bullet^\times \xrightarrow{u} (f_\bullet|_J)^\times R(f_\bullet|_J)_* (?)_J f_\bullet^\times \xrightarrow{c} (f_\bullet|_J)^\times (?)_J R(f_\bullet)_* f_\bullet^\times \xrightarrow{\varepsilon} (f_\bullet|_J)^\times (?)_J.$$

By definition, ξ is the conjugate map of $\theta(J, f_\bullet)$ in (14.2) if J is admissible.

14.4 Lemma. *Let $J_2 \subset J_1 \subset I$ be subcategories of I . Let S be a scheme, $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Assume both $X_\bullet$ and $Y_\bullet$ are quasi-compact separated. Then the composite map*

$$\begin{aligned} (?)_{J_2} f_\bullet^\times &\cong (?)_{J_2} (?)_{J_1} f_\bullet^\times \xrightarrow{\text{via } \xi(J_1)} (?)_{J_2} (f_\bullet|_{J_1})^\times (?)_{J_1} \\ &\xrightarrow{\text{via } \xi(J_2)} (f_\bullet|_{J_2})^\times (?)_{J_2} (?)_{J_1} = (f_\bullet|_{J_2})^\times (?)_{J_2} \end{aligned}$$

is equal to $\xi(J_2)$.

Proof. Straightforward (and tedious) diagram drawing. $\square$

14.5 Lemma. *Let S and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be as in Lemma 14.4. Let J be a subcategory of I . Assume that $Y_\bullet$ has flat arrows and $f_\bullet$ is of fiber type. Then $\xi(J, f_\bullet) : (?)_J f_\bullet^\times \rightarrow (f_\bullet|_J)^\times (?)_J$ is an isomorphism between functors from $D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$ to $D_{\text{LQco}(X_\bullet|_J)}(\text{Mod}(X_\bullet|_J))$.*

Proof. In view of Lemma 14.4, we may assume that $J = i$ for an object i of I . Then, as i is an admissible subcategory of I and $\xi(i, f_\bullet)$ is an conjugate map of $\theta(i, f_\bullet)$, it suffices to show that $(?)_i \theta(i, f_\bullet)$ is an isomorphism for any $j \in \text{ob}(I)$. As $Y_\bullet$ has flat arrows, $L_i : \text{Mod}(Y_i) \rightarrow \text{Mod}(Y_\bullet)$ is exact. As $f_\bullet$ is of fiber type, $L_i : \text{Mod}(X_i) \rightarrow \text{Mod}(X_\bullet)$ is also exact.

By Proposition 7.18, the composite

$$(\?)_j L_i R(f_i)_* \xrightarrow{(\?)_j \theta} (\?)_j R(f_\bullet)_* L_i \xrightarrow{c} R(f_j)_* (\?)_j L_i$$

agrees with the composite

$$\begin{aligned} (\?)_j L_i R(f_i)_* &\xrightarrow{\lambda_{i,j}} \bigoplus_{\phi \in I(i,j)} X_\phi^* R(f_i)_* \xrightarrow{\oplus \theta} \bigoplus_{\phi} R(f_j)_* Y_\phi^* \\ &\xrightarrow{C} R(f_j)_* \left(\bigoplus_{\phi} Y_\phi^* \right) \xrightarrow{\lambda_{i,j}^{-1}} R(f_j)_* (\?)_j L_i, \end{aligned}$$

where C is the canonical map. By Lemma 10.9, C is an isomorphism. As $f_\bullet$ is of fiber type, $\theta : X_\phi^* R(f_i)_* \rightarrow R(f_j)_* Y_\phi^*$ is an isomorphism for each $\phi \in I(i, j)$ by [23, (3.9.5)]. Hence the second composite is an isomorphism. As the first composite is an isomorphism and c is also an isomorphism, we have that $(\?)_j \theta(i, f_\bullet)$ is an isomorphism. $\square$

(14.6) Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$ and $Y_\bullet$ are quasi-compact separated.

14.7 Lemma. *Let J be a subcategory of I . Then the following hold:*

1 *The composite map*

$$\begin{aligned} (\?)_J &\xrightarrow{(\?)_J u} (\?)_J f_\bullet^\times R(f_\bullet)_* \xrightarrow{\text{via } \xi(J, f_\bullet)} \\ &\quad (f_\bullet|_J)^\times (\?)_J R(f_\bullet)_* \xrightarrow{\text{via } c_{J, f_\bullet}} (f_\bullet|_J)^\times R(f_\bullet|_J)_* (\?)_J \end{aligned}$$

agrees with $u(\?)_J$, where $u : \text{Id} \rightarrow (f_\bullet|_J)^\times R(f_\bullet|_J)_*$ is the unit of adjunction.

2 *The composite map*

$$\begin{aligned} (\?)_J R(f_\bullet)_* f_\bullet^\times &\xrightarrow{\text{via } c_{J, f_\bullet}} R(f_\bullet|_J)_* (\?)_J f_\bullet^\times \xrightarrow{\text{via } \xi(J, f_\bullet)} \\ &\quad R(f_\bullet|_J)_* (f_\bullet|_J)^\times (\?)_J \xrightarrow{\varepsilon(\?)_J} (\?)_J \end{aligned}$$

agrees with $(\?)_J \varepsilon$.

Proof. This consists in straightforward diagram drawings. $\square$

(14.8) Let I be a small category. For $i, j \in \text{ob}(I)$, we say that $i \leq j$ if $I(i, j) \neq \emptyset$. This definition makes $\text{ob}(I)$ a pseudo-ordered set. We say that I is *ordered* if $\text{ob}(I)$ is an ordered set with the pseudo-order structure above, and $I(i, i) = \{\text{id}\}$ for $i \in I$.

14.9 Lemma. *Let I be an ordered small category. Let J_0 and J_1 be full subcategories of I , such that $\text{ob}(J_0) \cup \text{ob}(J_1) = \text{ob}(I)$, $\text{ob}(J_0) \cap \text{ob}(J_1) = \emptyset$, and $I(j_1, j_0) = \emptyset$ for $j_1 \in J_1$ and $j_0 \in J_0$. Let $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Then, we have the following.*

- 1** *The unit of adjunction $u : \text{Id}_{\text{Mod}(X_\bullet|_{J_1})} \rightarrow (?)_{J_1} \circ L_{J_1}$ is an isomorphism.*
- 2** *$(?)_{J_0} \circ L_{J_1}$ is zero.*
- 3** *L_{J_1} is exact, and J_1 is an admissible subcategory of I .*
- 4** *For $\mathcal{M} \in \text{Mod}(X_\bullet)$, $\mathcal{M}_{J_0} = 0$ if and only if $\varepsilon : L_{J_1}\mathcal{M}_{J_1} \rightarrow \mathcal{M}$ is an isomorphism.*
- 5** *The counit of adjunction $(?)_{J_0} \circ R_{J_0} \rightarrow \text{Id}_{\text{Mod}(X_\bullet|_{J_0})}$ is an isomorphism.*
- 6** *$(?)_{J_1} \circ R_{J_0}$ is zero.*
- 7** *R_{J_0} is exact and preserves local-quasi-coherence.*
- 8** *For $\mathcal{M} \in \text{Mod}(X_\bullet)$, $\mathcal{M}_{J_1} = 0$ if and only if $u : \mathcal{M} \rightarrow R_{J_0}\mathcal{M}_{J_0}$ is an isomorphism.*
- 9** *The sequence*

$$0 \rightarrow L_{J_1}(?)_{J_1} \xrightarrow{\varepsilon} \text{Id} \xrightarrow{u} R_{J_0}(?)_{J_0} \rightarrow 0$$

is semisplit exact, and induces a distinguished triangle in $D(\text{Mod}(X_\bullet))$.

Proof. **1** This is obvious by Lemma 7.11.

2 The category $I_{j_0}^{(J_1 \hookrightarrow I)}$ is empty, if $j_0 \in J_0$, since $I(j_1, j_0) = \emptyset$ if $j_1 \in J_1$ and $j_0 \in J_0$. It follows that $(?)_{j_0} \circ L_{J_1} = 0$ if $j_0 \in J_0$. Hence, $(?)_{J_0} \circ L_{J_1} = 0$.

3 This is trivial by **1, 2** and their proof.

4 The ‘if’ part is trivial by **2**. We prove the ‘only if’ part. By assumption and **2**, both $\mathcal{M}_{J_0}$ and $(?)_{J_0} L_{J_1} \mathcal{M}_{J_1}$ are zero, and $(?)_{J_0} \varepsilon$ is an isomorphism. On the other hand, $((?)_{J_1} \varepsilon)(u(?)_{J_1}) = \text{id}$, and u is an isomorphism by **1**. Hence,

$$(?)_{J_1} \varepsilon : (?)_{J_1} L_{J_1} \mathcal{M}_{J_1} \rightarrow (?)_{J_1} \mathcal{M}$$

is an isomorphism. Hence, ε is an isomorphism.

The assertions **5,6,7,8,9** are similar, and we omit the proof. $\square$

14.10 Lemma. *Let I , S , J_1 and J_0 be as in Lemma 14.9. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$ and $Y_\bullet$ are quasi-compact separated. Then we have that $\theta(J_1, f_\bullet)$ and $\xi(J_1, f_\bullet)$ is an isomorphism.*

Proof. Note that J_1 is admissible by Lemma 14.9, **3**, and hence $\theta(J_1, f_\bullet)$ and $\xi(J_1, f_\bullet)$ are defined. Since we have $\xi(J_1, f_\bullet)$ is the conjugate of $\theta(J_1, f_\bullet)$ by definition, it suffices to show that $\theta(J_1, f_\bullet)$ is an isomorphism. It suffices to show that

$$(\?)_i LL_{J_1} R(f_\bullet|_{J_1})_* \xrightarrow{(\?)_i \theta} (\?)_i R(f_\bullet)_* LL_{J_1} \cong R(f_i)_* (\?)_i LL_{J_1}$$

is an isomorphism for any $i \in I$.

If $i \in J_0$, then both hand sides are zero functors, and it is an isomorphism. On the other hand, if $i \in J_1$, then the map in question is equal to the composite isomorphism

$$(\?)_i LL_{J_1} R(f_\bullet|_{J_1})_* \cong (\?)_i R(f_\bullet|_{J_1})_* \xrightarrow{c_{i, f_\bullet|_{J_1}}} R(f_i)_* (\?)_i \cong R(f_i)_* (\?)_i LL_{J_1}$$

by Proposition 7.18. Hence $\theta(J_1, f_\bullet)$ is an isomorphism, as desired. $\square$

(14.11) Let S be a scheme, I an ordered small category, $i \in I$, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Let J_1 be a filter of $\text{ob}(I)$ such that i is a minimal element of J_1 (e.g., $[i, \infty)$), and set $\Gamma_i := L_{I, J_1} \circ R_{J_1, i}$. Then we have $(\?)_j \Gamma_i = 0$ if $j \neq i$ and $(\?)_i \Gamma_i = \text{Id}$. It follows that Γ_i preserves arbitrary limits and colimits (hence is exact). Assume that X_i is quasi-compact and separated. Then $D_{\text{Qco}(X_i)}(\text{Mod}(X_i))$ is compactly generated, and the derived functor

$$\Gamma_i : D_{\text{Qco}(X_i)}(\text{Mod}(X_i)) \rightarrow D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$$

preserves coproducts. It follows that there is a right adjoint

$$\Sigma_i : D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet)) \rightarrow D_{\text{Qco}(X_i)}(\text{Mod}(X_i)).$$

As Γ_i is obviously way-out left, we have Σ_i is way-out right.

(14.12) Let S be a scheme, I a small category, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. We define $\mathcal{D}^+(X_\bullet)$ (resp. $\mathcal{D}^-(X_\bullet)$) to be the full subcategory of $D(\text{Mod}(X_\bullet))$ consisting of $\mathbb{F} \in D(\text{Mod}(X_\bullet))$ such that $\mathbb{F}_i$ is bounded below (resp. above) and has quasi-coherent cohomology groups for each $i \in I$. For a plump full subcategory $\mathcal{A}$ of $\text{LQco}(X_\bullet)$, we denote the triangulated subcategory of $\mathcal{D}^+(X_\bullet)$ (resp. $\mathcal{D}^-(X_\bullet)$) consisting of objects all of whose cohomology groups belong to $\mathcal{A}$ by $\mathcal{D}_{\mathcal{A}}^+(X_\bullet)$ (resp. $\mathcal{D}_{\mathcal{A}}^-(X_\bullet)$).

(14.13) Let P be an ordered set. We say that P is upper Jordan-Dedekind (UJD for short) if for any $p \in P$, the subset

$$[p, \infty) := \{q \in P \mid q \geq p\}$$

is finite. We say that an ordered small category I is UJD if the ordered set $\text{ob}(I)$ is UJD, and $I(i, j)$ is finite for $i, j \in I$.

14.14 Proposition. *Let I be an ordered UJD small category. Let S be a scheme, and $g_\bullet : U_\bullet \rightarrow X_\bullet$ and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be morphisms in $\mathcal{P}(I, \text{Sch}/S)$. Assume that $Y_\bullet$ is separated noetherian which has flat arrows, $f_\bullet$ is proper, $g_\bullet$ is an open immersion such that $g_i(U_i)$ is dense in X_i for each $i \in I$, and $f_\bullet \circ g_\bullet$ is of fiber type. Then $g_\bullet$ is of fiber type, and for any $i \in I$ the composite natural map*

$$(\cdot)_i g_\bullet^* f_\bullet^\times \xrightarrow{\text{via } \theta^{-1}} g_i^*(\cdot)_i f_\bullet^\times \xrightarrow{\text{via } \xi(i)} g_i^* f_i^\times (\cdot)_i$$

is an isomorphism between functors $\mathcal{D}^+(Y_\bullet) \rightarrow D_{\text{Qco}(U_i)}(\text{Mod}(U_i))$, where $\theta : g_i^*(\cdot)_i \rightarrow (\cdot)_i g_\bullet^*$ is the canonical isomorphism.

Proof. Note that $U_\bullet$ has flat arrows, since $Y_\bullet$ has flat arrows and $f_\bullet \circ g_\bullet$ is of fiber type.

We prove that $g_\bullet$ is of fiber type. Let $\phi : i \rightarrow j$ be a morphism in I . Then, the canonical map $(U_\phi, f_j g_j) : U_j \rightarrow U_i \times_{Y_i} Y_j$ is an isomorphism by assumption. This map factors through $(U_\phi, g_j) : U_j \rightarrow U_i \times_{X_i} X_j$, and hence (U_ϕ, g_j) is a closed immersion. On the other hand, it is an image dense open immersion, as can be seen easily, and hence it is an isomorphism. So $g_\bullet$ is of fiber type.

Set $J_1 := [i, \infty)$ and $J_0 := \text{ob}(I) \setminus J_1$. By Lemma 14.10, $\xi(J_1, f_\bullet)$ is an isomorphism. By Lemma 14.4, we may replace I by J_1 , and we may assume that I is an ordered finite category, and i is a minimal element of $\text{ob}(I)$. Now we have $\mathcal{D}^+(Y_\bullet)$ agrees with $D_{\text{LQco}(Y_\bullet)}^+(\text{Mod}(Y_\bullet))$. Since we have $\text{ob}(I)$

is finite, it is easy to see that $R(f_\bullet)_*$ is way-out both. It follows that $f_\bullet^\times$ is way-out right.

It suffices to show

$$g_i^* \xi(i) : g_i^*(?)_i f_\bullet^\times \rightarrow g_i^* f_i^\times (?)_i$$

is an isomorphism of functors from $D_{\text{LQco}(Y_\bullet)}^+(\text{Mod}(Y_\bullet))$ to $D_{\text{Qco}(U_i)}^+(\text{Mod}(U_i))$. As $g_i^* R(g_i)_* \cong \text{Id}$, it suffices to show that $R(g_i)_* g_i^* \xi(i)$ is an isomorphism. This is equivalent to say that for any perfect complex $\mathbb{P} \in C(\text{Qco}(X_i))$, we have

$$\begin{aligned} R(g_i)_* g_i^* \xi(i) : \text{Hom}_{D(\text{Mod}(X_i))}(\mathbb{P}, R(g_i)_* g_i^*(?)_i f_\bullet^\times) \\ \rightarrow \text{Hom}_{D(\text{Mod}(X_i))}(\mathbb{P}, R(g_i)_* g_i^* f_i^\times (?)_i) \end{aligned}$$

is an isomorphism. By [36, Lemma 2], this is equivalent to say that the canonical map

$$\begin{aligned} \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(R(f_\bullet)_* LL_i(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \\ \rightarrow \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(LL_i R(f_i)_*(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \end{aligned}$$

induced by the conjugate $\theta(i, f_\bullet)$ of $\xi(i)$ is an isomorphism, where $\mathcal{J}$ is a defining ideal sheaf of the closed subset $X_i \setminus U_i$ in X_i .

As I is ordered and $\text{ob}(I)$ is finite, we may label

$$\text{ob}(I) = \{i = i(0), i(1), i(2), \dots\}$$

so that $I(i(s), i(t)) \neq \emptyset$ implies that $s \leq t$. Let $J(r)$ denote the full subcategory of I whose object set is $\{i(r), i(r+1), \dots\}$.

By descending induction on t , we prove that the map

$$\begin{aligned} \text{via } \theta(i, f_\bullet) : \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(L_{J(t)}(?)_{J(t)} R(f_\bullet)_* LL_i(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \\ \rightarrow \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(L_{J(t)}(?)_{J(t)} LL_i R(f_i)_*(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \end{aligned}$$

is an isomorphism. This is enough to prove the proposition, since $L_{J(1)}(?)_{J(1)} = \text{Id}$.

Since the sequence

$$0 \rightarrow L_{J(t+1)}(?)_{J(t+1)} \xrightarrow{\text{via } \varepsilon} L_{J(t)}(?)_{J(t)} \rightarrow \Gamma_{i(t)}(?)_{i(t)} \rightarrow 0$$

is an exact sequence of exact functors, it suffices to prove that the map

$$\begin{aligned} \text{via } \theta(i, f_\bullet) : \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(\Gamma_{i(t)}(?)_{i(t)} R(f_\bullet)_* L L_i(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \\ \rightarrow \varinjlim \text{Hom}_{D(\text{Mod}(Y_\bullet))}(\Gamma_{i(t)}(?)_{i(t)} L L_i R(f_i)_*(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), ?) \end{aligned}$$

is an isomorphism by induction assumption and the five lemma. By Proposition 7.18, this is equivalent to say that the map

$$\begin{aligned} \text{via } \theta : \varinjlim \text{Hom}_{D(\text{Mod}(Y_{i(t)}))}(R(f_{i(t)})_* \bigoplus_{\phi} L X_{\phi}^*(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), \Sigma_i(?)) \\ \rightarrow \varinjlim \text{Hom}_{D(\text{Mod}(Y_{i(t)}))}(\bigoplus_{\phi} Y_{\phi}^* R(f_i)_*(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n), \Sigma_i(?)) \end{aligned}$$

is an isomorphism, where the sum is taken over the *finite* set $I(i, i(t))$. It suffices to prove that the map

$$\begin{aligned} \text{via } \xi : \varinjlim \text{Hom}_{D(\text{Mod}(X_i))}(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n, R(X_{\phi})_* f_{i(t)}^{\times} \Sigma_i(?)) \\ \rightarrow \varinjlim \text{Hom}_{D(\text{Mod}(X_i))}(\mathbb{P} \otimes_{\mathcal{O}_{X_i}}^{\bullet, L} \mathcal{J}^n, f_i^{\times} R(Y_{\phi})_* \Sigma_i(?)) \end{aligned}$$

induced by the map $\xi : R(X_{\phi})_* f_{i(t)}^{\times} \rightarrow f_i^{\times} R(Y_{\phi})_*$, which is conjugate to

$$\theta : Y_{\phi}^* R(f_i)_* \rightarrow R(f_{i(t)})_* L X_{\phi}^*,$$

is an isomorphism for $\phi \in I(i, i(t))$. Since both $R(X_{\phi})_* f_{i(t)}^{\times} \Sigma_i$ and $f_i^{\times} R(Y_{\phi})_* \Sigma_i$ are way-out right, it suffices to show the canonical map

$$g_i^* \xi(Y_{\phi} f_{i(t)} = f_i X_{\phi}) : g_i^* R(X_{\phi})_* f_{i(t)}^{\times} \rightarrow g_i^* f_i^{\times} R(Y_{\phi})_*$$

is an isomorphism between functors from $D_{\text{Qco}(Y_{i(t)})}^+(\text{Mod}(Y_{i(t)}))$ to $D_{\text{Qco}(U_i)}^+(\text{Mod}(U_i))$. Let $X' := X_i \times_{Y_i} Y_{i(t)}$, $p_1 : X' \rightarrow X_i$ be the first projection, $p_2 : X' \rightarrow Y_{i(t)}$ the second projection, and $\pi : X_{i(t)} \rightarrow X'$ be the map $(X_{\phi}, f_{i(t)})$. It is easy to see that $\xi(Y_{\phi} f_{i(t)} = f_i X_{\phi})$ equals the composite map

$$R(X_{\phi})_* f_{i(t)}^{\times} \cong R(p_1)_* R\pi_* \pi^{\times} p_2^{\times} \xrightarrow{\varepsilon} R(p_1)_* p_2^{\times} \cong f_i^{\times} R(Y_{\phi})_*.$$

Note that the last map is an isomorphism since Y_{ϕ} is flat. As we have

$$U_{i(t)} \rightarrow U_i \times_{Y_i} Y_{i(t)} \cong U_i \times_{X_i} X'$$

is an isomorphism and g_i is an open immersion by assumption, the canonical map

$$g_i^* R(p_1)_* \rightarrow R(U_\phi)_*(\pi \circ g_{i(t)})^*$$

is an isomorphism. So it suffices to prove that

$$(\pi \circ g_{i(t)})^* R\pi_* \pi^\times \xrightarrow{\text{via } \varepsilon} (\pi \circ g_{i(t)})^*$$

is an isomorphism.

Consider the fiber square

$$\begin{array}{ccc} U_{i(t)} & \xrightarrow{g_{i(t)}} & X_{i(t)} \\ \downarrow \text{id} & \sigma & \downarrow \pi \\ U_{i(t)} & \xrightarrow{\pi \circ g_{i(t)}} & X'. \end{array}$$

By [36], $\zeta_0(\sigma) : g_{i(t)}^* \pi^\times \rightarrow (\pi \circ g_{i(t)})^*$ is an isomorphism. By definition (11.7), ζ_0 is the composite map

$$g_{i(t)}^* \pi^\times \xrightarrow{u} \text{id}^\times R\text{id}_* g_{i(t)}^* \pi^\times \xrightarrow{\cong} \text{id}_* (\pi \circ g_{i(t)})^* R\pi_* \pi^\times \xrightarrow{\varepsilon} (\pi \circ g_{i(t)})^*.$$

Since the first and the second maps are isomorphisms, the third map is an isomorphism. This was what we wanted to prove. $\square$

14.15 Corollary. *Under the same assumption as in the proposition, we have $g_\bullet^* f_\bullet^\times(\mathcal{D}^+(Y_\bullet)) \subset \mathcal{D}^+(U_\bullet)$.*

Proof. This is because $g_i^* f_i^\times(D_{\text{Qco}(Y_i)}^+(\text{Mod}(Y_i))) \subset D_{\text{Qco}(U_i)}^+(\text{Mod}(U_i))$ for each $i \in I$.

15 Open immersion base change

(15.1) Let S be a scheme, I a small category, and

$$\begin{array}{ccc} X'_\bullet & \xrightarrow{g'_\bullet} & X_\bullet \\ \downarrow f'_\bullet & \sigma & \downarrow f_\bullet \\ Y'_\bullet & \xrightarrow{g_\bullet} & Y_\bullet \end{array}$$

a fiber square in $\mathcal{P}(I, \text{Sch}/S)$. Assume that the all diagrams of schemes in the diagram are quasi-compact separated, and $g_\bullet$ is flat. By Lemma 12.4, the canonical map

$$\theta(g_\bullet, f_\bullet) : g_\bullet^* R(f_\bullet)_* \rightarrow R(f'_\bullet)_*(g'_\bullet)^*$$

is an isomorphism of functors from $D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$ to $D_{\text{LQco}(Y'_\bullet)}(\text{Mod}(Y'_\bullet))$. We define $\zeta(\sigma) = \zeta(g_\bullet, f_\bullet)$ to be the composite map

$$\zeta(\sigma) : (g'_\bullet)^* f_\bullet^\times \xrightarrow{u} (f'_\bullet)^\times R(f'_\bullet)_* (g'_\bullet)^* f_\bullet^\times \xrightarrow{\theta^{-1}} (f'_\bullet)^\times g_\bullet^* R(f_\bullet)_* f_\bullet^\times \xrightarrow{\varepsilon} (f'_\bullet)^\times g_\bullet^*.$$

15.2 Lemma. *Let σ be as above, and J a subcategory of I . Then the diagram*

$$\begin{array}{ccccc} (?)_J (g'_\bullet)^* f_\bullet^\times & \xrightarrow{\theta^{-1}} & (g'_\bullet|_J)^* (?)_J f_\bullet^\times & \xrightarrow{\xi} & (g'_\bullet|_J)^* (f_\bullet|_J)^\times (?)_J \\ \downarrow \zeta & & & & \downarrow \zeta \\ (?)_J (f'_\bullet)^\times g_\bullet^* & \xrightarrow{\xi} & (f'_\bullet|_J)^\times (?)_J g_\bullet^* & \xrightarrow{\theta^{-1}} & (f'_\bullet|_J)^\times (g_\bullet|_J)^* (?)_J \end{array}$$

is commutative.

Proof. Follows immediately from Lemma 14.7 and the commutativity of (12.5). $\square$

15.3 Theorem. *Let S be a scheme, I an ordered UJD small category, and*

$$\begin{array}{ccccc} V_\bullet & \xrightarrow{j} & U'_\bullet & \xrightarrow{i'_\bullet} & X'_\bullet \\ & & \downarrow p_\bullet^U \quad \sigma & & \downarrow p_\bullet^X \\ & & U_\bullet & \xrightarrow{i_\bullet} & X_\bullet \xrightarrow{q_\bullet} Y_\bullet \end{array}$$

be a diagram in $\mathcal{P}(I, \text{Sch}/S)$. Assume the following.

- 1 $Y_\bullet$ is separated noetherian.
- 2 $j_\bullet$, $i'_\bullet$ and $i_\bullet$ are image dense open immersions.
- 3 $q_\bullet$, $p_\bullet^X$ and $p_\bullet^U$ are proper.
- 4 $V_\bullet$, $U_\bullet$ and $Y_\bullet$ have flat arrows.
- 5 $p_\bullet^X i'_\bullet = i_\bullet p_\bullet^U$.
- 6 $q_\bullet i_\bullet$ and $p_\bullet^U j_\bullet$ are of fiber type.

Then σ is a fiber square, and the composite map

$$j_\bullet^* \zeta(\sigma) q_\bullet^\times : j_\bullet^* (i'_\bullet)^* (p_\bullet^X)^\times q_\bullet^\times \rightarrow j_\bullet^* (p_\bullet^U)^\times i_\bullet^* q_\bullet^\times$$

is an isomorphism of functors from $\mathcal{D}^+(Y_\bullet)$ to $\mathcal{D}^+(V_\bullet)$.

Proof. The square σ is a fiber square, since the canonical map $U'_\bullet \rightarrow U_\bullet \times_{X_\bullet} X'_\bullet$ is an image dense closed open immersion, and is an isomorphism.

To prove the theorem, it suffices to show that the map in question is an isomorphism after applying $(?)_i$ for any $i \in I$. By Proposition 14.14, Lemma 15.2, and [23, (3.7.2), (iii)], the problem is reduced to the flat base change theorem (in fact open immersion base change theorem is enough) for schemes [36, Theorem 2], and we are done. $\square$

16 The existence of compactification and composition data for finite diagrams of schemes

(16.1) Let S be a noetherian separated scheme, and I an ordered finite category which is non-empty. Let $\mathcal{A}$ denotes the category of I^{op} -diagrams of S -schemes separated of finite type over S . Let $\mathcal{P}$ denote the class of proper morphisms in $\text{Mor}(\mathcal{A})$. Let $\mathcal{I}$ denote the class of image dense open immersions in $\text{Mor}(\mathcal{A})$.

Define $\mathcal{D}(X_\bullet) := D_{\text{LQco}(X_\bullet)}(\text{Mod}(X_\bullet))$ for $X_\bullet \in \text{ob}(\mathcal{A})$. Define a pseudofunctor $(?)^\#$ on $\mathcal{A}_{\mathcal{P}}$ to be $(?)^\times$, where $X_\bullet^\# = \mathcal{D}(X_\bullet)$ for $X_\bullet \in \text{ob}(\mathcal{A}_{\mathcal{P}})$. Define a pseudofunctor $(?)^\flat$ on $\mathcal{A}_{\mathcal{I}}$ to be $(?)^*$, where $X_\bullet^\flat = \mathcal{D}(X_\bullet)$ for $X_\bullet \in \text{ob}(\mathcal{A}_{\mathcal{I}})$.

For a pi-square σ , define $\zeta(\sigma)$ to be the natural map defined in (15.1).

16.2 Lemma. *Let the notation be as above. Conditions 1–6 and 8–12 in Definition 11.1 are satisfied. Moreover, any pi-square is a fiber square.*

Proof. This is easy. $\square$

16.3 Proposition. *Let the notation be as in (16.1). Then the condition 7 in Definition 11.1 is satisfied. That is, for any morphism $f_\bullet$ in $\mathcal{A}$, there is a factorization $f_\bullet = p_\bullet j_\bullet$ with $p_\bullet \in \mathcal{P}$ and $j_\bullet \in \mathcal{I}$.*

Proof. Label the object set $\text{ob}(I)$ of I as $\{i(1), \dots, i(n)\}$ so that $I(i(s), i(t)) = \emptyset$ if $s > t$. Set $J(r)$ to be the full subcategory of I with $\text{ob}(J(r)) = \{i(1), \dots, i(r)\}$. By induction on r , we construct morphisms $h_\bullet(r) : X_\bullet|_{J(r)} \rightarrow Z_\bullet(r)$ in $\mathcal{P}(J(r), \text{Sch}/S)$ such that

- 1 $Z_\bullet(r)$ is proper over S ;
- 2 $h_\bullet(r)_i$ is an open immersion, whose scheme theoretic image is $Z_\bullet(r)_i$ for $i \in J(r)$;

3 $Z_{\bullet}(r)|_{J(j)} = Z_{\bullet}(j)$ and $h_{\bullet}(r)|_{J(j)} = h_{\bullet}(j)$ for $j < r$.

We may assume that the construction is done for $j < r$.

First, consider the case where $i(r)$ is a minimal element in $\text{ob}(I)$. By Nagata's compactification theorem [29], there is an S -morphism

$$h : X_{i(r)} \rightarrow Z$$

such that Z is proper over S and h is an open immersion whose scheme theoretic image is Z . Now define $Z_{\bullet}(r)_{i(r)} := Z$ and $Z_{\bullet}(r)_{\text{id}_{i(r)}} = \text{id}_Z$. Defining the other structures after **3**, we get $Z_{\bullet}(r)$, since $I(i(r), i(s)) = \emptyset = I(i(s), i(r))$ for $s < r$ and $I(i(r), i(r)) = \{\text{id}\}$ by assumption. Define $h_{\bullet}(r)_{i(r)} := h$ and by **3**, we get a morphism $h_{\bullet}(r) : X_{\bullet}|_{J(r)} \rightarrow Z_{\bullet}(r)$ by the same reason, and **1,2,3** are satisfied by induction assumption. So this case is OK.

Now assume that $i(r)$ is not minimal so that $\bigcup_{i < r} I(i(j), i(r)) \neq \emptyset$. Consider the S -morphism

$$\varphi : X_{i(r)} \xrightarrow{(X_{\phi})} \prod_{j < r} \prod_{\phi \in I(i(j), i(r))} X_{i(j)} \xrightarrow{(h_j)} \prod_{j < r} \prod_{\phi \in I(i(j), i(r))} Z_j,$$

where $Z_j := (Z_{\bullet}(j))_{i(j)}$, and $h_j := (h_{\bullet}(j))_{i(j)}$. By Nagata's compactification theorem [29], There is a factorization

$$X_{i(r)} \xrightarrow{h_r} Z_r \xrightarrow{p_r} \prod_{j < r} \prod_{\phi \in I(i(j), i(r))} Z_j$$

such that h_r is an open immersion S -morphism such that the scheme theoretic image of h_r agrees with Z_r , and p_r is a proper S -morphism. Set $Z_{\bullet}(r)_{i(j)} := Z_j$, and $Z_{\bullet}(r)_{i(r)} := Z_r$. For $j \leq j' < r$ and $\phi \in I(i(j), i(j'))$, define $(Z_{\bullet}(r))_{\phi} := Z_{\bullet}(r-1)_{\phi}$. For each $j < r$ and $\phi \in I(i(j), i(r))$, define $(Z_{\bullet}(r))_{\phi} : Z_r \rightarrow Z_j$ to be the composite

$$Z_r \xrightarrow{p_r} \prod_{j < r} \prod_{\phi} Z_j \xrightarrow{\text{projection}} Z_j.$$

Finally, define $(Z_{\bullet}(r))_{\text{id}_{i(r)}} := \text{id}_{Z_r}$. Almost by definition, we have $h_j \circ X_{\phi} = (Z_{\bullet}(r))_{\phi} \circ h_r$ for $j < r$ and $\phi \in I(i(j), i(r))$. As h_r is scheme theoretically dense, it is easy to see that $Z_{\bullet}(r)$ is a $J(r)^{\text{op}}$ diagram of S -schemes, and the conditions **1,2,3** are checked easily.

Hence there is a morphism $h_\bullet : X_\bullet \rightarrow Z_\bullet$ of $\mathcal{P}(I, \text{Sch}/S)$ such that $Z_\bullet$ is proper over S , and $h_\bullet$ is an image-dense open immersion.

Now let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{A}$. Consider the morphism

$$\psi_\bullet : X_\bullet \xrightarrow{\Gamma(f_\bullet)} X_\bullet \times Y_\bullet \xrightarrow{h_\bullet \times 1} Z_\bullet \times Y_\bullet.$$

We define $\bar{X}_\bullet$ as follows. For each $i \in I$, define $\bar{X}_i$ to be the scheme theoretic image of ψ_i , and $\bar{X}_\phi$ is the restriction of $Z_\phi \times Y_\phi$. As $\psi_\bullet$ is an immersion, we have a factorization

$$X_\bullet \xrightarrow{i_\bullet} \bar{X}_\bullet \xrightarrow{q_\bullet} Z_\bullet \times Y_\bullet$$

of $\psi_\bullet$ such that $i_\bullet \in \mathcal{I}$ and $q_\bullet$ is a closed immersion.

Let $p_2 : Z_\bullet \times Y_\bullet \rightarrow Y_\bullet$ be the second projection. Set $p_\bullet := p_2 \circ q_\bullet$. Then, as we have $p_2 \circ \psi_\bullet = f_\bullet$, we have $f_\bullet = p_\bullet i_\bullet$, and $p_\bullet \in \mathcal{P}$. This proves the lemma. $\square$

16.4 Theorem. *Let the notation be as in (16.1). Set $\mathcal{F}$ to be the subcategory of $\mathcal{A}$ whose objects are objects of $\mathcal{A}$ with flat arrows, and whose morphisms are morphisms in $\mathcal{A}$ of fiber type. Define $\mathcal{D}^+$ by $\mathcal{D}^+(X_\bullet) := D_{\text{LQco}(X_\bullet)}^+(\text{Mod}(X_\bullet))$. Then $(\mathcal{A}, \mathcal{F}, \mathcal{P}, \mathcal{I}, \mathcal{D}, \mathcal{D}^+, (?)^\#, (?)^\flat, \zeta)$ is a composition data of contravariant pseudofunctors.*

Proof. Conditions **1**–**12** in Definition 11.1 have already been checked. **13** follows from Lemma 8.15. **14** is trivial. Since I is finite, the definition of $\mathcal{D}^+(X_\bullet)$ is consistent with that in (14.12). Hence **15** is Corollary 14.15. **16** is Theorem 15.3. $\square$

We call the composite of $(?)^\#$ and $(?)^\flat$ defined by the composition data in the theorem the *equivariant twisted inverse pseudofunctor*, and denote it by $(?)^!$.

17 Flat base change

Let the notation be as in Theorem 16.4. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and J a subcategory of I . Let $f_\bullet = p_\bullet i_\bullet$ be a compactification.

17.1 Lemma. *The composite map*

$$(?)_J f_\bullet^! \xrightarrow{\text{via } \rho} (?)_J i_\bullet^* p_\bullet^\times \xrightarrow{\theta^{-1}} i_J^* (?)_J p_\bullet^\times \xrightarrow{\xi} i_J^* p_J^\times (?)_J \xrightarrow{\text{via } \rho} f_J^! (?)_J$$

is independent of choice of compactification $f_\bullet = p_\bullet i_\bullet$, where ρ 's are independence isomorphisms.

The proof utilizes Lemma 11.3, and left to the reader. We denote by $\bar{\xi} = \bar{\xi}(i, f_\bullet)$ the composite map in the lemma.

17.2 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and $K \subset J \subset I$ be subcategories. Then the composite map*

$$(\?)_K f_\bullet^! \cong (\?)_K (\?)_J f_\bullet^! \xrightarrow{\bar{\xi}} (\?)_K f_J^! (\?)_J \xrightarrow{\bar{\xi}} f_K^! (\?)_K (\?)_J \cong f_K^! (\?)_K$$

agrees with $\bar{\xi}(K, f_\bullet)$.

Proof. Follows easily from Lemma 14.4. $\square$

17.3 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and J a subcategory of I . Then, $\bar{\xi}(J, f_\bullet)$ is an isomorphism.*

Proof. It suffices to show that $(\?)_i \bar{\xi}(J, f_\bullet)$ is an isomorphism for any $i \in \text{ob}(J)$. By Lemma 17.2, we have

$$\bar{\xi}(i, f_J) \circ ((\?)_i \bar{\xi}(J, f_\bullet)) = \bar{\xi}(i, f_\bullet).$$

By Proposition 14.14, we have $\bar{\xi}(i, f_J)$ and $\bar{\xi}(i, f_\bullet)$ are isomorphisms. Hence the natural map $(\?)_i \bar{\xi}(J, f_\bullet)$ is also an isomorphism. $\square$

17.4 Lemma. *Let $f : X \rightarrow Y$ be a flat morphism of locally noetherian schemes, and U a dense open subset of Y . Then $f^{-1}(U)$ is a dense open subset of X .*

Proof. The question is local both on Y and X , and hence we may assume that both $Y = \text{Spec } A$ and $X = \text{Spec } B$ are affine. Let I be the radical ideal of A defining the closed subset $Y \setminus U$. By assumption, I is not contained in any minimal prime of A . Assume that $f^{-1}(U)$ is not dense in X . Then, there is a minimal prime P of B which contains IB . As we have $I \subset IB \cap A \subset P \cap A$ and $P \cap A$ is minimal by the going-down theorem (see [25, Theorem 9.5]), this is a contradiction. $\square$

(17.5) Let $b : S' \rightarrow S$ be a morphism of noetherian separated schemes. Let $\mathcal{A}'$ be the full subcategory of $\mathcal{P}(I, \text{Sch}/S')$ consisting of objects separated of finite type over S' . Let $\mathcal{F}'$ be the category of objects of $\mathcal{A}'$ with flat arrows and morphisms in $\mathcal{A}'$ of fiber type. Any S' -scheme is viewed as an S -scheme via b .

Let

$$\begin{array}{ccc} X'_\bullet & \xrightarrow{f'_\bullet} & Y'_\bullet \\ \downarrow g_\bullet^X \sigma & & \downarrow g_\bullet \\ X_\bullet & \xrightarrow{f_\bullet} & Y_\bullet \end{array}$$

be a diagram in $\mathcal{P}(I, \text{Sch}/S)$ such that

- 1 $X'_\bullet$ and $Y'_\bullet$ are in $\mathcal{F}'$, and $f'_\bullet$ is a morphism in $\mathcal{F}'$.
- 2 $f_\bullet$ is a morphism in $\mathcal{F}$;
- 3 σ is a fiber square;
- 4 $g_\bullet$ is flat.

By assumption, there is a diagram

$$\begin{array}{ccccc} X'_\bullet & \xrightarrow{i'_\bullet} & Z'_\bullet & \xrightarrow{p'_\bullet} & Y'_\bullet \\ \downarrow g_\bullet^X \sigma_1 & & \downarrow g_\bullet^Z \sigma_2 & & \downarrow g_\bullet \\ X_\bullet & \xrightarrow{i_\bullet} & Z_\bullet & \xrightarrow{p_\bullet} & Y_\bullet \end{array} \quad (17.6)$$

such that $f_\bullet = p_\bullet i_\bullet$ is a compactification, σ_1 and σ_2 are fiber squares, and the whole rectangle $\sigma_1 \sigma_2$ equals σ . By Lemma 17.4, we have that $f'_\bullet = p'_\bullet i'_\bullet$ is a compactification.

17.7 Lemma. *The composite map*

$$(g_\bullet^X)^* f_\bullet^! \xrightarrow{\rho} (g_\bullet^X)^* i_\bullet^* p_\bullet^\times \xrightarrow{d} (i'_\bullet)^* (g_\bullet^Z)^* p_\bullet^\times \xrightarrow{\zeta} (i'_\bullet)^* (p'_\bullet)^\times g_\bullet^* \xrightarrow{\rho} (f'_\bullet)^! g_\bullet^*$$

is independent of choice of the diagram (17.6), and depends only on σ , where ρ 's are independence isomorphisms.

Proof. Obvious by Lemma 11.3. □

We denote the composite map in the lemma by $\bar{\zeta} = \bar{\zeta}(\sigma)$.

17.8 Theorem. *Let the notation be as above. Then we have:*

- 1 *Let J be an admissible subcategory of I . Then the diagram*

$$\begin{array}{ccccccc} (?)_J (g_\bullet^X)^* f_\bullet^! & \xrightarrow{\theta^{-1}} & (g_J^X)^* (?)_J f_\bullet^! & \xrightarrow{\bar{\xi}} & (g_J^X)^* f_J^! (?)_J & & \\ \downarrow (?)_J \bar{\zeta}(\sigma) & & & & \downarrow \bar{\zeta}(\sigma_J)(?)_J & & (17.9) \\ (?)_J (f'_\bullet)^! (g_\bullet)^* & \xrightarrow{\bar{\xi}} & (f'_J)^! (?)_J (g_\bullet)^* & \xrightarrow{\theta^{-1}} & (f'_J)^! (g_J)^* (?)_J & & \end{array}$$

is commutative.

2 $\bar{\zeta}(\sigma)$ is an isomorphism.

Proof. **1** is an immediate consequence of Lemma 15.2 and [23, (3.7.2)].

2 Let i be an object of I . By Lemma 17.3, the horizontal arrows in the diagram (17.9) are isomorphisms. By Verdier's flat base change theorem [36, Theorem 2], we have that $\bar{\zeta}(\sigma_i)$ is an isomorphism. Hence, we have that $(?)_i \bar{\zeta}(\sigma)$ is an isomorphism for any $i \in I$ by **1** applied to $J = i$, and the assertion follows. $\square$

18 Preservation of Quasi-coherent cohomology

(18.1) Let the notation be as in (16.1). Let $\mathcal{F}$ be as in Theorem 16.4.

18.2 Lemma. *Let $f : X \rightarrow Y$ be a morphism of finite type between separated noetherian schemes. If $\mathbb{F} \in \mathcal{D}_{\text{Coh}(Y)}^+(\text{Mod}(Y))$, then $f^! \mathbb{F} \in \mathcal{D}_{\text{Coh}(X)}^+(\text{Mod}(X))$.*

Proof. We may assume that both Y and X are affine. So we may assume that f is either smooth or a closed immersion. The case where f is smooth is obvious by [36, Theorem 3]. The case where f is a closed immersion is also obvious by Proposition III.6.1 and Theorem III.6.7 in [15]. $\square$

18.3 Proposition. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and $\phi : i \rightarrow j$ a morphism in I . Then the composite map*

$$X_\phi^*(?)_i f_\bullet^! \xrightarrow{\alpha_\phi} (?)_j f_\bullet^! \xrightarrow{\bar{\xi}} f_j^! (?)_j$$

agrees with the composite map

$$X_\phi^*(?)_i f_\bullet^! \xrightarrow{\bar{\xi}} X_\phi^* f_i^! (?)_i \xrightarrow{\bar{\zeta}} f_j^! Y_\phi^*(?)_i \xrightarrow{\alpha_\phi} f_j^! (?)_j.$$

Proof. By Lemma 17.2, we may assume that I is the ordered category given by $\text{ob}(I) = \{i, j\}$ and $I(i, j) = \{\phi\}$.

Then it is easy to see that there is a compactification

$$X_\bullet \xrightarrow{i_\bullet} Z_\bullet \xrightarrow{p_\bullet} Y_\bullet$$

of $f_\bullet$ such that $p_\bullet$ is of fiber type. Note that $i_\bullet$ is of fiber type, and $Z_\bullet$ has flat arrows.

By the definition of $\bar{\xi}$ and $\bar{\zeta}$, it suffices to prove that the composite map

$$X_\phi^*(?)_i i_\bullet^* p_\bullet^\times \xrightarrow{\theta^{-1}} X_\phi^* i_i^*(?)_i p_\bullet^\times \xrightarrow{d} i_j^* Z_\phi^*(?)_i p_\bullet^\times \xrightarrow{\xi} i_j^* Z_\phi^* p_i^\times (?)_i \xrightarrow{\zeta} i_j^* p_j^\times Y_\phi^*(?)_i \xrightarrow{\alpha_\phi} i_j^* p_j^\times (?)_j$$

agrees with

$$X_\phi^*(?)_i i_\bullet^* p_\bullet^\times \xrightarrow{\alpha_\phi} (?)_j i_\bullet^* p_\bullet^\times \xrightarrow{\theta^{-1}} i_j^*(?)_j p_\bullet^\times \xrightarrow{\xi} i_j^* p_j^\times (?)_j.$$

By the “derived version” of (7.25), the composite map

$$X_\phi^*(?)_i i_\bullet^* p_\bullet^\times \xrightarrow{\theta^{-1}} X_\phi^* i_i^*(?)_i p_\bullet^\times \xrightarrow{d} i_j^* Y_\phi^*(?)_i p_\bullet^\times \xrightarrow{\alpha_\phi} i_j^*(?)_j p_\bullet^\times$$

agrees with

$$X_\phi^*(?)_i i_\bullet^* p_\bullet^\times \xrightarrow{\alpha_\phi} (?)_j i_\bullet^* p_\bullet^\times \xrightarrow{\theta^{-1}} i_j^*(?)_j p_\bullet^\times.$$

Hence it suffices to prove the map

$$Z_\phi^*(?)_i p_\bullet^\times \xrightarrow{\xi} Z_\phi^* p_i^\times (?)_i \xrightarrow{\zeta} p_j^\times Y_\phi^*(?)_i \xrightarrow{\alpha_\phi} p_j^\times (?)_j$$

agrees with

$$Z_\phi^*(?)_i p_\bullet^\times \xrightarrow{\alpha_\phi} (?)_j p_\bullet^\times \xrightarrow{\xi} p_j^\times (?)_j.$$

Now the proof consists in a straightforward diagram drawing utilizing the derived version of (7.14). $\square$

18.4 Corollary. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$. Then we have $f_\bullet^!(\mathcal{D}_{\text{Qco}}^+(Y_\bullet)) \subset \mathcal{D}_{\text{Qco}}^+(X_\bullet)$ and $f_\bullet^!(\mathcal{D}_{\text{Coh}}^+(Y_\bullet)) \subset \mathcal{D}_{\text{Coh}}^+(X_\bullet)$.*

Proof. Let $\phi : i \rightarrow j$ be a morphism in I . By Theorem 17.8, **2**, Lemma 17.3 and the proposition, we have $\alpha_\phi : X_\phi^*(?)_i f_\bullet^! \rightarrow (?)_j f_\bullet^!$ is an isomorphism if $\alpha_\phi : Y_\phi^*(?)_i \rightarrow (?)_j$ is an isomorphism. So $f_\bullet^!$ preserves equivariance of cohomology groups, and the first assertion follows.

On the other hand, by Lemma 18.2 and Proposition 14.14, $f^!$ preserves local coherence of cohomology groups. Hence it also preserves the coherence of cohomology groups, by the first paragraph. $\square$

19 Compatibility with derived direct image

(19.1) Let the notation be as in (17.5). Consider the diagram (17.6). Lipman's theta $\theta(\sigma_2) : g_{\bullet}^* R(p_{\bullet})_* \rightarrow R(p'_{\bullet})_*(g_{\bullet}^Z)^*$ induces the conjugate map

$$\xi(\sigma_2) : R(g_{\bullet}^Z)_*(p'_{\bullet})^{\times} \rightarrow p_{\bullet}^{\times} R(g_{\bullet})_*.$$

As σ_2 is a fiber square, $\theta(\sigma_2)$ is an isomorphism. Hence $\xi(\sigma_2)$ is also an isomorphism. Note that

$$\theta : i_{\bullet}^* R(g_{\bullet}^Z)_* \rightarrow R(g_{\bullet}^X)_*(i'_{\bullet})^*$$

is an isomorphism, since σ_1 is a fiber square. We define $\bar{\xi} : R(g_{\bullet}^X)_*(f'_{\bullet})^! \rightarrow f_{\bullet}^! R(g_{\bullet})_*$ to be the composite

$$R(g_{\bullet}^X)_*(f'_{\bullet})^! \xrightarrow{\rho} R(g_{\bullet}^X)_*(i'_{\bullet})^*(p'_{\bullet})^{\times} \xrightarrow{\theta^{-1}} i_{\bullet}^* R(g_{\bullet}^Z)_*(p'_{\bullet})^{\times} \xrightarrow{\xi} i_{\bullet}^* p_{\bullet}^{\times} R(g_{\bullet})_* \xrightarrow{\rho} f_{\bullet}^! R(g_{\bullet})_*.$$

As all the maps in the composition are isomorphisms, we have

19.2 Lemma. $\bar{\xi}$ is an isomorphism.

19.3 Lemma. For any subcategory J , the composite

$$(\?)_J R(g_{\bullet}^Z)_*(p'_{\bullet})^{\times} \xrightarrow{\xi} (\?)_J p_{\bullet}^{\times} R(g_{\bullet})_* \xrightarrow{\xi} (p_{\bullet}|_J)^{\times} (\?)_J R(g_{\bullet})_* \xrightarrow{c} (p_{\bullet}|_J)^{\times} R(g_{\bullet}|_J)_*(\?)_J$$

agrees with the composite

$$\begin{aligned} (\?)_J R(g_{\bullet}^Z)_*(p'_{\bullet})^{\times} &\xrightarrow{c} R(g_{\bullet}^Z|_J)_*(\?)_J (p'_{\bullet})^{\times} \xrightarrow{\xi} \\ &R(g_{\bullet}^Z|_J)_*(p'_{\bullet}|_J)^{\times} (\?)_J \xrightarrow{\xi} (p_{\bullet}|_J)^{\times} R(g_{\bullet}|_J)_*(\?)_J. \end{aligned}$$

Proof. Follows immediately from Lemma 14.7. $\square$

20 Compatibility with derived right inductions

(20.1) Let the notation be as in Theorem 16.4. Let $X_{\bullet} \in \mathcal{F}$, and J a subcategory of I . As I is assumed to be a finite category, $I_i^{(J \rightarrow I)}$ is a finite category for any $i \in \text{ob}(I)$. Hence for any $\mathcal{M} \in \text{LQco}(X_{\bullet}|_J)$, we

have that $R_J \mathcal{M} \in \text{LQco}(X_\bullet)$ by (7.10). It is easy to see that $RR_J : D_{\text{LQco}(X_\bullet|_J)}^+(\text{Mod}(X_\bullet|_J)) \rightarrow D_{\text{LQco}(X_\bullet)}^+(\text{Mod}(X_\bullet))$ is the right adjoint of $(?)_J$.

Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{A}$. As

$$c : (?)_J R(f_\bullet)_* \rightarrow R(f_\bullet|_J)_* (?)_J$$

is an isomorphism, its conjugate map

$$c' : RR_J(f_\bullet|_J)^\times \rightarrow f_\bullet^\times RR_J$$

is also an isomorphism.

Let $g_\bullet : U_\bullet \rightarrow X_\bullet$ be an image dense open immersion of fiber type in $\mathcal{A}$. Let σ be the canonical map

$$g_\bullet^* RR_J \xrightarrow{u} g_\bullet^* RR_J R(g_\bullet|_J)_* (g_\bullet|_J)^* \xrightarrow{\xi^{-1}} g_\bullet^* R(g_\bullet)_* RR_J(g_\bullet|_J)^* \xrightarrow{\varepsilon} RR_J(g_\bullet|_J)^*,$$

where $\xi : R(g_\bullet)_* RR_J \rightarrow RR_J R(g_\bullet|_J)_*$ is the conjugate of the isomorphism $\theta : (g_\bullet|_J)^* (?)_J \rightarrow (?)_J g_\bullet^*$.

20.2 Lemma. *Let the notation be as above. Then $\sigma : g_\bullet^* RR_J \rightarrow RR_J(g_\bullet|_J)^*$ is an isomorphism of functors from $D(\text{Mod}(X_\bullet|_J))$ to $D(\text{Mod}(U_\bullet))$.*

Proof. As $g_\bullet^*$, $(g_\bullet|_J)^*$, R_J , $(g_\bullet)_*$, and $(g_\bullet|_J)_*$ have exact left adjoints, it suffices to show that

$$(?)_i \sigma : (?)_i g_\bullet^* R_J \mathbb{I} \rightarrow (?)_i R_J(g_\bullet|_J)^* \mathbb{I}$$

is an isomorphism for any K -injective complex in $C(\text{Mod}(X_\bullet|_J))$ and $i \in \text{ob}(I)$. This map agrees with

$$\begin{aligned} (?)_i g_\bullet^* R_J \mathbb{I} &\xrightarrow{\theta^{-1}} g_i^* (?)_i R_J \mathbb{I} \cong g_i^* \varprojlim (X_\phi)_* \mathbb{I}_j \cong \varprojlim g_i^* (X_\phi)_* \mathbb{I}_j \\ &\xrightarrow{\theta} \varprojlim (U_\phi)_* g_j^* \mathbb{I}_j \xrightarrow{\theta} \varprojlim (U_\phi)_* (?)_j (g_\bullet|_J)^* \mathbb{I} \cong R_J(g_\bullet|_J)^* \mathbb{I}, \end{aligned}$$

which is obviously an isomorphism. $\square$

(20.3) Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and $f_\bullet = p_\bullet i_\bullet$ a compactification. We define $\bar{c} : f_\bullet^! RR_J \rightarrow RR_J(f_\bullet|_J)^!$ to be the composite

$$f_\bullet^! RR_J \xrightarrow{\rho} i_\bullet^* p_\bullet^\times RR_J \xrightarrow{c'} i_\bullet^* RR_J(p_\bullet|_J)^\times \xrightarrow{\sigma} RR_J(i_\bullet|_J)^* (p_\bullet|_J)^\times \xrightarrow{\rho} RR_J(f_\bullet|_J)^!.$$

By Lemma 20.2, we have

20.4 Lemma. $\bar{c} : f_\bullet^! RR_J \rightarrow RR_J(f_\bullet|_J)^!$ is an isomorphism of functors from $D_{\text{LQco}(Y_\bullet|_J)}^+(\text{Mod}(X_\bullet))$ to $D_{\text{LQco}(X_\bullet)}^+(\text{Mod}(X_\bullet))$.

21 Equivariant Grothendieck's duality

21.1 Theorem (Equivariant Grothendieck's duality). *Let I be a small category, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/\mathbb{Z})$. If $Y_\bullet$ is separated noetherian with flat arrows and $f_\bullet$ is proper of fiber type, then the composite*

$$\begin{aligned} \Theta(f_\bullet) : R(f_\bullet)_* R \underline{\text{Hom}}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, f_\bullet^\times \mathbb{G}) &\xrightarrow{H} R \underline{\text{Hom}}_{\text{Mod}(Y_\bullet)}^\bullet(R(f_\bullet)_* \mathbb{F}, R(f_\bullet)_* f_\bullet^\times \mathbb{G}) \\ &\xrightarrow{\varepsilon} R \underline{\text{Hom}}_{\text{Mod}(Y_\bullet)}^\bullet(R(f_\bullet)_* \mathbb{F}, \mathbb{G}) \end{aligned}$$

is an isomorphism for $\mathbb{F} \in \mathcal{D}_{\text{Coh}}^-(X_\bullet)$ and $\mathbb{G} \in \mathcal{D}^+(Y_\bullet)$.

Proof. It suffices to show that $(?)_i \Theta(f_\bullet)$ is an isomorphism for $i \in \text{ob}(I)$. By Lemma 2.10 and Lemma 14.7, **2**, it is easy to see that $(?)_i \Theta(f_\bullet)$ agrees with the composite

$$\begin{aligned} (?)_i R(f_\bullet)_* R \underline{\text{Hom}}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, f_\bullet^\times \mathbb{G}) &\xrightarrow{c} R(f_i)_* (?)_i R \underline{\text{Hom}}_{\text{Mod}(X_\bullet)}^\bullet(\mathbb{F}, f_\bullet^\times \mathbb{G}) \\ &\xrightarrow{H_i} R(f_i)_* R \underline{\text{Hom}}_{\text{Mod}(X_i)}^\bullet(\mathbb{F}_i, (?)_i f_\bullet^\times \mathbb{G}) \xrightarrow{\xi} R(f_i)_* R \underline{\text{Hom}}_{\text{Mod}(X_i)}^\bullet(\mathbb{F}_i, f_i^\times \mathbb{G}_i) \\ &\xrightarrow{\Theta(f_i)} R \underline{\text{Hom}}_{\text{Mod}(Y_i)}^\bullet(R(f_i)_* \mathbb{F}_i, \mathbb{G}_i). \end{aligned}$$

By Lemma 13.9, H_i is an isomorphism. By Lemma 14.5, ξ is an isomorphism. By Grothendieck's duality for usual schemes [30, section 6], $\Theta(f_i)$ is an isomorphism. Hence, $(?)_i \Theta(f_\bullet)$ is an isomorphism. $\square$

22 Morphisms of finite flat dimension

(22.1) Let $((?)^*, (?)_*)$ be a monoidal adjoint pair over a category $\mathcal{S}$. For a morphism $f : X \rightarrow Y$ in $\mathcal{S}$, we define the projection morphism $\Pi = \Pi(f)$ to be the composite

$$f_* a \otimes b \xrightarrow{u} f_* f^*(f_* a \otimes b) \xrightarrow{\Delta} f_*(f^* f_* a \otimes f^* b) \xrightarrow{\varepsilon} f_*(a \otimes f^* b),$$

where $a \in M_X$ and $b \in M_Y$ (see section 2 for notation).

22.2 Lemma. *Let the notation be as above, and $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in $\mathcal{S}$. For $x \in M_X$ and $z \in M_Z$, the composite*

$$(gf)_* x \otimes z \xrightarrow{c} g_*(f_* x) \otimes z \xrightarrow{\Pi(g)} g_*(f_* x \otimes g^* z) \xrightarrow{\Pi(f)} g_* f_*(x \otimes f^* g^* z) \xrightarrow{\cong} (gf)_*(x \otimes (gf)^* z)$$

agrees with $\Pi(gf)$.

Proof. Left to the reader. $\square$

(22.3) Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/S)$.

22.4 Lemma (Projection Formula). *Assume that $X_\bullet$ and $Y_\bullet$ are quasi-compact and separated. Then the natural map*

$$\Pi = \Pi(f_\bullet) : (Rf_\bullet)_* \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G} \rightarrow (Rf_\bullet)_* (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G})$$

is an isomorphism for $\mathbb{F} \in D_{\text{LQco}}(X_\bullet)(\text{Mod}(X_\bullet))$ and $\mathbb{G} \in D_{\text{LQco}}(Y_\bullet)(\text{Mod}(Y_\bullet))$.

Proof. For each $i \in \text{ob}(I)$, the composite

$$\begin{aligned} R(f_i)_* \mathbb{F}_i \otimes_{\mathcal{O}_{Y_i}}^L \mathbb{G}_i &\xrightarrow{\Pi(f_i)} R(f_i)_* (\mathbb{F}_i \otimes_{\mathcal{O}_{X_i}}^L Lf_i^* \mathbb{G}_i) \xrightarrow{\theta(f_\bullet, i)} R(f_i)_* (\mathbb{F}_i \otimes_{\mathcal{O}_{X_i}}^L (?)_i Lf_i^* \mathbb{G}) \\ &\xrightarrow{m} R(f_i)_* (?)_i (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}) \xrightarrow{c} (?)_i R(f_\bullet)_* (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}) \end{aligned}$$

is an isomorphism by [30, Proposition 5.3] and Lemma 9.7, **1**. On the other hand, it is straightforward to check that this composite isomorphism agrees with the composite

$$\begin{aligned} R(f_i)_* \mathbb{F}_i \otimes_{\mathcal{O}_{Y_i}}^L \mathbb{G}_i &\xrightarrow{c} (?)_i R(f_\bullet)_* \mathbb{F} \otimes_{\mathcal{O}_{Y_i}}^L \mathbb{G}_i \xrightarrow{m} (?)_i (R(f_\bullet)_* \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \\ &\xrightarrow{(?)_i \Pi(f_\bullet)} (?)_i R(f_\bullet)_* (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}). \end{aligned}$$

It follows that $(?)_i \Pi(f_\bullet)$ is an isomorphism for any $i \in \text{ob}(I)$. Hence, $\Pi(f_\bullet)$ is an isomorphism. $\square$

(22.5) Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{P}(I, \text{Sch}/S)$, and assume that both $X_\bullet$ and $Y_\bullet$ are quasi-compact separated. Define $\Phi = \Phi(f_\bullet)$ to be the composite

$$\begin{aligned} f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G} &\xrightarrow{u} f_\bullet^\times R(f_\bullet)_* (f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}) \\ &\xrightarrow{\Pi(f_\bullet)^{-1}} f_\bullet^\times (R(f_\bullet)_* f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \xrightarrow{\varepsilon} f_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}), \end{aligned}$$

where $\mathbb{F}, \mathbb{G} \in D_{\text{LQco}}(Y_\bullet)(\text{Mod}(Y_\bullet))$.

Utilizing the commutativity as in the proof of Lemma 22.4 and Lemma 14.7, it is not so difficult to show the following.

22.6 Lemma. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be as in (22.5). For a subcategory J of I , the composite

$$\begin{aligned} (?)_J f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet|J}}^{\bullet, L} (?)_J Lf_\bullet^* \mathbb{G} &\xrightarrow{\xi \otimes \theta^{-1}} (f_\bullet|_J)^\times \mathbb{F}_J \otimes_{\mathcal{O}_{X_\bullet|J}}^{\bullet, L} L(f_\bullet|_J)^* \mathbb{G}_J \\ &\xrightarrow{\Phi(f_\bullet|_J)} (f_\bullet|_J)^\times (\mathbb{F}_J \otimes_{\mathcal{O}_{Y_\bullet|J}} \mathbb{G}_J) \xrightarrow{m} (f_\bullet|_J)^\times (?)_J (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}} \mathbb{G}) \end{aligned}$$

agrees with

$$\begin{aligned} (?)_J f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet|J}}^{\bullet, L} (?)_J Lf_\bullet^* \mathbb{G} &\xrightarrow{m} (?)_J (f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}) \\ &\xrightarrow{\Phi(f_\bullet)} (?)_J f_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \xrightarrow{\xi} (f_\bullet|_J)^\times (?)_J (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}). \end{aligned}$$

22.7 Lemma. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be as in (22.5). The composite

$$\begin{aligned} f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* (\mathbb{G} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{H}) &\xrightarrow{\Delta} f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{H} \\ &\xrightarrow{\Phi} f_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{H} \xrightarrow{\Phi} f_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{H}) \end{aligned}$$

agrees with Φ .

22.8 Lemma. Let S , I and σ be as in (15.1). For $\mathbb{F} \in D_{\text{LQco}(Y_\bullet)}(\text{Mod}(Y_\bullet))$, the composite

$$\begin{aligned} (g'_\bullet)^* f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X'_\bullet}}^{\bullet, L} (g'_\bullet)^* Lf_\bullet^* \mathbb{G} &\xrightarrow{\Delta^{-1}} (g'_\bullet)^* (f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} Lf_\bullet^* \mathbb{G}) \\ &\xrightarrow{\Phi} (g'_\bullet)^* f_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \xrightarrow{\zeta(\sigma)} (f'_\bullet)^\times g_\bullet^* (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \end{aligned}$$

agrees with

$$\begin{aligned} (g'_\bullet)^* f_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X'_\bullet}}^{\bullet, L} (g'_\bullet)^* Lf_\bullet^* \mathbb{G} &\xrightarrow{\zeta(\sigma) \otimes d} (f'_\bullet)^\times g_\bullet^* \mathbb{F} \otimes_{\mathcal{O}_{X'_\bullet}}^{\bullet, L} L(f'_\bullet)^* g_\bullet^* \mathbb{G} \\ &\xrightarrow{\Phi} (f'_\bullet)^\times (g_\bullet^* \mathbb{F} \otimes_{\mathcal{O}_{Y'_\bullet}}^{\bullet, L} g_\bullet^* \mathbb{G}) \xrightarrow{\Delta^{-1}} (f'_\bullet)^\times g_\bullet^* (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}). \end{aligned}$$

22.9 Lemma. Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y_\bullet \rightarrow Z_\bullet$ be morphisms in $\mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$, $Y_\bullet$ and $Z_\bullet$ are quasi-compact separated. Then the composite

$$\begin{aligned} (g_\bullet f_\bullet)^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} (g_\bullet f_\bullet)^* \mathbb{G} &\cong f_\bullet^\times g_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} f_\bullet^* g_\bullet^* \mathbb{G} \xrightarrow{\Phi(f_\bullet)} f_\bullet^\times (g_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} g_\bullet^* \mathbb{G}) \\ &\xrightarrow{\Phi(g_\bullet)} f_\bullet^\times g_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Z_\bullet}}^{\bullet, L} \mathbb{G}) \cong (g_\bullet f_\bullet)^\times (\mathbb{F} \otimes_{\mathcal{O}_{Z_\bullet}}^{\bullet, L} \mathbb{G}) \end{aligned}$$

agrees with $\Phi(g_\bullet f_\bullet)$.

The proof of the lemmas above are left to the reader.

(22.10) Let the notation be as in Theorem 16.4. Let $i_\bullet : U_\bullet \rightarrow X_\bullet$ be a morphism in $\mathcal{I}$, and $p_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}$.

We define $\bar{\Phi} = \bar{\Phi}(p_\bullet, i_\bullet)$ to be the composite

$$\begin{aligned} \bar{\Phi} : i_\bullet^* p_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{U_\bullet}}^{\bullet, L} L(p_\bullet i_\bullet)^* \mathbb{G} &\xrightarrow{d^{-1}} i_\bullet^* p_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{U_\bullet}}^{\bullet, L} i_\bullet^* L p_\bullet^* \mathbb{G} \\ &\xrightarrow{\Delta^{-1}} i_\bullet^* (p_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} L p_\bullet^* \mathbb{G}) \xrightarrow{i_\bullet^* \Phi} i_\bullet^* p_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}). \end{aligned}$$

22.11 Lemma. *Let $f_\bullet$ be a morphism in $\mathcal{F}$, and $f_\bullet = p_\bullet i_\bullet = q_\bullet j_\bullet$ an independence square. Then the composite*

$$i_\bullet^* p_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{U_\bullet}}^{\bullet, L} L(p_\bullet i_\bullet)^* \mathbb{G} \xrightarrow{\bar{\Phi}} i_\bullet^* p_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}) \xrightarrow{\rho(p_\bullet i_\bullet = q_\bullet j_\bullet)} j_\bullet^* q_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G})$$

agrees with

$$i_\bullet^* p_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{U_\bullet}}^{\bullet, L} L(p_\bullet i_\bullet)^* \mathbb{G} \xrightarrow{\rho \otimes 1} j_\bullet^* q_\bullet^\times \mathbb{F} \otimes_{\mathcal{O}_{U_\bullet}}^{\bullet, L} L(q_\bullet j_\bullet)^* \mathbb{G} \xrightarrow{\bar{\Phi}} j_\bullet^* q_\bullet^\times (\mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} \mathbb{G}).$$

Proof. As ρ is constructed from ζ and d by definition, the assertion follows easily from Lemma 22.8 and Lemma 22.9. $\square$

(22.12) Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$. We define $\bar{\Phi}(f_\bullet)$ to be $\bar{\Phi}(p_\bullet, i_\bullet)$, where $f_\bullet = p_\bullet i_\bullet$ is the (fixed) compactification of $f_\bullet$.

By Lemma 22.11, $\bar{\Phi}(f_\bullet)$ is an isomorphism if and only if there exists some compactification $f_\bullet = q_\bullet j_\bullet$ such that $\bar{\Phi}(q_\bullet, j_\bullet)$ is an isomorphism.

22.13 Lemma. *Let the notation be as in Theorem 16.4, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ and $g_\bullet : Y_\bullet \rightarrow Z_\bullet$ morphisms in $\mathcal{F}$. Then the composite*

$$\begin{aligned} (g_\bullet f_\bullet)^! \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} L(g_\bullet f_\bullet)^* \mathbb{G} &\cong f_\bullet^! g_\bullet^! \mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} L f_\bullet^* L g_\bullet^* \mathbb{G} \xrightarrow{\bar{\Phi}(f_\bullet)} f_\bullet^! (g_\bullet^! \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} L g_\bullet^* \mathbb{G}) \\ &\xrightarrow{\bar{\Phi}(g_\bullet)} f_\bullet^! g_\bullet^! (\mathbb{F} \otimes_{\mathcal{O}_{Z_\bullet}}^{\bullet, L} \mathbb{G}) \cong (g_\bullet f_\bullet)^! (\mathbb{F} \otimes_{\mathcal{O}_{Z_\bullet}}^{\bullet, L} \mathbb{G}) \end{aligned}$$

agrees with $\bar{\Phi}(g_\bullet f_\bullet)$.

22.14 Theorem. *Let the notation be as in Theorem 16.4, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{F}$. If $f_\bullet$ is of finite flat dimension, then*

$$\bar{\Phi}(f_\bullet) : f_\bullet^! \mathbb{F} \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} L f_\bullet^* \mathbb{G} \rightarrow f_\bullet^! (\mathbb{F} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{G})$$

is an isomorphism for $\mathbb{F}, \mathbb{G} \in D_{\text{LQco}(Y_\bullet)}^+(\text{Mod}(Y_\bullet))$.

Proof. Let $f_\bullet = p_\bullet i_\bullet$ be a compactification of $f_\bullet$. It suffices to show that $\bar{\Phi}(p_\bullet, i_\bullet)$ is an isomorphism. In view of Proposition 14.14 and Lemma 22.6, it suffices to show that

$$\bar{\Phi}(f_j) : i_j^* p_j^\times \mathbb{F}_j \otimes_{\mathcal{O}_{X_j}}^{\bullet, L} Lf_j^* \mathbb{G}_i \rightarrow i_j^* p_j^\times (\mathbb{F}_j \otimes_{\mathcal{O}_{Y_j}}^{\bullet, L} \mathbb{G}_j)$$

is an isomorphism for any $j \in \text{ob}(I)$. So we may assume that $I = j$.

By the flat base change theorem and Lemma 22.8, the question is local on Y_j . Clearly, the question is local on U_j . Hence we may assume that Y_j and U_j are affine. Set $f = f_j$, $Y = Y_j$, and $U = U_j$. Note that f is a closed immersion defined by an ideal of finite projective dimension, followed by an affine n -space.

By Lemma 22.13, it suffices to prove that $\bar{\Phi}(f)$ is an isomorphism if f is a closed immersion defined by an ideal of finite projective dimension or an affine n -space.

By Lemma 22.7, we may assume that $\mathbb{F} = \mathcal{O}_Y$. Both cases are proved easily, using [30, Theorem 5.4]. $\square$

23 Finite morphisms of fiber type

(23.1) Let I be a small category, S a scheme, and $f_\bullet : X_\bullet \rightarrow Y_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Let Z denote the ringed site $(\text{Zar}(Y_\bullet), (f_\bullet)_*(\mathcal{O}_{X_\bullet}))$. Assume that $Y_\bullet$ is locally noetherian. There are obvious admissible ringed continuous functors $i : \text{Zar}(Y_\bullet) \rightarrow Z$ and $g : Z \rightarrow \text{Zar}(X_\bullet)$ such that $gi = f_\bullet^{-1}$. It is easy to see that $g_\# : \text{Mod}(Z) \rightarrow \text{Mod}(X_\bullet)$ is an exact functor.

23.2 Lemma. *If $f_\bullet$ is affine, then the counit*

$$\varepsilon : g_\# Rg^\# \mathbb{F} \rightarrow \mathbb{F}$$

is an isomorphism for $\mathbb{F} \in D_{\text{LQco}(X_\bullet)}^+(\text{Mod}(X_\bullet))$.

Proof. The construction of ε is compatible with restrictions. So we may assume that $f_\bullet = f : X \rightarrow Y$ is an affine morphism of single schemes. Further, the question is local on Y , and hence we may assume that $Y = \text{Spec } A$ is affine. As f is affine, $X = \text{Spec } B$ is affine. By the way-out lemma, we may assume that $\mathbb{F}$ is a single quasi-coherent sheaf, say $\tilde{M}$, where M is a B -module. As $R^i f_* \tilde{M} = 0$ for $i > 0$, it suffices to show that $\varepsilon : g_\# g^\# \tilde{M} \rightarrow \tilde{M}$ is an isomorphism. As $g_\# g^\#$ is exact on $\tilde{M}$ and preserves direct sums, we may assume that $M = B$, which case is trivial. $\square$

(23.3) Let $I, S, f_\bullet : X_\bullet \rightarrow Y_\bullet, Z, g$, and i be as in (23.1). Assume that $f_\bullet$ is finite of fiber type.

We say that an $\mathcal{O}_Z$ -module $\mathcal{M}$ is locally quasi-coherent (resp. quasi-coherent, coherent) if $i^\# \mathcal{M}$ is. The corresponding full subcategory of $\text{Mod}(Z)$ is denoted by $\text{LQco}(Z)$ (resp. $\text{Qco}(Z), \text{Coh}(Z)$).

23.4 Lemma. *Let the notation be as above. Then an $\mathcal{O}_Z$ -module $\mathcal{M}$ is locally quasi-coherent if and only if for any $i \in \text{ob}(I)$ and any affine open subscheme U of Y_i , there exists an exact sequence of $((\mathcal{O}_Z)_i)|_U$ -modules*

$$(((\mathcal{O}_Z)_i)|_U)^{(T)} \rightarrow (((\mathcal{O}_Z)_i)|_U)^{(\Sigma)} \rightarrow \mathcal{M}_i|_U \rightarrow 0.$$

Proof. As we assume that $f_\bullet$ is finite of fiber type, $\mathcal{O}_Z$ is coherent. Hence the existence of such exact sequences implies that $\mathcal{M}$ is locally quasi-coherent.

We prove the converse. Let $i \in \text{ob}(I)$ and U an affine open subset of Y_i . Set $C := \Gamma(U, (\mathcal{O}_Z)_i) = \Gamma(f_i^{-1}(U), \mathcal{O}_{X_i})$ and $M := \Gamma(U, \mathcal{M}_i)$. There is a canonical map $(g_i|_{f_i^{-1}(U)})^\#(\tilde{M}) \rightarrow \mathcal{M}_i|_U$, where $\tilde{M}$ is the quasi-coherent sheaf over $\text{Spec } C \subset X_i$ associated with the C -module M . When we apply $i^\#$ to this map, we get $\tilde{M}_0 \rightarrow ((i^\# \mathcal{M})_i)_U$, where M_0 is M viewed as a $\Gamma(U, \mathcal{O}_{Y_i})$ -module. This is an isomorphism, since $((i^\# \mathcal{M})_i)_U$ is quasi-coherent and U is affine. As $i^\#$ is faithful and exact, we have $(g_i|_{f_i^{-1}(U)})^\#(\tilde{M}) \cong \mathcal{M}_i|_U$.

Take an exact sequence of the form

$$C^{(T)} \rightarrow C^{(\Sigma)} \rightarrow M \rightarrow 0.$$

Applying the exact functor $(g_i|_{f_i^{-1}(U)})^\# \circ ?$, we get an exact sequence of the desired type. $\square$

23.5 Corollary. *Under the same assumption as in the lemma, the functor $g_\#$ preserves local quasi-coherence.*

Proof. As $g_\#$ is compatible with restrictions, we may assume that I consists of one object and one morphism. Further, as the question is local, we may assume that $Y_\bullet = Y$ is an affine scheme. By the lemma, it suffices to show that $g_\# \mathcal{O}_Z$ is quasi-coherent, since $g_\#$ is exact and preserves direct sums. As $g_\# \mathcal{O}_Z = g_\# g^\# \mathcal{O}_X \cong \mathcal{O}_X$, we are done. $\square$

23.6 Lemma. *Let the notation be as above. The unit of adjunction $u : \mathbb{F} \rightarrow Rg^\# g_\# \mathbb{F}$ is an isomorphism for $\mathbb{F} \in D_{\text{LQco}(Z)}^+(\text{Mod}(Z))$.*

Proof. We may assume that I consists of one object and one morphism, and $Y_\bullet = Y$ is affine. By the way-out lemma, we may assume that $\mathbb{F}$ is a single quasi-coherent sheaf. By Corollary 23.5, we may assume that $\mathbb{F} = \mathcal{O}_Z = g^\# \mathcal{O}_X$ and it suffices to prove that $ug^\# : g^\# \mathcal{O}_X \rightarrow g^\# g_\# g^\# \mathcal{O}_X$ is an isomorphism. As $\text{id} = (g^\# \varepsilon)(ug^\#)$ and ε is an isomorphism, we are done. $\square$

For $\mathcal{N} \in \text{Mod}(Y_\bullet)$ and $\mathcal{M} \in \text{Mod}(Z)$, the sheaf $\underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(\mathcal{M}, \mathcal{N})$ on $Y_\bullet$ has a structure of $\mathcal{O}_Z$ -module, and it belongs to $\text{Mod}(Z)$. There is an obvious isomorphism of functors

$$\kappa : i^\# \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(\mathcal{M}, \mathcal{N}) \cong \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(i^\# \mathcal{M}, \mathcal{N}).$$

For $\mathcal{M}, \mathcal{M}' \in \text{Mod}(Z)$, there is a natural map

$$v : \underline{\text{Hom}}_{\text{Mod}(Z)}(\mathcal{M}, \mathcal{M}') \rightarrow \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(\mathcal{M}, i^\# \mathcal{M}').$$

Note that the composite

$$i^\# \underline{\text{Hom}}_{\text{Mod}(Z)}(\mathcal{M}, \mathcal{M}') \xrightarrow{i^\# v} i^\# \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(\mathcal{M}, i^\# \mathcal{M}') \xrightarrow{\kappa} \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}(i^\# \mathcal{M}, i^\# \mathcal{M}')$$

agrees with H .

(23.7) Let $I, S, f_\bullet : X_\bullet \rightarrow Y_\bullet, Z, g$, and i be as in (23.1). Assume that $f_\bullet$ is finite of fiber type, and $Y_\bullet$ has flat arrows. Define $f_\bullet^\natural : \mathcal{D}^+(Y_\bullet) \rightarrow D(\text{Mod}(X_\bullet))$ by

$$f_\bullet^\natural(\mathbb{F}) := g_\# R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_Z, \mathbb{F}).$$

As $f_\bullet$ is finite of fiber type, $\mathcal{O}_Z$ is coherent. By Lemma 13.11, $i^\# R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_Z, \mathbb{F}) \in \mathcal{D}^+(Y_\bullet)$. It follows that $f_\bullet^\natural(\mathbb{F}) \in \mathcal{D}^+(X_\bullet)$, and $f_\bullet^\natural$ is a functor from $\mathcal{D}^+(Y_\bullet)$ to $\mathcal{D}^+(X_\bullet)$.

Define $\varepsilon : R(f_\bullet)_* f_\bullet^\natural \rightarrow \text{Id}_{\mathcal{D}^+(Y_\bullet)}$ by

$$\begin{aligned} R(f_\bullet)_* f_\bullet^\natural \mathbb{F} &= i^\# R g^\# g_\# R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_Z, \mathbb{F}) \xrightarrow{u^{-1}} i^\# R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_Z, \mathbb{F}) \\ &\xrightarrow{\kappa} R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(i^\# \mathcal{O}_Z, \mathbb{F}) = R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet((f_\bullet)_* \mathcal{O}_{X_\bullet}, \mathbb{F}) \xrightarrow{\eta} R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_{Y_\bullet}, \mathbb{F}) \cong \mathbb{F}. \end{aligned}$$

Define $u : \text{Id}_{\mathcal{D}^+(X_\bullet)} \rightarrow f_\bullet^\natural R(f_\bullet)_*$ by

$$\begin{aligned} \mathbb{F} &\cong R \underline{\text{Hom}}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathcal{O}_{X_\bullet}, \mathbb{F}) \xrightarrow{\varepsilon^{-1}} g_\# R g^\# R \underline{\text{Hom}}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathcal{O}_{X_\bullet}, \mathbb{F}) \\ &\xrightarrow{H} g_\# R \underline{\text{Hom}}_{\text{Mod}(Z)}^\bullet(g^\# \mathcal{O}_{X_\bullet}, R g^\# \mathbb{F}) \xrightarrow{v} g_\# R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(\mathcal{O}_Z, R(f_\bullet)_* \mathbb{F}) = f_\bullet^\natural R(f_\bullet)_* \mathbb{F}. \end{aligned}$$

23.8 Theorem. *Let the notation be as above. Then $f_{\bullet}^{\natural}$ is right adjoint to $R(f_{\bullet})_*$, and ε and u defined above are the counit and unit of adjunction, respectively. In particular, if moreover $Y_{\bullet}$ is quasi-compact separated, then $f_{\bullet}^{\natural}$ is isomorphic to $f_{\bullet}^{\times}$.*

Proof. It is easy to see that the composite

$$\begin{aligned} R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, \mathbb{F}) &\cong R\mathbf{Hom}_{\mathbf{Mod}(Z)}^{\bullet}(\mathcal{O}_Z, R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, \mathbb{F})) \\ &\xrightarrow{v} R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, i^{\#} R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, \mathbb{F})) \xrightarrow{\kappa} R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(i^{\#}\mathcal{O}_Z, \mathbb{F})) \\ &\xrightarrow{\eta} R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_{Y_{\bullet}}, \mathbb{F})) \cong R\mathbf{Hom}_{\mathcal{O}_{Y_{\bullet}}}^{\bullet}(\mathcal{O}_Z, \mathbb{F}) \end{aligned}$$

is the identity. Utilizing this, $(f_{\bullet}^{\natural}\varepsilon) \circ (uf_{\bullet}^{\natural}) = \text{id}$ and $(\varepsilon R(f_{\bullet})_*) \circ (R(f_{\bullet})_* u) = \text{id}$ are checked directly.

The last assertion is obvious, as the right adjoint functor is unique. $\square$

24 Regular embeddings and Smooth morphisms of fiber type

(24.1) Let I be a small category, S a scheme, and $X_{\bullet} \in \mathcal{P}(I, \text{Sch}/S)$. An $\mathcal{O}_{X_{\bullet}}$ -module sheaf $\mathcal{M} \in \mathbf{Mod}(X_{\bullet})$ is said to be locally free (resp. invertible) if $\mathcal{M}$ is coherent and $\mathcal{M}_i$ is locally free (resp. invertible) for any $i \in \text{ob}(I)$. A perfect complex of $X_{\bullet}$ is a bounded complex in $C^b(\mathbf{Mod}(X_{\bullet}))$ each of whose terms is locally free.

A point of $X_{\bullet}$ is a pair (i, x) such that $i \in \text{ob}(I)$ and $x \in X_i$. A stalk of a sheaf $\mathcal{M} \in \mathbf{AB}(X_{\bullet})$ at the point (i, x) is defined to be $(\mathcal{M}_i)_x$, and we denote it by $\mathcal{M}_{i,x}$.

A connected component of $X_{\bullet}$ is an equivalence class with respect to the equivalence relation of the set of points of $X_{\bullet}$ generated by the following relations.

- 1 (i, x) and (i', x') are equivalent if $i = i'$ and x and x' belong to the same connected component of X_i .
- 2 (i, x) and (i', x') are equivalent if there exists some $\phi : i \rightarrow i'$ such that $X_{\phi}(x') = x$.

We say that $X_\bullet$ is d -connected if $X_\bullet$ consists of one connected component (note that the word ‘connected’ is reserved for componentwise connectedness). If $X_\bullet$ is locally noetherian, then a connected component of $X_\bullet$ is a closed open subdiagram of schemes in a natural way. If this is the case, the rank function $(i, x) \mapsto \text{rank}_{\mathcal{O}_{X_i, x}} \mathcal{F}_{i, x}$ of a locally free sheaf $\mathcal{F}$ is constant on a connected component of $X_\bullet$.

24.2 Lemma. *Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Let $\mathbb{F}$ be a perfect complex of $X_\bullet$. Then we have*

1 *The canonical map*

$$H_J : (?)_J R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G}) \rightarrow R\text{Hom}_{\mathcal{O}_{X_\bullet|J}}^\bullet(\mathbb{F}_J, \mathbb{G}_J)$$

is an isomorphism for $\mathbb{G} \in D(\text{Mod}(X_\bullet))$.

2 *The canonical map*

$$R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G}) \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{H} \rightarrow R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{H})$$

is an isomorphism for $\mathbb{G}, \mathbb{H} \in D(\text{Mod}(X_\bullet))$.

Proof. **1** It suffices to show that

$$H_J : (?)_J \text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G}) \rightarrow \text{Hom}_{\mathcal{O}_{X_\bullet|J}}^\bullet(\mathbb{F}_J, \mathbb{G}_J)$$

is an isomorphism of complexes if $\mathbb{G}$ is a K -injective complex in $C(\text{Mod}(X_\bullet))$, since $\mathbb{F}_J$ is K -flat and $\mathbb{G}_J$ is weakly K -injective. The assertion follows immediately by Lemma 7.28.

2 We may assume that $\mathbb{F}$ is a single locally free sheaf. By **1**, we may assume that $X = X_\bullet$ is a single scheme. We may assume that X is affine and $\mathbb{F} = \mathcal{O}_X^n$ for some n . This case is trivial. $\square$

(24.3) Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. An $\mathcal{O}_{X_\bullet}$ -module $\mathcal{M}$ is said to be locally of finite projective dimension if $\mathcal{M}_{i, x}$ is of finite projective dimension as an $\mathcal{O}_{X_i, x}$ -module for any point (i, x) of $X_\bullet$. We say that $\mathcal{M}$ has finite projective dimension if there exists some non-negative integer d such that $\text{proj.dim}_{\mathcal{O}_{X_i, x}} \mathcal{M}_{i, x} \leq d$ for any point (i, x) of $X_\bullet$.

24.4 Lemma. *Let I be a small category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$ has flat arrows and is locally noetherian. If $\mathbb{F}$ is a complex*

in $C(\text{Mod}(X_\bullet))$ with bounded coherent cohomology groups which have finite projective dimension, then the canonical map

$$R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G}) \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{H} \rightarrow R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G} \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} \mathbb{H})$$

is an isomorphism for $\mathbb{G}, \mathbb{H} \in D(\text{Mod}(X_\bullet))$.

Proof. We may assume that $\mathbb{G} = \mathcal{O}_{X_\bullet}$. By the way-out lemma, we may assume that $\mathbb{F}$ is a single coherent sheaf which has finite projective dimension, say d . By Lemma 13.10, it is easy to see that $\text{Ext}_{\mathcal{O}_{X_\bullet}}^i(\mathbb{F}, G) = 0$ ($i > d$) for $G \in \text{Mod}(X_\bullet)$. In particular, $R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, ?)$ is way-out in both directions. On the other hand, as $R\text{Hom}_{\mathcal{O}_{X_\bullet}}(\mathbb{F}, \mathcal{O}_{X_\bullet})$ has finite flat dimension, and hence $R\text{Hom}_{\mathcal{O}_{X_\bullet}}(\mathbb{F}, \mathcal{O}_{X_\bullet}) \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet, L} ?$ is also way-out in both directions. By the way-out lemma, we may assume that $\mathbb{H}$ is a single $\mathcal{O}_{X_\bullet}$ -module. By Lemma 13.9, we may assume that $X = X_\bullet$ is a single scheme. The question is local, and we may assume that $X = \text{Spec } A$ is affine. Moreover, we may assume that $\mathbb{F}$ is a complex of sheaves associated with a finite projective resolution of a single finitely generated module. As $\mathbb{F}$ is perfect, the result follows from Lemma 24.2. $\square$

(24.5) Let S , I , and $X_\bullet$ be as above. For a locally free sheaf $\mathcal{F}$ over $X_\bullet$, we denote $\text{Hom}_{\mathcal{O}_{X_\bullet}}(\mathcal{F}, \mathcal{O}_{X_\bullet})$ by $\mathcal{F}^\vee$. It is easy to see that $\mathcal{F}^\vee$ is again locally free. If $\mathcal{L}$ is an invertible sheaf, then

$$\mathcal{O}_{X_\bullet} \xrightarrow{\text{tr}} \text{Hom}_{\mathcal{O}_{X_\bullet}}(\mathcal{L}, \mathcal{L}) \cong \mathcal{L}^\vee \otimes_{\mathcal{O}_{X_\bullet}} \mathcal{L}$$

are isomorphisms.

(24.6) Let I be a small category, S a scheme, and $i_\bullet : Y_\bullet \rightarrow X_\bullet$ a closed immersion in $\mathcal{P}(I, \text{Sch}/S)$. Then the canonical map $\eta : \mathcal{O}_{X_\bullet} \rightarrow (i_\bullet)_* \mathcal{O}_{Y_\bullet}$ is an epimorphism in $\text{LQco}(X_\bullet)$. Set $\mathcal{I} := \text{Ker } \eta$. Then $\mathcal{I}$ is a locally quasi-coherent ideal of $\mathcal{O}_{X_\bullet}$. Conversely, if $\mathcal{I}$ is a given locally quasi-coherent ideal of $\mathcal{O}_{X_\bullet}$, then

$$Y_\bullet := \underline{\text{Spec}}_\bullet \mathcal{O}_{X_\bullet} / \mathcal{I} \xrightarrow{i_\bullet} X_\bullet$$

is defined appropriately, and $i_\bullet$ is a closed immersion. Thus the isomorphism classes of closed immersions to $X_\bullet$ in the category $\mathcal{P}(I, \text{Sch}/S)/X_\bullet$ and locally quasi-coherent ideals of $\mathcal{O}_{X_\bullet}$ are in one-to-one correspondence. We call $\mathcal{I}$ the defining ideal sheaf of $Y_\bullet$.

Obviously, $i_\bullet$ is of fiber type if and only if $\mathcal{I}$ is equivariant.

(24.7) Let $X_\bullet$ be locally noetherian. A morphism $i_\bullet : Y_\bullet \rightarrow X_\bullet$ is said to be a regular embedding, if $i_\bullet$ is a closed immersion of fiber type such that $i_j : Y_j \rightarrow X_j$ is a regular embedding for each $j \in \text{ob}(I)$, or equivalently, $\mathcal{I}$ is coherent and $\mathcal{I}_{j,x}$ is a complete intersection ideal of $\mathcal{O}_{X_j,x}$ for any $j \in \text{ob}(I)$ and $x \in X_j$. If this is the case, we say that $\mathcal{I}$ is a local complete intersection ideal sheaf.

A closed immersion of fiber type $i_\bullet : Y_\bullet \rightarrow X_\bullet$ with $X_\bullet$ locally noetherian is a regular embedding if and only if $i_\bullet^* \mathcal{I}$ is locally free and $\mathcal{I}_{i,x}$ is of finite projective dimension as an $\mathcal{O}_{X_i,x}$ -module.

Note that $i_\bullet^* \mathcal{I} \cong \mathcal{I}/\mathcal{I}^2$, and we have

$$\text{ht}_{\mathcal{O}_{X_i,x}} \mathcal{I}_x = \text{rank}_{\mathcal{O}_{Y_i,y}} (i_\bullet^* \mathcal{I})_{i,y}$$

for any point (i, y) of $Y_\bullet$, where $x = i_i(y)$. We call these numbers the codimension of $\mathcal{I}$ at (i, y) .

24.8 Proposition. *Let I be a small category, S a scheme, and $i_\bullet : Y_\bullet \rightarrow X_\bullet$ a morphism in $\mathcal{P}(I, \text{Sch}/S)$. Assume that $X_\bullet$ is locally noetherian with flat arrows and $i_\bullet$ is a regular embedding. Let $\mathcal{I}$ be the defining ideal of $Y_\bullet$, and assume that $Y_\bullet$ has a constant codimension d . Then we have the following.*

1 $\underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^i((i_\bullet)_* \mathcal{O}_{Y_\bullet}, \mathcal{O}_{X_\bullet}) = 0$ for $i \neq d$.

2 The canonical map

$$\underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^d((i_\bullet)_* \mathcal{O}_{Y_\bullet}, \mathcal{O}_{X_\bullet}) \rightarrow \underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^d((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet})$$

is an isomorphism.

3 The Yoneda algebra

$$\underline{\text{Ext}}_{\mathcal{O}_{Y_\bullet}}^\bullet((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}) := \bigoplus_{j \geq 0} \underline{\text{Ext}}_{\mathcal{O}_{Y_\bullet}}^j((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet})$$

is isomorphic to the exterior algebra $(i_\bullet)_* \bigwedge^\bullet (i_\bullet^* \mathcal{I})^\vee$ as graded $\mathcal{O}_{X_\bullet}$ -algebras.

4 There is an isomorphism

$$i_\bullet^\sharp \mathcal{O}_{X_\bullet} \cong \bigwedge^d (i_\bullet^* \mathcal{I})^\vee [-d].$$

5 For $\mathbb{F} \in \mathcal{D}^+(X_\bullet)$, there is a functorial isomorphism

$$i_\bullet^! \mathbb{F} \cong \bigwedge^d (i_\bullet^* \mathcal{I})^\vee \otimes_{\mathcal{O}_{Y_\bullet}}^{\bullet, L} Li_\bullet^* \mathbb{F}[-d].$$

Proof. 1 is trivial, since $\mathcal{I}_{i,x}$ is a complete intersection ideal of the local ring $\mathcal{O}_{X_{i,x}}$ of codimension d for any point (i, x) of $X_\bullet$.

2 Note that $(i_\bullet)_* \mathcal{O}_{Y_\bullet} \cong \mathcal{O}_{X_\bullet} / \mathcal{I}$. From the short exact sequence

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_{X_\bullet} \rightarrow \mathcal{O}_{X_\bullet} / \mathcal{I} \rightarrow 0,$$

we get an isomorphism

$$\underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^1((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}) \cong \underline{\mathrm{Hom}}_{\mathcal{O}_{X_\bullet}}(\mathcal{I}, \mathcal{O}_{X_\bullet} / \mathcal{I}) \cong (i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee.$$

The canonical map

$$(i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee \cong \underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^1((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}) \hookrightarrow \underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^\bullet((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet})$$

is uniquely extended to an $\mathcal{O}_{X_\bullet}$ -algebra map

$$T_\bullet((i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee) \rightarrow \underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^\bullet((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}),$$

where $T_\bullet$ denotes the tensor algebra. It suffices to prove that this map is an epimorphism, which induces an isomorphism

$$\bigwedge^\bullet((i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee) \rightarrow \underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^\bullet((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}).$$

In fact, the exterior algebra is compatible with base change, and

$$\begin{aligned} \bigwedge^\bullet((i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee) &\cong (i_\bullet)_* i_\bullet^* \bigwedge^\bullet((i_\bullet)_*(i_\bullet^* \mathcal{I})^\vee) \\ &\cong (i_\bullet)_* \bigwedge^\bullet((i_\bullet^* (i_\bullet)_*)(i_\bullet^* \mathcal{I})^\vee) \cong (i_\bullet)_* \bigwedge^\bullet(i_\bullet^* \mathcal{I})^\vee. \end{aligned}$$

To verify this, we may assume that $i_\bullet : Y_\bullet \rightarrow X_\bullet$ is a morphism of single schemes, $X_\bullet = \mathrm{Spec} A$ affine, and $\mathcal{I} = \tilde{I}$ generated by an A -sequence. The proof for this case is essentially the same as [17, Lemma IV.1.1.8], and we omit it.

4 Let Z denote the ringed site $(\mathrm{Zar}(X_\bullet), (i_\bullet)_* \mathcal{O}_{Y_\bullet})$, and $g : Z \rightarrow \mathrm{Zar}(Y_\bullet)$ the associated admissible ringed continuous functor. By 2–3, there is a sequence of isomorphisms of $\mathrm{Coh}(X_\bullet)$

$$(i_\bullet)_* \bigwedge^d (i_\bullet^* \mathcal{I})^\vee \cong \underline{\mathrm{Ext}}_{\mathcal{O}_{Y_\bullet}}^d((i_\bullet)_* \mathcal{O}_{Y_\bullet}, (i_\bullet)_* \mathcal{O}_{Y_\bullet}) \cong \underline{\mathrm{Ext}}_{\mathcal{O}_{X_\bullet}}^d((i_\bullet)_* \mathcal{O}_{Y_\bullet}, \mathcal{O}_{X_\bullet}).$$

In view of **1**, there is an isomorphism

$$R(g_{\bullet})^{\#} \bigwedge^d (i_{\bullet}^* \mathcal{I})^{\vee} \cong R \underline{\mathrm{Hom}}_{\mathcal{O}_{X_{\bullet}}}^{\bullet}(\mathcal{O}_Z, \mathcal{O}_{X_{\bullet}})[d]$$

in $D_{\mathrm{Coh}}^b(\mathrm{Mod}(Z))$. Applying $(g_{\bullet})_{\#}$ to both sides, we get

$$\bigwedge^d (i_{\bullet}^* \mathcal{I})^{\vee} \cong i_{\bullet}^! \mathcal{O}_{X_{\bullet}}[d].$$

5 is an immediate consequence of **4** and Lemma 24.4. $\square$

(24.9) Let I and S be as in (24.6). Let $f_{\bullet} : X_{\bullet} \rightarrow Y_{\bullet}$ be a morphism in $\mathcal{P}(I, \mathrm{Sch}/S)$. Assume that $f_{\bullet}$ is separated so that the diagonal $\Delta_{X_{\bullet}/Y_{\bullet}} : X_{\bullet} \rightarrow X_{\bullet} \times_{Y_{\bullet}} X_{\bullet}$ is a closed immersion. Define $\Omega_{X_{\bullet}/Y_{\bullet}} := i_{\bullet}^* \mathcal{I}$, where $\mathcal{I} := \mathrm{Ker}(\eta : \mathcal{O}_{X_{\bullet} \times_{Y_{\bullet}} X_{\bullet}} \rightarrow (\Delta_{X_{\bullet}/Y_{\bullet}})_* \mathcal{O}_{X_{\bullet}})$. Note that $(\Delta_{X_{\bullet}/Y_{\bullet}})_* \Omega_{X_{\bullet}/Y_{\bullet}} \cong \mathcal{I}/\mathcal{I}^2$.

24.10 Lemma. *Let the notation be as above. Then we have*

1 $\Omega_{X_{\bullet}/Y_{\bullet}}$ is locally quasi-coherent.

2 If $f_{\bullet}$ is of fiber type, then $\Omega_{X_{\bullet}/Y_{\bullet}}$ is quasi-coherent.

3 For $i \in \mathrm{ob}(I)$, there is a canonical isomorphism $\Omega_{X_i/Y_i} \cong (\Omega_{X_{\bullet}/Y_{\bullet}})_i$.

Proof. Easy. $\square$

24.11 Theorem. *Let I be a finite category, S a separated noetherian scheme, and $f_{\bullet} : X_{\bullet} \rightarrow Y_{\bullet}$ a morphism in $\mathcal{P}(I, \mathrm{Sch}/S)$. Assume that $Y_{\bullet}$ is separated of finite type over S , and $f_{\bullet}$ is separated smooth of finite type and fiber type. Assume that $f_{\bullet}$ has a constant relative dimension d . Then for any $\mathbb{F} \in D_{\mathrm{LQco}(Y_{\bullet})}^+(\mathrm{Mod}(X_{\bullet}))$, there is a functorial isomorphism*

$$\bigwedge^d \Omega_{X_{\bullet}/Y_{\bullet}}[d] \otimes_{\mathcal{O}_{X_{\bullet}}}^{\bullet} f_{\bullet}^* \mathbb{F} \cong f_{\bullet}^! \mathbb{F},$$

where $[d]$ denotes the shift of degree.

Proof. In view of Theorem 22.14, it suffices to show that there is an isomorphism $f_{\bullet}^! \mathcal{O}_{Y_{\bullet}} \cong \bigwedge^d \Omega_{X_{\bullet}/Y_{\bullet}}[d]$. Consider the commutative diagram

$$\begin{array}{ccccc} X_{\bullet} & \xrightarrow{\Delta} & X_{\bullet} \times_{Y_{\bullet}} X_{\bullet} & \xrightarrow{p_2} & X_{\bullet} \\ & \searrow \mathrm{id} & \downarrow p_1 & & \downarrow f_{\bullet} \\ & & X_{\bullet} & \xrightarrow{f_{\bullet}} & Y_{\bullet} \end{array}$$

By Lemma 8.16 and Lemma 8.15, all the morphisms in the diagrams are of fiber type. As p_1 is smooth of finite type of relative dimension d , Δ is a regular embedding of the constant codimension d .

By Theorem 17.8, Theorem 23.8, and Proposition 24.8, we have

$$\begin{aligned} \mathcal{O}_{X_\bullet} &\cong f_\bullet^* \mathcal{O}_{Y_\bullet} \cong \Delta^! p_1^! f_\bullet^* \mathcal{O}_{Y_\bullet} \cong \Delta^! p_2^* f_\bullet^! \mathcal{O}_{Y_\bullet} \cong \\ &\bigwedge^d \Omega_{X_\bullet/Y_\bullet}^\vee[-d] \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet L} L\Delta^* p_2^* f_\bullet^! \mathcal{O}_{Y_\bullet} \cong \bigwedge^d \Omega_{X_\bullet/Y_\bullet}^\vee[-d] \otimes_{\mathcal{O}_{X_\bullet}}^{\bullet L} f_\bullet^! \mathcal{O}_{Y_\bullet}. \end{aligned}$$

As $\bigwedge^d \Omega_{X_\bullet/Y_\bullet}$ is an invertible sheaf, we are done. $\square$

25 Simplicial objects

(25.1) For $n \in \mathbb{Z}$ with $n \geq -1$, we define $[n]$ to be the totally ordered finite set $\{0 < 1 < \dots < n\}$. Thus, $[-1] = \emptyset$, $[0] = \{0\}$, $[1] = \{0 < 1\}$, and so on. We define (Δ^+) to be the small category given by $\text{ob}(\Delta^+) := \{[n] \mid n \in \mathbb{Z}, n \geq -1\}$ and

$$\text{Mor}(\Delta^+) := \{\text{monotone maps}\}.$$

For a subset S of $\{-1, 0, 1, \dots\}$, we define $(\Delta^+)_S$ to be the full subcategory of (Δ^+) such that $\text{ob}((\Delta^+)_S) = \{[n] \mid n \in S\}$. We define $(\Delta) := (\Delta^+)_{[0, \infty)}$. If $-1 \notin S$, then $(\Delta^+)_S$ is also denoted by $(\Delta)_S$.

We define $(\Delta^+)^{\text{mon}}$ to be the subcategory of (Δ^+) by $\text{ob}((\Delta^+)^{\text{mon}}) := \text{ob}(\Delta^+)$ and

$$\text{Mor}((\Delta^+)^{\text{mon}}) := \{\text{injective monotone maps}\}.$$

For $S \subset \{-1, 0, 1, \dots\}$, the full subcategories $(\Delta^+)_S^{\text{mon}}$ and $(\Delta)_S^{\text{mon}}$ of $(\Delta^+)^{\text{mon}}$ are defined similarly.

We denote $(\Delta)_{\{0,1,2\}}^{\text{mon}}$ and $(\Delta^+)_{\{-1,0,1,2\}}^{\text{mon}}$ by Δ_M and Δ_M^+ , respectively.

Let $\mathcal{C}$ be a category. We call an object of $\mathcal{P}((\Delta^+), \mathcal{C})$ (resp. $\mathcal{P}((\Delta), \mathcal{C})$), an augmented simplicial object (resp. simplicial object) of $\mathcal{C}$.

For a subcategory $\mathcal{D}$ of (Δ^+) and an object $X_\bullet \in \mathcal{P}(\mathcal{D}, \mathcal{C})$, we denote $X_{[n]}$ by X_n .

As $[-1]$ is the initial object of Δ^+ , an augmented simplicial object $X_\bullet$ of $\mathcal{C}$ with $X_{-1} = c$ is identified with a simplicial object of $\mathcal{C}/c$.

We define some particular morphisms in $\tilde{\Delta}$. The unique map $[-1] \rightarrow [n]$ is denoted by $\varepsilon(n)$. The unique injective monotone map $[n-1] \rightarrow [n]$ such that i is not in the image is denoted by $\delta_i(n)$ for $i \in [n]$. The unique surjective

monotone map $[n+1] \rightarrow [n]$ such that i has two inverse images is denoted by $\sigma_i(n)$ for $i \in [n]$. The unique map $[0] \rightarrow [n]$ such that i is in the image is denoted by $\rho_i(n)$. The unique map $[n] \rightarrow [0]$ is denoted by λ_n .

Let $\mathcal{D}$ be a subcategory of (Δ^+) . For $X_\bullet \in \mathcal{P}(\mathcal{D}, \mathcal{C})$, we denote $X_\bullet(\varepsilon(n))$ (resp. $X_\bullet(\delta_i(n))$, $X_\bullet(\sigma_i(n))$, $X_\bullet(\rho_i(n))$, and $X_\bullet(\lambda_n)$) by $e(n, X_\bullet)$ (resp. $d_i(n, X_\bullet)$, $s_i(n, X_\bullet)$, $r_i(n, X_\bullet)$, and $l_n(X_\bullet)$), or simply by $e(n)$ (resp. $d_i(n)$, $s_i(n)$, $r_i(n)$, l_n), if there is no confusion.

Note that Δ is generated by $\delta_i(n)$, $\sigma_i(n)$ for various i and n .

(25.2) Note that $\Delta^+([m], [n])$ is a finite set for any m, n . Assume that $\mathcal{C}$ has finite limits and let $f : X \rightarrow Y$ be a morphism in $\mathcal{C}$. Then the *Čech nerve* is defined to be $\text{Nerve}(f) := \text{cosk}_{(\Delta^+)_{\{0,1\}}}^+(f)$, where $\text{cosk}_{(\Delta^+)_{\{0,1\}}}^+$ is the right adjoint of the restriction, which exists. It is described as follows. $\text{Nerve}(f)_n = X \times_Y \times \cdots \times_Y X$ ($(n+1)$ -fold fiber product) for $n \geq 0$, and $\text{Nerve}(f)_{-1} = Y$. Note that $d_i(n)$ is given by

$$d_i(n)(x_n, \dots, x_1, x_0) = (x_n, \overset{i}{\cdots}, x_1, x_0),$$

and $s_i(n)$ is given by

$$s_i(n)(x_n, \dots, x_1, x_0) = (x_n, \dots, x_{i+1}, x_i, x_i, x_{i-1}, \dots, x_1, x_0)$$

if $\mathcal{C} = \underline{\text{Set}}$.

(25.3) Let S be a scheme. A simplicial object (resp. augmented simplicial object) in Sch/S , in other words, an object of $\mathcal{P}((\Delta), \text{Sch}/S)$ (resp. $\mathcal{P}((\Delta^+), \text{Sch}/S)$), is called a simplicial (resp. augmented simplicial) S -scheme. The following is well-known.

25.4 Lemma. *Let $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$. Then the restriction $(?)_{\Delta_M} : \text{EM}(X_\bullet) \rightarrow \text{EM}(X_\bullet|_{\Delta_M})$ is an equivalence. With the equivalence, quasi-coherent sheaves correspond to quasi-coherent sheaves.*

Proof. We define a third category $\mathcal{A}$ as follows. An object of $\mathcal{A}$ is a pair $(\mathcal{M}_0, \varphi)$ such that, $\mathcal{M}_0 \in \text{Mod}(X_0)$, $\varphi \in \text{Hom}_{\text{Mod}(X_1)}(d_0^*(\mathcal{M}_0), d_1^*(\mathcal{M}_0))$, φ an isomorphism, and that $d_1^*(\varphi) = d_2^*(\varphi) \circ d_0^*(\varphi)$. Note that applying l_2^* to the last equality, we get $l_1^*(\varphi) = l_1^*(\varphi) \circ l_1^*(\varphi)$. As φ is an isomorphism, we get $l_1^*(\varphi) = \text{id}$.

A morphism $\gamma_0 : (\mathcal{M}_0, \varphi) \rightarrow (\mathcal{N}_0, \psi)$ is an element of

$$\gamma_0 \in \text{Hom}_{\text{Mod}(X_0)}(\mathcal{M}_0, \mathcal{N}_0)$$

such that

$$\psi \circ d_0^*(\gamma_0) = d_1^*(\gamma_0) \circ \varphi.$$

We define a functor $\Phi : \text{EM}(X_\bullet|_{\Delta_M}) \rightarrow \mathcal{A}$ by

$$\Phi(\mathcal{M}) := (\mathcal{M}_0, \alpha_{d_1(1)}^{-1} \circ \alpha_{d_0(1)}).$$

It is easy to verify that this gives a well-defined functor.

Now we define a functor $\Psi : \mathcal{A} \rightarrow \text{EM}(X_\bullet)$. Note that an object $\mathcal{M}$ of $\text{EM}(X_\bullet)$ is identified with a family $(\mathcal{M}_n, \alpha_w)_{[n] \in (\Delta), w \in \text{Mor}((\Delta))}$ such that $\mathcal{M}_n \in \text{Mod}(X_n)$,

$$\alpha_w \in \text{Hom}_{\text{Mod}(X_n)}((X_w)_{\text{Mod}}^*(\mathcal{M}_m), \mathcal{M}_n)$$

for $w \in \Delta(m, n)$, α_w is an isomorphism, and

$$\alpha_{ww'} = \alpha_{w'} \circ X_{w'}^* \alpha_w \quad (25.5)$$

whenever ww' is defined, see (5.6).

For $(\mathcal{M}_0, \varphi) \in \mathcal{A}$, we define $\mathcal{M}_{n,i} := (r_i(n))^*(\mathcal{M}_0)$, and $\mathcal{M}_n := \mathcal{M}_{n,0}$ (as we have $r_0(0) = \text{id}$, this will not cause a double-meaning of $\mathcal{M}_0$) for $n \geq 0$ and $0 \leq i \leq n$. We define $\psi_i(n) : \mathcal{M}_{n,i+1} \rightarrow \mathcal{M}_{n,i}$ to be $(X_{q(i,n)})^*(\varphi)$ for $n \geq 1$ and $0 \leq i < n$, where $q(i, n) : [1] \rightarrow [n]$ is the unique injective monotone map with $\{i, i+1\} = \text{Im } q(i, n)$. We define $\varphi_i(n) : \mathcal{M}_{n,i} \cong \mathcal{M}_n$ to be the composite map

$$\varphi_i(n) := \psi_0(n) \circ \psi_1(n) \circ \cdots \circ \psi_{i-1}(n)$$

for $n \geq 0$ and $0 \leq i \leq n$.

Now we define

$$\alpha_w \in \text{Hom}_{\text{Mod}(X_n)}(X_w^*(\mathcal{M}_m), \mathcal{M}_n)$$

to be the map

$$X_w^* \mathcal{M}_m = X_w^* r_0(m)^* \mathcal{M}_0 = r_{w(0)}(n)^* \mathcal{M}_0 = \mathcal{M}_{n,w(0)} \xrightarrow{\varphi_{w(0)}(n)} \mathcal{M}_n$$

for $w \in \Delta(m, n)$.

Thus $(\mathcal{M}_0, \varphi)$ yields a family $(\mathcal{M}_n, \alpha_w)$, and this gives the definition of $\Psi : \mathcal{A} \rightarrow \text{EM}(X_\bullet)$. The details of the proof of the well-definedness is left to the reader.

It is also straightforward to check that $(?)_{\Delta_M}$, Φ , and Ψ give the equivalence of these three categories. The proof is also left to the reader.

The last assertion is obvious from the construction. $\square$

26 Descent theory

Let S be a scheme.

(26.1) Consider the functor $\text{shift} : (\Delta^+) \rightarrow (\Delta)$ given by $\text{shift}[n] := [n+1]$, $\text{shift}(\delta_i(n)) := \delta_{i+1}(n+1)$, $\text{shift}(\sigma_i(n)) := \sigma_{i+1}(n+1)$, and $\text{shift}(\varepsilon(0)) := \delta_1(1)$. We have a natural transformation $(\delta_0^+) : \text{Id}_{(\Delta^+)} \rightarrow \iota \circ \text{shift}$ given by $(\delta_0^+)_n := \delta_0(n+1)$ for $n \geq 0$ and $(\delta_0^+)_{-1} := \varepsilon(0)$, where $\iota : (\Delta) \hookrightarrow (\Delta^+)$ is the inclusion. We denote $(\delta_0^+)_\iota$ by (δ_0) . Note that (δ_0) can be viewed as a natural map $(\delta_0) : \text{Id}_{(\Delta)} \rightarrow \text{shift} \iota$.

Let $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$. We define $X'_\bullet$ to be the augmented simplicial scheme $\text{shift}^\#(X_\bullet) = X_\bullet \cdot \text{shift}$. The natural map

$$X_\bullet(\delta_0) : X'_\bullet|_{(\Delta)} = X_\bullet \cdot \text{shift} \iota \rightarrow X_\bullet$$

is denoted by $(d_0)(X_\bullet)$ or (d_0) . Similarly, if $Y_\bullet \in \mathcal{P}((\Delta^+), \text{Sch}/S)$, then

$$(d_0^+)(Y_\bullet) : (Y_\bullet|_{(\Delta)})' = Y_\bullet \cdot \text{shift} \xrightarrow{Y_\bullet(\delta_0^+)} Y_\bullet$$

is defined as well.

(26.2) We say that $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$ is a simplicial groupoid of S -schemes if there is a faithfully flat morphism of S -schemes $g : Z \rightarrow Y$ such that there is a faithfully flat morphism $f_\bullet : Z_\bullet \rightarrow X_\bullet$ of fiber type of $\mathcal{P}((\Delta), \text{Sch}/S)$, where $Z_\bullet = \text{Nerve}(g)|_{(\Delta)}$.

26.3 Lemma. *Let $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$.*

- 1 *If $X_\bullet \cong \text{Nerve}(g)|_{(\Delta)}$ for some faithfully flat morphism g of S -schemes, then $X_\bullet$ is a simplicial groupoid.*
- 2 *If $f_\bullet : Z_\bullet \rightarrow X_\bullet$ is a faithfully flat morphism of fiber type of simplicial S -schemes and $Z_\bullet$ is a simplicial groupoid, then we have $X_\bullet$ is also a simplicial groupoid.*
- 3 *$X_\bullet$ is a simplicial groupoid if and only if $(d_0) : X'_\bullet|_{(\Delta)} \rightarrow X_\bullet$ is of fiber type, the canonical unit map*

$$X'_\bullet \rightarrow \text{Nerve}(d_1(1))$$

is an isomorphism, and $d_0(1)$ and $d_1(1)$ are flat.

- 4 If $f_\bullet : Z_\bullet \rightarrow X_\bullet$ is a morphism of fiber type of simplicial S -schemes and $X_\bullet$ is a simplicial groupoid, then $Z_\bullet$ is a simplicial groupoid.
- 5 A simplicial groupoid has flat $(\Delta)^{\text{mon}}$ -arrows.
- 6 If $X_\bullet$ is a simplicial groupoid of S -schemes such that $d_0(1)$ and $d_1(1)$ are separated (resp. quasi-compact, of finite type, smooth, étale), then $X_\bullet$ has separated (resp. quasi-compact, of finite type, smooth, étale) $(\Delta)^{\text{mon}}$ -arrows, and $(d_0) : X'_\bullet|_{(\Delta)} \rightarrow X_\bullet$ is separated (resp. quasi-compact, of finite type, smooth, étale).

Proof. **1** and **2** are obvious by definition.

We prove **3**. We prove the ‘if’ part. As $d_0(1)s_0(0) = \text{id} = d_1(1)s_0(0)$, we have that $d_0(1)$ and $d_1(1)$ are faithfully flat by assumption. As $d_0(1) = (d_0)_0$ is faithfully flat and (d_0) is of fiber type, it is easy to see that (d_0) is also faithfully flat. So this direction is obvious.

We prove the ‘only if’ part. As $X_\bullet$ is a simplicial groupoid, there is a faithfully flat S -morphism $g : Z \rightarrow Y$ and a faithfully flat morphism $f_\bullet : Z_\bullet \rightarrow X_\bullet$ of simplicial S -schemes of fiber type, where $Z_\bullet = \text{Nerve}(g)|_{(\Delta)}$. It is easy to see that $(d_0) : Z'_\bullet|_{(\Delta)} \rightarrow Z_\bullet$ is nothing but the base change by g , and it is faithfully flat of fiber type. It is also obvious that $Z'_\bullet \cong \text{Nerve}(d_1(1)(Z_\bullet))$ and $d_0(1)(Z_\bullet)$ and $d_1(1)(Z_\bullet)$ are flat. It is obvious that $f'_\bullet : Z'_\bullet \rightarrow X'_\bullet$ is faithfully flat of fiber type. Now by Lemma 8.15, $(d_0)(X_\bullet)$ is of fiber type. As $f_\bullet$ is faithfully flat of fiber type and $d_0(1)(Z_\bullet)$ and $d_1(1)(Z_\bullet)$ are flat, we have that $d_0(1)(X_\bullet)$ and $d_1(1)(X_\bullet)$ are flat. When we base change $X'_\bullet \rightarrow \text{Nerve}(d_1(1)(X_\bullet))$ by $f_0 : Z_0 \rightarrow X_0$, then we have the isomorphism $Z'_\bullet \cong \text{Nerve}(d_1(1)(Z_\bullet))$. As f_0 is faithfully flat, we have that $X'_\bullet \rightarrow \text{Nerve}(d_1(1))$ is also an isomorphism.

The assertions **4**, **5** and **6** are proved easily. $\square$

(26.4) Let $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$. Then we define $F : \text{Zar}(X'_\bullet) \rightarrow \text{Zar}(X_\bullet)$ by $F(i, U) = (\text{shift } i, U)$. The corresponding pull-back $F_{\text{Mod}}^\#$ is denoted by $(?)'$. It is easy to see that $(?)'$ have a left and a right adjoint. It also preserves equivariant and locally quasi-coherent sheaves.

Let $\mathcal{M} \in \text{Mod}(X_\bullet)$. Then we define $(\alpha) : (d_0)^*\mathcal{M} \rightarrow \mathcal{M}'_{(\Delta)}$ by

$$(\alpha)_n : ((d_0)^*\mathcal{M})_n = d_0(n)^*\mathcal{M}_n \xrightarrow{\alpha_{\delta_0(n)}} \mathcal{M}_{n+1} = \mathcal{M}'_n.$$

It is easy to see that $(\alpha) : (d_0)^* \rightarrow (?)_{(\Delta)} \circ (?)'$ is a natural map. Similarly, for $Y_\bullet \in \mathcal{P}((\Delta^+), \text{Sch}/S)$,

$$(\alpha^+) : (d_0^+)^* \rightarrow (?)' \circ (?)_{(\Delta)}$$

is defined.

(26.5) Let $X_\bullet \in \mathcal{P}((\Delta^+), \text{Sch}/S)$, and $\mathcal{M} \in \text{Mod}(X_\bullet|_{(\Delta)})$. Then, we have a cosimplicial object $\text{Cos}(\mathcal{M})$ of $\text{Mod}(X_{-1})$ (i.e., a simplicial object of $\text{Mod}(X_{-1})^{\text{op}}$). We have $\text{Cos}(\mathcal{M})_n := e(n)_*(\mathcal{M}_n)$, and

$$w(\text{Cos}(\mathcal{M})) := e(m)_*(\beta_w) : e(m)_*(\mathcal{M}_m) \rightarrow e(m)_*(w_*(\mathcal{M}_n)) \cong e(n)_*(\mathcal{M}_n)$$

for a morphism $w : m \rightarrow n$ in Δ . Similarly, the augmented cosimplicial object $\text{Cos}^+(\mathcal{N})$ of $\text{Mod}(X_{-1})$ is defined for $\mathcal{N} \in \text{Mod}(X_\bullet)$.

By (7.10), it is easy to see that $\mathcal{M}^+ := R_{(\Delta)}^{\text{Mod}}(\mathcal{M})$ is $\mathcal{M}$ on $X_\bullet|_{(\Delta)}$, $\mathcal{M}_{-1}^+$ is $\varprojlim \text{Cos}(\mathcal{M})$, and $\beta_{\varepsilon(n)}(\mathcal{M}^+)$ is nothing but the canonical map

$$\varprojlim \text{Cos}(\mathcal{M}) \rightarrow \text{Cos}(\mathcal{M})_n = e(n)_*(\mathcal{M}_n) = e(n)_*(\mathcal{M}_n^+).$$

Note also that we have

$$\varprojlim \text{Cos}(\mathcal{M}) = \text{Ker}(e(0)_*\beta_{\delta_0(1)} - e(0)_*\beta_{\delta_1(1)}), \quad (26.6)$$

which is determined only by $\mathcal{M}_{(\Delta)_{\{0,1\}}}$.

26.7 Lemma. *Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism of $\mathcal{P}((\Delta^+), \text{Sch}/S)$. If $f_\bullet|_{(\Delta^+)_{\{-1,0,1\}}}$ is of fiber type, flat quasi-compact separated and $Y_\bullet$ has flat $(\Delta^+)_{\{-1,0,1\}}$ -arrows, then the canonical map*

$$f_\bullet^* \circ R_{(\Delta)} \rightarrow R_{(\Delta)} \circ (f_\bullet|_{(\Delta)})^*$$

is an isomorphism of functors from $\text{LQco}(Y_\bullet|_{(\Delta)})$ to $\text{LQco}(X_\bullet)$.

Proof. To prove that the map in question is an isomorphism, it suffices to show that the map is an isomorphism after applying the functor $(?)_n$ for $n \geq -1$. This is trivial if $n \geq 0$. On the other hand, if $n = -1$, the map restricted at -1 and evaluated by $\mathcal{M} \in \text{LQco}(Y_\bullet)$ is nothing but

$$f_{-1}^*(H^0(\text{Cos}(\mathcal{M}))) \cong H^0(f_{-1}^*(\text{Cos}(\mathcal{M}))) \rightarrow H^0(\text{Cos}((f_\bullet)^*(\mathcal{M}))).$$

The first map is an isomorphism as f_{-1} is flat. Although the map $f_{-1}^*(\text{Cos}(\mathcal{M})) \rightarrow \text{Cos}((f_\bullet)^*(\mathcal{M}))$ may not be a chain isomorphism, it is an isomorphism at the degrees $-1, 0, 1$, and it induces the isomorphism of H^0 . $\square$

26.8 Lemma. *Let $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$, and $\mathcal{M} \in \text{Mod}(X_\bullet)$. Then the (associated chain complex of the) augmented cosimplicial object $\text{Cos}^+(\mathcal{M}')$ of $\mathcal{M}' \in \text{Mod}(X'_\bullet)$ is split exact. In particular, the unit map $u : \mathcal{M}' \rightarrow R_{(\Delta)}\mathcal{M}'_{(\Delta)}$ is an isomorphism.*

Proof. Define $s_n : \text{Cos}^+(\mathcal{M}')_n \rightarrow \text{Cos}^+(\mathcal{M}')_{n-1}$ to be

$$\begin{aligned} \text{Cos}^+(\mathcal{M}')_n &= (r_0)(n+1)_*(\mathcal{M}_{n+1}) \xrightarrow{(r_0)(n+1)_*\beta\sigma_0(n)} \\ &= (r_0)(n+1)_*s_0(n)_*(\mathcal{M}_n) = (r_0)(n)_*(\mathcal{M}_n) \text{Cos}^+(\mathcal{M}')_{n-1} \end{aligned}$$

for $n \geq 0$, and $s_{-1} : \text{Cos}^+(\mathcal{M}')_{-1} \rightarrow 0$ to be 0. It is easy to verify that s is a chain deformation of $\text{Cos}^+(\mathcal{M}')$. $\square$

26.9 Corollary. *Let the notation be as in the lemma. Then there is a functorial isomorphism*

$$R_{(\Delta)}(d_0)^*(\mathcal{M}) \rightarrow \mathcal{M}' \quad (26.10)$$

for $\mathcal{M} \in \text{EM}(X_\bullet)$. In particular, there is a functorial isomorphism

$$(R_{(\Delta)}(d_0)^*(\mathcal{M}))_{-1} \rightarrow \mathcal{M}_0. \quad (26.11)$$

Proof. The first map (26.10) is defined to be the composite

$$R_{(\Delta)}(d_0)^* \xrightarrow{R_{(\Delta)}(\alpha)} R_{(\Delta)}(?)'_{(\Delta)} \xrightarrow{u^{-1}} (?)'.$$

As $(\alpha)(\mathcal{M})$ is an isomorphism if $\mathcal{M}$ is equivariant, this is an isomorphism. The second map (26.11) is obtained from (26.10), applying $(?)_{-1}$. $\square$

The following well-known theorem in descent theory contained in [28] is now easy to prove.

26.12 Proposition. *Let $f : X \rightarrow Y$ be a morphism of S -schemes, and set $X_\bullet^+ := \text{Nerve}(f)$, and $X_\bullet := X_\bullet^+|_{(\Delta)}$. Let $\mathcal{M} \in \text{Mod}(X_\bullet)$. Then we have the following.*

0 *The counit of adjunction*

$$\varepsilon : (R_{(\Delta)}\mathcal{M})_{(\Delta)} \rightarrow \mathcal{M}$$

is an isomorphism.

1 *If f is quasi-compact separated and $\mathcal{M} \in \text{LQco}(X_\bullet)$, then $R_{(\Delta)}\mathcal{M} \in \text{LQco}(X_\bullet^+)$.*

- 2** If f is faithfully flat quasi-compact separated and $\mathcal{M} \in \mathbf{Qco}(X_\bullet)$, then we have $R_{(\Delta)}\mathcal{M} \in \mathbf{Qco}(X_\bullet^+)$.
- 3** If f is faithfully flat quasi-compact separated, $\mathcal{N} \in \mathbf{EM}(X_\bullet^+)$, and $\mathcal{N}|_{(\Delta)} \in \mathbf{Qco}(X_\bullet)$, then the unit of adjunction

$$u : \mathcal{N} \rightarrow R_{(\Delta)}(\mathcal{N}_{(\Delta)})$$

is an isomorphism. In particular, $\mathcal{N}$ is quasi-coherent.

- 4** If f is faithfully flat quasi-compact separated, then the restriction functor

$$(\cdot)_{(\Delta)} : \mathbf{Qco}(X_\bullet^+) \rightarrow \mathbf{Qco}(X_\bullet|_{(\Delta)})$$

is an equivalence, with $R_{(\Delta)}$ its quasi-inverse.

Proof. The assertion **0** follows from Lemma 7.11.

We prove **1**. By **0**, it suffices to prove that

$$(R_{(\Delta)}\mathcal{M})_{-1} = \text{Ker}(e(0)_*\beta_{\delta_0(1)} - e(0)_*\beta_{\delta_1(1)})$$

is quasi-coherent. This is obvious by [11, (9.2.2)].

Now we assume that f is faithfully flat quasi-compact separated, to prove the assertions **2**, **3**, and **4**.

We prove **2**. As we already know that $R_{(\Delta)}\mathcal{M}$ is locally quasi-coherent, it suffices to show that it is equivariant. As $(d_0^+) : (X_\bullet|_{(\Delta)})' \rightarrow X_\bullet$ is faithfully flat, it suffices to show that $(d_0^+)^*R_{(\Delta)}\mathcal{M}$ is equivariant, by Lemma 8.21. Now the assertion is obvious by Lemma 26.7 and Corollary 26.9, as $\mathcal{M}'$ is quasi-coherent.

We prove **3**. Note that the composite map

$$(d_0^+)^*\mathcal{N} \xrightarrow{(d_0^+)^*u} (d_0^+)^*R_{(\Delta)}(\mathcal{N}_{(\Delta)}) \rightarrow R_{(\Delta)}(d_0^*)\mathcal{N}_{(\Delta)} \cong R_{(\Delta)}((d_0^+)^*\mathcal{N})_{(\Delta)} \quad (26.13)$$

is nothing but the unit of adjunction $u((d_0^+)^*\mathcal{N})$. As $(\alpha^+) : (d_0^+)^*\mathcal{N} \rightarrow (\mathcal{N}_{(\Delta)})'$ is an isomorphism since $\mathcal{N}$ is equivariant, we have that $u((d_0^+)^*\mathcal{N})$ is an isomorphism by Lemma 26.8. As the second arrow in (26.13) is an isomorphism by Lemma 26.7, we have that $(d_0^+)^*u$ is an isomorphism. As (d_0^+) is faithfully flat, we have that $u : \mathcal{N} \rightarrow R_{(\Delta)}(\mathcal{N}_{(\Delta)})$ is an isomorphism, as desired. The last assertion is obvious by **2**, and **3** is proved.

The assertion **4** is a consequence of **0**, **2** and **3**. $\square$

26.14 Corollary. *Let $f : X \rightarrow Y$ be a faithfully flat quasi-compact separated morphism of S -schemes, and $\mathcal{M} \in \text{Mod}(Y)$. Then $\mathcal{M}$ is quasi-coherent if and only if $f^*\mathcal{M}$ is.*

Proof. Only if part is trivial.

If $f^*\mathcal{M}$ is quasi-coherent, then $\mathcal{N} \in L_{-1}\mathcal{M}$ satisfies the assumption of **2** of the proposition, as can be seen easily. So $\mathcal{M} \cong (\mathcal{N})_{-1}$ is quasi-coherent. $\square$

26.15 Corollary. *Let the notation be as in the proposition, and assume that f is faithfully flat quasi-compact separated. The composite functor*

$$\mathbb{A} := (?)_{(\Delta)} \circ L_{-1} : \text{Qco}(Y) \rightarrow \text{Qco}(X_\bullet)$$

is an equivalence with

$$\mathbb{D} := (?)_{-1} \circ R_{(\Delta)}$$

its quasi-inverse.

Proof. Follows immediately by the proposition and Lemma 7.30, **2**, since $[-1]$ is the initial object of (Δ^+) . $\square$

We call $\mathbb{A}$ in the corollary the ascent functor, and $\mathbb{D}$ the descent functor.

26.16 Corollary. *Let the notation be as in the proposition. Then the composite functor*

$$\mathbb{A} \circ \mathbb{D} : \text{LQco}(X_\bullet) \rightarrow \text{Qco}(X_\bullet)$$

is the right adjoint functor of the inclusion $\text{Qco}(X_\bullet) \hookrightarrow \text{LQco}(X_\bullet)$.

Proof. Note that $\mathbb{D} : \text{LQco}(X_\bullet) \rightarrow \text{Qco}(Y)$ is a well-defined functor, and hence $\mathbb{A} \circ \mathbb{D}$ is a functor from $\text{LQco}(X_\bullet)$ to $\text{Qco}(X_\bullet)$.

For $\mathcal{M} \in \text{Qco}(X_\bullet)$ and $\mathcal{N} \in \text{LQco}(X_\bullet)$, we have

$$\begin{aligned} \text{Hom}_{\text{Qco}(X_\bullet)}(\mathcal{M}, \mathbb{A}\mathbb{D}\mathcal{N}) &\cong \text{Hom}_{\text{Qco}(Y)}(\mathbb{D}\mathcal{M}, \mathbb{D}\mathcal{N}) \cong \text{Hom}_{\text{LQco}(X_\bullet^+)}(R_{(\Delta)}\mathcal{M}, R_{(\Delta)}\mathcal{N}) \\ &\cong \text{Hom}_{\text{LQco}(X_\bullet)}((R_{(\Delta)}\mathcal{M})_{(\Delta)}, \mathcal{N}) \cong \text{Hom}_{\text{LQco}(X_\bullet)}(\mathcal{M}, \mathcal{N}) \end{aligned}$$

by the proposition, Corollary 26.15, and Lemma 7.30, **1**. $\square$

26.17 Corollary. *Let $X_\bullet$ be a simplicial groupoid of S -schemes, and assume that $d_0(1)$ and $d_1(1)$ are separated and quasi-compact. Then*

$$(d_0)_*^{\text{Qco}} \circ \mathbb{A} : \text{Qco}(X_0) \rightarrow \text{Qco}(X_\bullet)$$

is a right adjoint of $(?)_0 : \text{Qco}(X_\bullet) \rightarrow \text{Qco}(X_0)$, where $\mathbb{A} : \text{Qco}(X_0) \rightarrow \text{Qco}(X'_\bullet|_{(\Delta)})$ is the ascent functor defined in Corollary 26.15.

Proof. Note that $(d_0)_*^{\text{Qco}}$ is well-defined, because (d_0) is quasi-compact separated of fiber type and $X_\bullet$ has flat $(\Delta)^{\text{mon}}$ -arrows by Lemma 26.3 and the assumption. It is obvious that $\mathbb{D} \circ (d_0)_{\text{Qco}}^*$ is the left adjoint of $(d_0)_*^{\text{Qco}} \circ \mathbb{A}$ by Corollary 26.15. On the other hand, we have $(?)_0 \cong \mathbb{D} \circ (d_0)_{\text{Qco}}^*$ by Corollary 26.9. Hence, $(d_0)_*^{\text{Qco}} \circ \mathbb{A}$ is a right adjoint of $(?)_0$, as desired. $\square$

27 Local noetherian property

An abelian category $\mathcal{A}$ is called *locally noetherian* if it is a $\mathcal{U}$ -category, satisfies the (AB5) condition, and has a small set of noetherian generators [8]. For a locally noetherian category $\mathcal{A}$, we denote the full subcategory of $\mathcal{A}$ consisting of its noetherian objects by $\mathcal{A}_f$.

27.1 Lemma. *Let $\mathcal{A}$ be an abelian $\mathcal{U}$ -category which satisfies the (AB3) condition, and $\mathcal{B}$ a locally noetherian category. Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be a faithful exact functor, and G its right adjoint. If G preserves filtered inductive limits, then the following hold.*

1 $\mathcal{A}$ is locally noetherian

2 $a \in \mathcal{A}$ is a noetherian object if and only if Fa is a noetherian object.

Proof. The ‘if’ part of **2** is obvious, as F is faithful and exact. Note that $\mathcal{A}$ satisfies the (AB5) condition, as F is faithful exact and colimit preserving, and $\mathcal{B}$ satisfies the (AB5) condition.

Note also that, for $a \in \mathcal{A}$, the set of subobjects of a is small, because the set of subobjects of Fa is small [10] and F is faithful exact.

Let S be a small set of noetherian generators of $\mathcal{B}$. As any noetherian object is a quotient of a finite sum of objects in S , we may assume that any noetherian object in $\mathcal{B}$ is isomorphic to an element of S , replacing S by some larger small set, if necessary. For each $s \in S$, the set of subobjects of Gs is small by the last paragraph. Hence, there is a small subset T of $\text{ob}(\mathcal{A})$ such that, any element $t \in T$ admits a monomorphism $t \rightarrow Gs$ for some $s \in S$, Ft is noetherian, and that if $a \in \mathcal{A}$ admits a monomorphism $a \rightarrow Gs$ for some $s \in S$ and Fa noetherian then $a \cong t$ for some $t \in T$.

We claim that any $a \in \mathcal{A}$ is a filtered inductive limit $\varinjlim a_\lambda$ of subobjects a_λ of a , with each a_λ isomorphic to some element in T .

If the claim is true, then **1** is obvious, as T is a small set of noetherian generators of $\mathcal{A}$, and $\mathcal{A}$ satisfies the (AB5) condition, as we have already seen.

The ‘only if’ part of **2** is also true if the claim is true, since if $a \in \mathcal{A}$ is noetherian, then it is a quotient of a finite sum of elements of T , and hence Fa is noetherian.

It suffices to prove the claim. As $\mathcal{B}$ is locally noetherian, we have $Fa = \varinjlim b_\lambda$, where (b_λ) is the filtered inductive system of noetherian subobjects of Fa .

Let $u : \text{Id} \rightarrow GF$ be the unit of adjunction, and $\varepsilon : FG \rightarrow \text{Id}$ be the counit of adjunction. It is well-known that we have $(\varepsilon F) \circ (Fu) = \text{id}_F$. As Fu is a split monomorphism, u is also a monomorphism. We define $a_\lambda := u(a)^{-1}(Gb_\lambda)$. As G preserves filtered inductive limits and $\mathcal{A}$ satisfies the (AB5) condition, we have

$$\varinjlim a_\lambda = u(a)^{-1}(G \varinjlim b_\lambda) = u(a)^{-1}(GFa) = a.$$

Note that $a_\lambda \rightarrow Gb_\lambda$ is a monomorphism, with b_λ being noetherian.

It remains to show that Fa_λ is noetherian. Let $i_\lambda : a_\lambda \hookrightarrow a$ be the inclusion map, and $j_\lambda : b_\lambda \rightarrow Fa$ the inclusion. Then the diagram

$$\begin{array}{ccccc} Fa & \xrightarrow{Fu(a)} & FGFa & \xrightarrow{\varepsilon F(a)} & Fa \\ Fi_\lambda \uparrow & & \uparrow FGj_\lambda & & \uparrow j_\lambda \\ Fa_\lambda & \longrightarrow & FGb_\lambda & \xrightarrow{\varepsilon(b_\lambda)} & b_\lambda \end{array}$$

is commutative. As the composite of the first row is the identity map and Fi_λ is a monomorphism, we have that the composite of the second row $Fa_\lambda \rightarrow b_\lambda$ is a monomorphism. As b_λ is noetherian, we have that Fa_λ is also noetherian, as desired. $\square$

(27.2) Let S a scheme, and $X_\bullet \in \mathcal{P}((\Delta), \text{Sch}/S)$.

27.3 Lemma. *The restriction functor $(?)_0 : \text{EM}(X_\bullet) \rightarrow \text{Mod}(X_0)$ is faithful exact.*

Proof. This is obvious, because for any $[n] \in (\Delta)$, there is a morphism $[0] \rightarrow [n]$.

27.4 Lemma. *Let $X_\bullet$ be a simplicial groupoid of S -schemes, and assume that $d_0(1)$ and $d_1(1)$ are quasi-compact and separated. Assume moreover that $\text{Qco}(X_0)$ is locally noetherian. Then we have*

1 $\mathrm{Qco}(X_\bullet)$ is locally noetherian.

2 $\mathcal{M} \in \mathrm{Qco}(X_\bullet)$ is a noetherian object if and only if $\mathcal{M}_0$ is a noetherian object.

Proof. Let $F := (?)_0 : \mathrm{Qco}(X_\bullet) \rightarrow \mathrm{Qco}(X_0)$ be the restriction. By Lemma 27.3, F is faithful exact. Let $G := (d_0)_*^{\mathrm{Qco}} \circ \mathbb{A}$ be the right adjoint of F , see Corollary 26.17. As $\mathbb{A}$ is an equivalence and $(d_0)_*^{\mathrm{Qco}}$ preserves filtered inductive limits by Lemma 8.19, G preserves filtered inductive limits. As $\mathrm{Qco}(X_\bullet)$ satisfies (AB3) by Lemma 8.7, the assertion is obvious by Lemma 27.1. $\square$

The following is well-known, see [16, pp. 126–127].

27.5 Corollary. *Let Y be a noetherian separated scheme. Then $\mathrm{Qco}(Y)$ is locally noetherian, and $\mathcal{M} \in \mathrm{Qco}(Y)$ is a noetherian object if and only if it is coherent.*

Proof. This is obvious if $Y = \mathrm{Spec} A$ is affine. Now consider the general case. Let $(U_i)_{1 \leq i \leq r}$ be an affine open covering of Y , and set $X := \coprod_i U_i$. Let $p : X \rightarrow Y$ be the canonical map, and set $X_\bullet := \mathrm{Nerve}(f)|_{(\Delta)}$. Note that p is faithfully flat, quasi-compact and separated. By assumption and the lemma, we have that $\mathrm{Qco}(X_\bullet)$ is locally noetherian, and $\mathcal{M} \in \mathrm{Qco}(X_\bullet)$ is a noetherian object if and only if $\mathcal{M}_0$ is noetherian, i.e., coherent. As $\mathbb{A} : \mathrm{Qco}(Y) \rightarrow \mathrm{Qco}(X_\bullet)$ is an equivalence, we have that $\mathrm{Qco}(Y)$ is locally noetherian, and $\mathcal{M} \in \mathrm{Qco}(Y)$ is a noetherian object if and only if $(\mathbb{A}\mathcal{M})_0 = p^*\mathcal{M}$ is coherent if and only if $\mathcal{M}$ is coherent. $\square$

Now the following is obvious.

27.6 Corollary. *Let $X_\bullet$ be a simplicial groupoid of S -schemes, with $d_0(1)$ and $d_1(1)$ quasi-compact and separated. If X_0 is separated noetherian, then $\mathrm{Qco}(X_\bullet)$ is locally noetherian, and $\mathcal{M} \in \mathrm{Qco}(X_\bullet)$ is a noetherian object if and only if $\mathcal{M}_0$ is coherent.*

27.7 Lemma. *Let I be a finite category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \mathrm{Sch}/S)$. If $X_\bullet$ is separated noetherian, then $\mathrm{Mod}(X_\bullet)$ and $\mathrm{LQco}(X_\bullet)$ are locally noetherian.*

Proof. Let J be the discrete subcategory of I such that $\mathrm{ob}(J) = \mathrm{ob}(I)$. Obviously, the restriction $(?)_J$ is faithful and exact. For $i \in \mathrm{ob}(I)$, there is an isomorphism of functors $(?)_i R_J \cong \prod_{j \in \mathrm{ob}(J)} \prod_{\phi \in I(i,j)} (X_\phi)_* (?)_j$. The product is a finite product, as I is finite. As each X_ϕ is quasi-compact,

$(X_\phi)_*(?)_j$ preserves filtered inductive limits. Hence R_J preserves filtered inductive limits. Note also that R_J preserves local quasi-coherence.

Hence we may assume that I is a discrete finite category, which case is trivial by [15, Theorem II.7.8] and Corollary 27.5. $\square$

28 Groupoid of schemes

(28.1) Let $\mathcal{C}$ be a category with finite limits. A $\mathcal{C}$ -groupoid X_* is a functor from $\mathcal{C}^{\text{op}}$ to the category of $\mathcal{U}$ -small groupoids (i.e., $\mathcal{U}$ -small category all of whose morphisms are isomorphisms) such that the set valued functors $X_0 := \text{ob} \circ X_*$ and $X_1 := \text{Mor} \circ X_*$ are representable.

Let X_* be a $\mathcal{C}$ -groupoid. Let us denote the source (resp. target) $X_1 \rightarrow X_0$ by d_1 (resp. d_0). Then $X_2 := X_1 \times_{d_1 \times d_0} X_1$ represents the functor of pairs (f, g) of morphisms of X_* such that $f \circ g$ is defined. Let $d'_0 : X_2 \rightarrow X_1$ (resp. $d'_2 : X_2 \rightarrow X_1$) be the first (resp. second) projection, and $d'_1 : X_2 \rightarrow X_1$ be the composition.

By Yoneda's lemma, d_0 , d_1 , d'_0 , d'_1 , and d'_2 are morphisms of $\mathcal{C}$. $d_1 : X_1(T) \rightarrow X_0(T)$ is surjective for any $T \in \text{ob}(\mathcal{C})$. Note that the squares

$$\begin{array}{ccccc} X_2 & \xrightarrow{d'_0} & X_1 & & X_2 & \xrightarrow{d'_1} & X_1 & & X_2 & \xrightarrow{d'_1} & X_1 \\ \downarrow d'_2 & & \downarrow d_1 & & \downarrow d'_2 & & \downarrow d_1 & & \downarrow d'_0 & & \downarrow d_0 \\ X_1 & \xrightarrow{d_0} & X_0 & & X_1 & \xrightarrow{d_1} & X_0 & & X_1 & \xrightarrow{d_0} & X_0 \end{array} \quad (28.2)$$

are fiber squares. In particular,

$$X_* := X_2 \begin{array}{c} \xrightarrow{d'_0} \\ \xrightarrow{d'_1} \\ \xrightarrow{d'_2} \end{array} X_1 \begin{array}{c} \xrightarrow{d_0} \\ \xrightarrow{d_1} \end{array} X_0 \quad (28.3)$$

forms an object of $\mathcal{P}(\Delta_M, \mathcal{C})$. Finally, by the associativity,

$$\circ(\circ \times 1) = \circ(1 \times \circ), \quad (28.4)$$

where $\circ : X_1 \times_{d_1 \times d_0} X_1 \rightarrow X_1$ denotes the composition, or $\circ$ is the composite

$$X_1 \times_{d_1 \times d_0} X_1 \cong X_2 \xrightarrow{d'_1} X_1.$$

Conversely, a diagram $X_* \in \mathcal{P}(\Delta_M, \mathcal{C})$ as in (28.3) such that the squares in (28.2) are fiber squares, $d_1(T) : X_1(T) \rightarrow X_0(T)$ are surjective for all $T \in \text{ob}(\mathcal{C})$, and the associativity (28.4) holds gives a $\mathcal{C}$ -groupoid [9]. In the sequel, we mainly consider that a $\mathcal{C}$ -groupoid is an object of $\mathcal{P}(\Delta_M, \mathcal{C})$.

Let S be a scheme. We say that $X_\bullet \in \mathcal{P}(\Delta_M, \text{Sch}/S)$ is an S -groupoid, if $X_\bullet$ is a (Sch/S) -groupoid with flat arrows.

(28.5) Let $X_\bullet$ be a (Sch/S) -groupoid, and set $X_n := X_1 \times_{d_1 \times d_0} X_1 \times_{d_1 \times d_0} \cdots \times_{d_1 \times d_0} X_1$ (X_1 appears n times) for $n \geq 2$. For $n \geq 2$, $d_i : X_n \rightarrow X_{n-1}$ is defined by $d_0(x_{n-1}, \dots, x_1, x_0) = (x_{n-1}, \dots, x_1)$, $d_n(x_{n-1}, \dots, x_1, x_0) = (x_{n-2}, \dots, x_0)$, and $d_i(x_{n-1}, \dots, x_1, x_0) = (x_{n-2}, \dots, x_i \circ x_{i-1}, \dots, x_0)$ for $0 < i < n$. $s_i : X_n \rightarrow X_{n+1}$ is defined by $s_i(x_{n-1}, \dots, x_1, x_0) = (x_{n-1}, \dots, x_i, \text{id}, x_{i-1}, \dots, x_0)$.

It is easy to see that this gives a simplicial S -scheme $\Sigma(X_\bullet)$ such that $\Sigma(X_\bullet)|_{\Delta_M} = X_\bullet$.

For any simplicial S -scheme $Z_\bullet$ and $\psi_\bullet : Z_\bullet|_{\Delta_M} \rightarrow X_\bullet$, there exists some unique $\varphi_\bullet : Z_\bullet \rightarrow \Sigma(X_\bullet)$ such that $\varphi|_{\Delta_M} : Z_\bullet|_{\Delta_M} \rightarrow \Sigma(X_\bullet)|_{\Delta_M} = X_\bullet$ equals ψ . Indeed, φ is given by

$$\varphi_n(z) = (\psi_1(Q_{n-1}(z)), \dots, \psi_1(Q_0(z))),$$

where $q_i : [1] \rightarrow [n]$ is the injective monotone map such that $\text{Im } q_i = \{i, i+1\}$ for $0 \leq i < n$, and $Q_i : Z_n \rightarrow Z_1$ is the associated morphism. This shows that $\Sigma(X_\bullet) \cong \text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet$, and the counit map $(\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet)|_{\Delta_M} \rightarrow X_\bullet$ is an isomorphism.

Note that under the identification $\Sigma(X_\bullet)_{n+1} \cong X_{nr_0 \times d_0} X_1$, the morphism $d_0 : \Sigma(X_\bullet)_{n+1} \rightarrow \Sigma(X_\bullet)_n$ is nothing but the first projection. So $(d_0) : \Sigma(X_\bullet)' \rightarrow \Sigma(X_\bullet)$ is of fiber type. If moreover $d_0(1)$ is flat, then (d_0) is faithfully flat.

We construct an isomorphism $h_\bullet : \Sigma(X_\bullet)' \rightarrow \text{Nerve}(d_1(1))$. Define $h_{-1} = \text{id}$ and $h_0 = \text{id}$. Define h_1 to be the composite

$$X_1 \times_{d_1 \times d_0} X_1 \xrightarrow{(d_0 \boxtimes d_2)^{-1}} X_2 \xrightarrow{d_1 \boxtimes d_2} X_1 \times_{d_1 \times d_1} X_1.$$

Now define h_n to be the composite

$$\begin{aligned} X_1 \times_{d_1 \times d_0} X_1 \times_{d_1 \times d_0} \cdots \times_{d_1 \times d_0} X_1 \times_{d_1 \times d_0} X_1 &\xrightarrow{1 \times h_1} X_1 \times_{d_1 \times d_0} X_1 \times_{d_1 \times d_0} \cdots \times_{d_1 \times d_0} X_1 \times_{d_1 \times d_1} X_1 \\ &\xrightarrow{\text{via } h_1} \cdots \xrightarrow{h_1 \times 1} X_1 \times_{d_1 \times d_1} X_1 \times_{d_1 \times d_1} \cdots \times_{d_1 \times d_1} X_1 \times_{d_1 \times d_1} X_1. \end{aligned}$$

It is straightforward to check that this gives a well-defined isomorphism $h_\bullet : \Sigma(X_\bullet)' \rightarrow \text{Nerve}(d_1(1))$. In conclusion, we have

28.6 Lemma. *If $X_\bullet$ is an S -groupoid, then $\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet$ is a simplicial S -groupoid, and the counit $\varepsilon : (\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet)|_{\Delta_M} \rightarrow X_\bullet$ is an isomorphism.*

Conversely, the following hold.

28.7 Lemma. *If $Y_\bullet$ is a simplicial S -groupoid, then $Y_\bullet|_{\Delta_M}$ is an S -groupoid, and the unit map $u : Y_\bullet \rightarrow \text{cosk}_{\Delta_M}^{(\Delta)}(Y_\bullet|_{\Delta_M})$ is an isomorphism.*

Proof. It is obvious that $Y_\bullet|_{\Delta_M}$ has flat arrows. So it suffices to show that $Y_\bullet|_{\Delta_M}$ is a (Sch/ S)-groupoid. Since $Y'_\bullet$ is isomorphic to Nerve d_1 , the square

$$\begin{array}{ccc} Y_2 & \xrightarrow{d_2} & Y_1 \\ \downarrow d_1 & & \downarrow d_1 \\ Y_1 & \xrightarrow{d_1} & Y_0 \end{array}$$

is a fiber square. Since $(d_0) : Y'_\bullet|_{(\Delta)} \rightarrow Y_\bullet$ is a morphism of fiber type, the squares

$$\begin{array}{ccc} Y_2 & \xrightarrow{d_1} & Y_1 \\ \downarrow d_0 & & \downarrow d_0 \\ Y_1 & \xrightarrow{d_0} & Y_0 \end{array} \quad \begin{array}{ccc} Y_2 & \xrightarrow{d_2} & Y_1 \\ \downarrow d_0 & & \downarrow d_0 \\ Y_1 & \xrightarrow{d_1} & Y_0 \end{array}$$

are fiber squares.

As $d_1 s = \text{id}$, $d_1(T) : X_1(T) \rightarrow X_0(T)$ is surjective for any S -scheme T .

Let us denote the composite

$$X_1 \xrightarrow{d_1 \times d_0} X_1 \cong X_2 \xrightarrow{d_1} X_1$$

by $\circ$. It remains to show the associativity.

As the three squares in the diagram

$$\begin{array}{ccccc} X_3 & \xrightarrow{d_3} & X_2 & \xrightarrow{d_2} & X_1 \\ \downarrow d_0 & & \downarrow d_0 & & \downarrow d_0 \\ X_2 & \xrightarrow{d_2} & X_1 & \xrightarrow{d_1} & X_0 \\ \downarrow d_0 & & \downarrow d_0 & & \\ X_1 & \xrightarrow{d_1} & X_0 & & \end{array}$$

are all fiber squares, the canonical map $Q := Q_2 \boxtimes Q_1 \boxtimes Q_0 : X_3 \rightarrow X_1 \xrightarrow{d_1 \times d_0} X_1$ is an isomorphism. So it suffices to show that the maps

$$X_3 \xrightarrow{Q} X_1 \xrightarrow{d_1 \times d_0} X_1 \xrightarrow{d_1 \times d_0} X_1 \xrightarrow{\circ \times 1} X_1 \xrightarrow{d_1 \times d_0} X_1 \xrightarrow{\circ} X_1$$

and

$$X_3 \xrightarrow{Q} X_1 \times_{d_1 \times d_0} X_1 \xrightarrow{1 \times \circ} X_1 \times_{d_1 \times d_0} X_1 \xrightarrow{\circ} X_1$$

agree. But it is not so difficult to show that the first map is $d_1 d_2$, while the second one is $d_1 d_1$. So $Y_\bullet|_{\Delta_M}$ is an S -groupoid.

Set $Z_\bullet := \text{cosk}_{\Delta_M}^{(\Delta)}(Y_\bullet|_{\Delta_M})$, and we are to show that the unit $u_\bullet : Y_\bullet \rightarrow Z_\bullet$ is an isomorphism. Since $Y_\bullet|_{\Delta_M}$ is an S -groupoid, $\varepsilon_\bullet : Z_\bullet|_{\Delta_M} \rightarrow Y_\bullet|_{\Delta_M}$ is an isomorphism. It follows that $u_\bullet|_{\Delta_M} : Y_\bullet|_{\Delta_M} \rightarrow Z_\bullet|_{\Delta_M}$ is also an isomorphism. Hence $\text{Nerve}(d_1(1)(u_\bullet)) : \text{Nerve}(d_1(1)(Y_\bullet)) \rightarrow \text{Nerve}(d_1(1)(Z_\bullet))$ is also an isomorphism. As both $Y_\bullet$ and $Z_\bullet$ are simplicial S -groupoids by Lemma 28.6, $u'_\bullet : Y'_\bullet \rightarrow Z'_\bullet$ is an isomorphism. So $u_n : Y_n \rightarrow Z_n$ are all isomorphisms, and we are done. $\square$

28.8 Lemma. *Let S be a scheme, and $X_\bullet$ an S -groupoid, with $d_0(1)$ and $d_1(1)$ quasi-compact and separated. If X_0 is noetherian, then $\text{Qco}(X_\bullet)$ is locally noetherian, and $\mathcal{M} \in \text{Qco}(X_\bullet)$ is a noetherian object if and only if $\mathcal{M}_0$ is coherent.*

Proof. This is immediate by Corollary 27.6 and Lemma 25.4. $\square$

(28.9) Let $f : X \rightarrow Y$ be a faithfully flat quasi-compact separated S -morphism. Set $X_\bullet^+ := (\text{Nerve}(f))|_{\Delta_M^+}$ and $X_\bullet := (X_\bullet^+)|_{\Delta_M}$. We define the descent functor

$$\mathbb{D} : \text{LQco}(X_\bullet) \rightarrow \text{Qco}(Y)$$

to be the composite $(?)_{[-1]} R_{\Delta_M}$. The left adjoint $(?)_{\Delta_M} L_{[-1]}$ is denoted by $\mathbb{A}$, and called the ascent functor.

28.10 Lemma. *Let the notation be as above. Then $\mathbb{D} : \text{Qco}(X_\bullet) \rightarrow \text{Qco}(Y)$ is an equivalence, with $\mathbb{A}$ its quasi-inverse. The composite*

$$\mathbb{A} \circ \mathbb{D} : \text{LQco}(X_\bullet) \rightarrow \text{Qco}(X_\bullet)$$

is the right adjoint of the inclusion $\text{Qco}(X_\bullet) \hookrightarrow \text{LQco}(X_\bullet)$.

Proof. Follows easily from Corollary 26.15 and Corollary 26.16. $\square$

28.11 Lemma. *Let $X_\bullet$ be an S -groupoid, and assume that $d_0(1)$ and $d_1(1)$ are separated and quasi-compact. Set $Y_\bullet^+ := ((\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet)')|_{\Delta_M^+}$, and $Y_\bullet := (Y_\bullet^+)|_{\Delta_M}$. Let $(d_0) : Y_\bullet \rightarrow X_\bullet$ be the canonical map*

$$Y_\bullet = ((\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet)')|_{\Delta_M} \xrightarrow{(d_0)|_{\Delta_M}} (\text{cosk}_{\Delta_M}^{(\Delta)} X_\bullet)|_{\Delta_M} \cong X_\bullet.$$

Then (d_0) is quasi-compact separated faithfully flat of fiber type, and

$$(d_0)_*^{\text{Qco}} \circ \mathbb{A} : \text{Qco}(X_0) \rightarrow \text{Qco}(X_\bullet)$$

is a right adjoint of $(?)_0 : \text{Qco}(X_\bullet) \rightarrow \text{Qco}(X_0)$, where $\mathbb{A} : \text{Qco}(X_0) \rightarrow \text{Qco}(Y_\bullet)$ is the ascent functor.

Proof. Follows easily from Corollary 26.17. $\square$

29 Comparison of local Ext sheaves

(29.1) Let S be a scheme, and $X_\bullet$ an S -groupoid. Assume that $d_0(1)$ and $d_1(1)$ are affine faithfully flat of finite type, and X_0 is noetherian. Let $i : D(\text{Qco}(X_\bullet)) \rightarrow D(\text{Mod}(X_\bullet))$ be the inclusion. Note that $\text{Qco}(X_\bullet)$ is locally noetherian (Lemma 28.8). So any complex in $C(\text{Qco}(X_\bullet))$ admits a K -injective resolution with each term an injective object. The following is a generalization of [17, Theorem II.1.1.12].

29.2 Lemma. Let $\mathbb{F} \in K_{\text{Coh}(X_\bullet)}^-(\text{Mod}(X))$ and $\mathbb{G} \in K^+(\text{Qco}(X_\bullet))$. If $\mathbb{G}$ is a bounded below complex consisting of injective objects of $\text{Qco}(X_\bullet)$, then $\mathbb{G}$ is $\underline{\text{Hom}}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, ?)$ -acyclic.

Proof. By the way-out lemma, we may assume that $\mathbb{F}$ is a single coherent sheaf, and $\mathbb{G}$ is a single injective object of $\text{Qco}(X_\bullet)$.

To prove this case, it suffices to show that

$$\underline{\text{Ext}}_{\mathcal{O}_{X_\bullet}}^i(\mathbb{F}, \mathbb{G}) = 0$$

for $i > 0$.

Let $\{U_1, \dots, U_n\}$ be a finite affine open covering of X_0 , and set $Y := \coprod_{i=1}^n U_i$. Let $\varphi : Y \rightarrow X_0$ be the associated morphism. Set $f' : Y \times_{X_0} d_1(1) \rightarrow Y$ to be the first projection, and set $Y_\bullet^+ := \text{Nerve}(f')|_{\Delta_M^+}$ and $Y_\bullet := \text{Nerve}(f')|_{\Delta_M}$. Let $\psi_\bullet : Y_\bullet \rightarrow X'_\bullet$ be the canonical map, where $X'_\bullet := \text{Nerve}(d_1(1))|_{\Delta_M}$. As $(d_0)_* \mathbb{A} \varphi_* \cong (d_0)_*(\psi_\bullet)_* \mathbb{A}$ has a faithful left adjoint $\varphi^*(?)_0$, there exists some injective object $\mathbb{I}$ of Y such that $\mathbb{G}$ is a direct summand of $(d_0)_*(\psi_\bullet)_* \mathbb{A} \mathbb{I}$. Set $q_\bullet := (d_0) \circ \psi_\bullet$. Note that $q_\bullet$ is affine faithfully flat of finite type and fiber type. We may assume that $\mathbb{G} = (q_\bullet)_* \mathbb{A} \mathbb{I}$. As

$$R \underline{\text{Hom}}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, R(q_\bullet)_* \mathbb{A} \mathbb{I}) \cong R(q_\bullet)_* R \underline{\text{Hom}}_{\mathcal{O}_{Y_\bullet}}^\bullet(q_\bullet^* \mathbb{F}, \mathbb{A} \mathbb{I}),$$

$q_\bullet$ is affine, and $R\text{Hom}_{\mathcal{O}_{Y_\bullet}}^\bullet(q_\bullet^*\mathbb{F}, \mathbb{A}^1)$ has quasi-coherent cohomology groups, we may assume that $X_\bullet = \text{Nerve}(f)|_{\Delta_M}$ for some faithfully flat morphism $f : X \rightarrow Y$ of finite type between affine noetherian S -schemes. For each l , we have that

$$\begin{aligned} (?)_l R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{G}) &\cong R\text{Hom}_{\mathcal{O}_{X_l}}^\bullet((?)_l \mathbb{A}^1 \mathbb{D}\mathbb{F}, (?)_l \mathbb{A}^1 \mathbb{D}\mathbb{G}) \cong \\ &R\text{Hom}_{\mathcal{O}_{X_l}}^\bullet(e(l)^* \mathbb{D}\mathbb{F}, e(l)^* \mathbb{D}\mathbb{G}) \cong e(l)^* R\text{Hom}_{\mathcal{O}_Y}^\bullet(\mathbb{D}\mathbb{F}, \mathbb{D}\mathbb{G}). \end{aligned}$$

As $\mathbb{D} : \text{Qco}(X_\bullet) \rightarrow \text{Qco}(Y)$ is an equivalence, $\mathbb{D}\mathbb{G}$ is an injective object of $\text{Qco}(Y)$. Hence $\text{Ext}_{\mathcal{O}_Y}^i(\mathbb{D}\mathbb{F}, \mathbb{D}\mathbb{G}) = 0$ for $i > 0$ by Lemma 10.1, as desired. $\square$

30 Group schemes flat of finite type

(30.1) Let S be a noetherian separated scheme.

(30.2) Let $\mathcal{F}$ (resp. $\mathcal{F}_M$) denote the subcategory of $\mathcal{P}((\Delta), \text{Sch}/S)$ (resp. subcategory of $\mathcal{P}(\Delta_M, \text{Sch}/S)$) consisting of objects separated of finite type over S with flat arrows and morphisms of finite type and of fiber type.

Let G be a flat S -group scheme separated of finite type. Note that G is faithfully flat over S . Set $\mathcal{A}_G$ to be the category of S -schemes separated of finite type with left G -actions.

For $X \in \mathcal{A}_G$, we associate a simplicial scheme $B_G(X)$ by $B_G(X)_n = G^n \times X$. For $n \geq 1$, $d_i(n) : G^n \times X \rightarrow G^{n-1} \times X$ is the projection $p \times 1_{G^{n-1} \times X}$ if $i = n$, where $p : G \rightarrow S$ is the structure morphism. While $d_i(n) = 1_{G^{n-1} \times X}$ if $i = 0$, where $a : G \times X \rightarrow X$ is the action. If $0 < i < n$, then $d_i(n) = 1_{G^{n-i-1} \times \mu \times 1_{G^{i-1} \times X}}$, where $\mu : G \times G \rightarrow G$ is the product. For $n \geq 0$, $s_i(n) : G^n \times X \rightarrow G^{n+1} \times X$ is given by

$$s_i(n)(g_n, \dots, g_1, x) = (g_n, \dots, g_{i+1}, e, g_i, \dots, g_1, x),$$

where $e : S \rightarrow G$ is the unit element.

Note that $(B_G(X)')|_{(\Delta)}$ is canonically isomorphic to $B_G(G \times X)$, where $G \times X$ is viewed as a principal G -action. Note also that there is an isomorphism from $B_G(X)'$ to $\text{Nerve}(p_2 : G \times X \rightarrow X)$ given by

$$\begin{aligned} B_G(X)'_n &= G^{n+1} \times X \rightarrow (G \times X) \times_X \cdots \times_X (G \times X) = \text{Nerve}(p_2)_n \\ &(g_n, \dots, g_0, x) \mapsto ((g_n \cdots g_0, x), (g_{n-1} \cdots g_0, x), \dots, (g_0, x)). \end{aligned}$$

Hence we have

30.3 Lemma. *Let G and X be as above. Then $B_G(X)$ is a simplicial S -groupoid with $d_0(1)$ and $d_1(1)$ faithfully flat separated of finite type.*

We denote the restriction $B_G(X)|_{\Delta_M}$ by $B_G^M(X)$. Obviously, $B_G^M(X)$ is an S -groupoid with $d_0(1)$ and $d_1(1)$ faithfully flat separated of finite type.

For a morphism $f : X \rightarrow Y$ in $\mathcal{A}_G$, we define $B_G(f) : B_G(X) \rightarrow B_G(Y)$ by $(B_G(f))_n = 1_{G^n} \times f$. It is easy to check that B_G is a functor from $\mathcal{A}_G$ to $\mathcal{F}$. Thus B_G^M is a functor from $\mathcal{A}_G$ to $\mathcal{F}_M$.

By definition, a $(G, \mathcal{O}_X)$ -module (or a G -linearized $\mathcal{O}_X$ -module [27] or G -equivariant $\mathcal{O}_X$ -module) is an equivariant $\mathcal{O}_{B_G^M(X)}$ -module, see [5] and [17]. Note that the category of equivariant $\mathcal{O}_{B_G(X)}$ -modules and the category of equivariant $\mathcal{O}_{B_G^M(X)}$ -modules are equivalent (Lemma 25.4). Thus a quasi-coherent (resp. coherent) $(G, \mathcal{O}_X)$ -module is nothing but a quasi-coherent (resp. coherent) $\mathcal{O}_{B_G^M(X)}$ -module. We denote the category of $(G, \mathcal{O}_X)$ -modules by $\text{EM}(G, X)$. The category of quasi-coherent (resp. coherent) $(G, \mathcal{O}_X)$ -modules is denoted by $\text{Qco}(G, X)$ (resp. $\text{Coh}(G, X)$).

The discussion on derived categories of categories of sheaves over diagrams of schemes are interpreted to the derived categories of the categories of G -equivariant modules.

By Lemma 30.3 and Lemma 28.8, we have

30.4 Lemma. *Let $X \in \mathcal{A}_G$. Then $\text{Qco}(G, X)$ is a locally noetherian abelian category.*

Let $\mathcal{M}$ be a $(G, \mathcal{O}_X)$ -module. If there is no danger of confusion, we may write $\mathcal{M}_0$ instead of $\mathcal{M}$. For example, $\mathcal{O}_X$ means $\mathcal{O}_{B_G^M(X)}$, since $\mathcal{O}_{B_G^M(X)}$ is equivariant and $(\mathcal{O}_{B_G(X)})_0 = \mathcal{O}_X$. This abuse of notation is what we always do when $S = X = \text{Spec } k$ and G is an affine algebraic group over k . A G -module and its underlying vector space are denoted by the same symbol. Similarly, an object of $D(\text{Mod}(B_G^M(X)))$ and its restriction to $B_G^M(X)_0 = X$ are sometimes denoted by the same symbol. Moreover, for a morphism f in $\mathcal{A}_G$, we denote for example $R(B_G^M(f))_*$ by Rf_* , and $B_G^M(f)^\dagger$ by $f^\dagger$.

Thus, as a corollary to Theorem 21.1, we have

30.5 Theorem (G -Grothendieck's duality). *Let S be a noetherian separated scheme, and G a flat S -group scheme separated of finite type. Let X and Y be S -schemes separated of finite type with G -actions, and $f : X \rightarrow Y$*

a proper G -morphism. Then the composite

$$\begin{aligned} \Theta(f) : Rf_* R\underline{\mathrm{Hom}}_{\mathrm{Mod}(B_G^M(X))}^\bullet(\mathbb{F}, f^\times \mathbb{G}) &\xrightarrow{H} R\underline{\mathrm{Hom}}_{\mathrm{Mod}(B_G^M(Y))}^\bullet(Rf_* \mathbb{F}, Rf_* f^\times \mathbb{G}) \\ &\xrightarrow{\varepsilon} R\underline{\mathrm{Hom}}_{\mathrm{Mod}(B_G^M(Y))}^\bullet(Rf_* \mathbb{F}, \mathbb{G}) \end{aligned}$$

is an isomorphism in $D_{\mathrm{Qco}(G,Y)}^+(B_G^M(Y))$ for $\mathbb{F} \in D_{\mathrm{Coh}(G,X)}^-(B_G^M(X))$ and $\mathbb{G} \in D_{\mathrm{Qco}(G,Y)}^+(B_G^M(Y))$.

31 Compatibility with derived G -invariance

(31.1) Let S be a separated noetherian scheme, and G a flat separated S -group scheme of finite type. Let X be an S -scheme separated of finite type with a trivial G -action. That is, $a : G \times X \rightarrow X$ agrees with the second projection p_2 . In other words, $d_0(1) = d_1(1)$ in $B_G^M(X)$.

For an object $\mathcal{M}$ of $\mathrm{Mod}(B_G^M(X))$, we define the G -invariance of $\mathcal{M}$ to be the kernel of the natural map

$$\beta_{d_0(1)} - \beta_{d_1(1)} : \mathcal{M}_0 \rightarrow d_0(1)_* \mathcal{M}_1 = d_1(1)_* \mathcal{M}_1,$$

and we denote it by $\mathcal{M}^G$.

(31.2) Let X be as in (31.1). Define $\tilde{B}_G^M(X)$ to be the augmented diagram

$$\begin{array}{ccccc} & & \xrightarrow{1_G \times a} & & \\ G \times_S G \times_S X & \xrightarrow{\mu \times 1_X} & G \times_S X & \xrightarrow[p_2]{a} & X \xrightarrow{\mathrm{id}} X. \\ & \xrightarrow{p_{23}} & & & \end{array}$$

Note that $\tilde{B}_G^M(X)$ is an object of $\mathcal{P}(\tilde{\Delta}_M, S)$.

31.3 Lemma. The functor $(?)^G : \mathrm{Mod}(B_G^M(X)) \rightarrow \mathrm{Mod}(X)$ agrees with $(?)_{-1} R_{\Delta_M}$.

Proof. Follows easily from (7.10). $\square$

(31.4) We say that an object $\mathcal{M}$ of $B_G^M(X)$ is G -trivial if $\mathcal{M}$ is equivariant, and the canonical inclusion $\mathcal{M}^G \rightarrow \mathcal{M}_0$ is an isomorphism. Note that $(?)_{\Delta_M} L_{-1}$ is the exact left adjoint of $(?)^G$. Note also that $\mathcal{M}$ is G -trivial if and only if the counit of adjunction $\varepsilon : (?)_{\Delta_M} L_{-1} \mathcal{M}^G \rightarrow \mathcal{M}$ is an isomorphism if and only if $\mathcal{M} \cong \mathcal{N}_{\Delta_M}$ for some $\mathcal{N} \in \mathrm{EM}_{\tilde{B}_G^M(X)}$.

Let $\text{triv}(G, X)$ denote the full subcategory of $\text{Mod}(B_G(X))$ consisting of G -trivial objects. Note that $(?)^G : \text{triv}(G, X) \rightarrow \text{Mod}(X)$ is an equivalence, whose quasi-inverse is $(?)_{\Delta_M} L_{-1}$.

Note that if $\mathcal{M}$ is locally quasi-coherent, then $\mathcal{M}^G$ is quasi-coherent. Thus, we get a derived functor $R(?)^G : D_{\text{LQco}(B_G^M(X))}^+(\text{Mod}(B_G^M(X))) \rightarrow D_{\text{Qco}(X)}^+(\text{Mod}(X))$.

31.5 Proposition. *Let X and Y be S -schemes separated of finite type with trivial G -actions, and $f : X \rightarrow Y$ an S -morphism, which is automatically a G -morphism. Then there is a canonical isomorphism*

$$f^! R(?)^G \cong R(?)^G f^!$$

between functors from $D_{\text{LQco}}^+(B_G^M(Y))$ to $D_{\text{Qco}}^+(X)$.

Proof. As $(?)_{-1}$ is exact, we have $R(?)^G \cong (?)_{-1} R R_{\Delta_M}$ by Lemma 31.3. Thus, we have a composite isomorphism

$$f^! R(?)^G \cong f^! (?)_{-1} R R_{\Delta_M} \xrightarrow{\bar{\xi}^{-1}} (?)_{-1} (\tilde{B}_M^G(f))^! R R_{\Delta_M} \xrightarrow{\bar{\xi}} (?)_{-1} R R_{\Delta_M} f^! \cong R(?)^G f^!$$

by Proposition 14.14 and Lemma 20.4. $\square$

32 Equivariant dualizing complexes and canonical modules

(32.1) Let I be a finite ordered category, S a scheme, and $X_\bullet \in \mathcal{P}(I, \text{Sch}/S)$.

32.2 Lemma. *Assume that $X_\bullet$ has flat arrows. Then an object $\mathbb{I} \in K(\text{Mod}(X_\bullet))$ is K -injective if and only if $\mathbb{I}_i$ is K -injective for any $i \in \text{ob}(I)$.*

Proof. The ‘only if’ part is obvious, since $(?)_i$ has an exact left adjoint L_i .

We prove the converse by induction on the number of objects of I . We may assume that I has at least two objects. Let i be a minimal element of $\text{ob}(I)$, and set $J := \text{ob}(I) \setminus \{i\}$. Let $\mathbb{F}$ be an exact complex in $K(\text{Mod}(X_\bullet))$. Consider a distinguished triangle

$$L_i \mathbb{F}_i \xrightarrow{\varepsilon} \mathbb{F} \rightarrow \mathbb{C} \xrightarrow{(1)} \Sigma L_i \mathbb{F}_i.$$

As

$$\text{Hom}_{\text{Mod}(X_\bullet)}^\bullet(L_i \mathbb{F}_i, \mathbb{I}) \cong \text{Hom}_{\text{Mod}(X_i)}^\bullet(\mathbb{F}_i, \mathbb{I}_i)$$

is exact by assumption, it suffices to show that $\mathrm{Hom}_{\mathrm{Mod}(X_\bullet)}^\bullet(\mathbb{C}, \mathbb{I})$ is exact. Note that $\mathbb{C}$ is exact.

Consider the distinguished triangle

$$L_J \mathbb{C}_J \xrightarrow{\varepsilon} \mathbb{C} \xrightarrow{u} R_i \mathbb{C}_i \xrightarrow{(1)} \Sigma L_J \mathbb{C}_J.$$

As

$$\mathrm{Hom}_{\mathrm{Mod}(X_\bullet)}^\bullet(L_J \mathbb{C}_J, \mathbb{I}) \cong \mathrm{Hom}_{\mathrm{Mod}(X_\bullet|_J)}^\bullet(\mathbb{C}_J, \mathbb{I}_J),$$

is exact by induction assumption, it suffices to show that $\mathrm{Hom}_{\mathrm{Mod}(X_\bullet)}(R_i \mathbb{C}_i, \mathbb{I})$ is exact. To verify this, it suffices to show that $\mathbb{C}_i$ is null-homotopic. But this is trivial, since

$$(\?)_i L_i \mathbb{F}_i \xrightarrow{(\?)_i \varepsilon} \mathbb{F}_i \rightarrow \mathbb{C}_i \xrightarrow{(1)} (\?)_i \Sigma L_i \mathbb{F}_i$$

is a distinguished triangle, and $(\?)_i \varepsilon$ is an isomorphism. $\square$

32.3 Corollary. *Let I , S , and $X_\bullet$ be as in the lemma. Let J be a full subcategory of I such that $\mathrm{ob}(J)$ is a filter of $\mathrm{ob}(I)$. If $\mathbb{J}$ is a K -injective complex of $K(\mathrm{Mod}(X_\bullet|_J))$, then so is $L_J \mathbb{J}$.*

32.4 Corollary. *Let I , S , and $X_\bullet$ be as in the lemma. Let $\mathbb{F}$ be an object of $K(\mathrm{Mod}(X_\bullet))$. Then $\mathbb{F}$ has finite injective dimension if and only if $\mathbb{F}_i$ has finite injective dimension for any $i \in \mathrm{ob}(I)$.*

Proof. We prove the corollary by induction on the number of objects of I . We may assume that I has at least two objects. Let i be a minimal element of $\mathrm{ob}(I)$, and J the full subcategory whose object set is $\mathrm{ob}(I) \setminus \{i\}$. Then

$$L_J \mathbb{F}_J \xrightarrow{\varepsilon} \mathbb{F} \xrightarrow{u} R_i \mathbb{F}_i \xrightarrow{(1)} \Sigma L_J \mathbb{F}_J$$

is a distinguished triangle. As $\mathbb{F}_J$ has finite injective dimension by induction assumption, $L_J \mathbb{F}_J$ has finite injective dimension by Corollary 32.3. Since R_i is exact and has an exact left adjoint, $R_i \mathbb{F}_i$ has finite injective dimension. Hence $\mathbb{F}$ also has finite injective dimension. $\square$

(32.5) Let the notation be as in Theorem 16.4. Let $X_\bullet$ be an object of $\mathcal{F}$ (i.e., an I^{op} -diagram of separated S -schemes of finite type with flat arrows). We say that $\mathbb{F} \in D(\mathrm{Mod}(X_\bullet))$ is a *dualizing complex* of $X_\bullet$ if $\mathbb{F} \in D_{\mathrm{Coh}(X_\bullet)}^b(\mathrm{Mod}(X_\bullet))$, $\mathbb{F}$ has finite injective dimension, and the canonical map

$$\mathcal{O}_{X_\bullet} \xrightarrow{\mathrm{tr}} R \underline{\mathrm{Hom}}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{F})$$

is an isomorphism.

32.6 Lemma. *Let the notation be as in (32.5). An object $\mathbb{F} \in D(\text{Mod}(X_\bullet))$ is a dualizing complex of $X_\bullet$ if and only if $\mathbb{F}$ has equivariant cohomology groups and $\mathbb{F}_i \in D(\text{Mod}(X_i))$ is a dualizing complex of X_i for any $i \in \text{ob}(I)$.*

Proof. This is obvious by Lemma 13.9 and Corollary 32.4. $\square$

32.7 Corollary. *Let the notation be as in (32.5). If $X_\bullet$ is Gorenstein with finite Krull dimension, then $\mathcal{O}_{X_\bullet}$ is a dualizing complex of $X_\bullet$.*

Proof. This is clear by the lemma and [15, (V.10)]. $\square$

32.8 Lemma. *Let the notation be as in (32.5). If $X_\bullet$ has a dualizing complex $\mathbb{F}$, then $X_\bullet$ has finite Krull dimension, and $X_\bullet$ has Gorenstein arrows.*

Proof. As $\mathbb{F}_i$ is a dualizing complex of X_i for each $i \in \text{ob}(I)$, $X_\bullet$ has finite Krull dimension by [15, Corollary V.7.2].

Let $\phi : i \rightarrow j$ be a morphism of I . As X_ϕ is flat, $\alpha_\phi : X_\phi^* \mathbb{F}_i \rightarrow \mathbb{F}_j$ is an isomorphism of $D(\text{Mod}(X_j))$. As $X_\phi^* \mathbb{F}_i$ is a dualizing complex of X_j , X_ϕ is Gorenstein by [4, (5.1)]. $\square$

32.9 Proposition. *Let the notation be as above, and $\mathbb{I}$ a dualizing complex of $X_\bullet$. Let $\mathbb{F} \in D_{\text{Coh}(X_\bullet)}(\text{Mod}(X_\bullet))$. Then we have*

1 $R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{I}) \in D_{\text{Coh}(X_\bullet)}(\text{Mod}(X_\bullet))$.

2 *The canonical map*

$$\mathbb{F} \rightarrow R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(\mathbb{F}, \mathbb{I}), \mathbb{I})$$

is an isomorphism for $\mathbb{F} \in D_{\text{Coh}(X_\bullet)}(\text{Mod}(X_\bullet))$.

Proof. 1 As $\mathbb{I}$ has finite finite injective dimension, $R\text{Hom}_{\mathcal{O}_{X_\bullet}}^\bullet(?, \mathbb{I})$ is way-out in both directions. Hence by [15, Proposition I.7.3], we may assume that $\mathbb{F}$ is bounded. This case is trivial by Lemma 13.9 and Lemma 13.11.

2 Using Lemma 13.9 twice, we may assume that $X_\bullet$ is a single scheme. This case is [15, Proposition V.2.1]. $\square$

32.10 Lemma. *Let the notation be as in (32.5). Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be a morphism in $\mathcal{F}$, and let $\mathbb{I}$ be a dualizing complex of $Y_\bullet$. Then $f_\bullet^!(\mathbb{I})$ is a dualizing complex of $X_\bullet$.*

Proof. By Corollary 18.4, $f_{\bullet}^!(\mathbb{I})$ has coherent cohomology groups. By Lemma 32.4, $f_{\bullet}^!(\mathbb{I})$ has finite injective dimension.

By Lemma 32.6 and Proposition 14.14, we may assume that $f : X \rightarrow Y$ is a morphism of single schemes, and it suffices to show that

$$\mathcal{O}_X \xrightarrow{\text{tr}} R\mathbf{Hom}_{\mathcal{O}_X}^{\bullet}(f^!\mathbb{I}, f^!\mathbb{I})$$

is an isomorphism. As the question is local both on Y and X , We may assume that both Y and X are affine, and f is either an affine n -space or a closed immersion. This case is done in [15, Chapter V]. $\square$

32.11 Lemma. *Let the notation be as in (32.5), and $\mathbb{I}$ and $\mathbb{J}$ dualizing complexes on $X_{\bullet}$. If $X_{\bullet}$ is connected and X_i is non-empty for some $i \in \text{ob}(I)$, then there exist a unique invertible sheaf $\mathcal{L}$ and a unique integer n such that*

$$\mathbb{J} \cong \mathbb{I} \otimes_{\mathcal{O}_{X_{\bullet}}}^{\bullet, L} \mathcal{L}[n].$$

Such $\mathcal{L}$ and n are determined by

$$\mathcal{L}[n] \cong R\mathbf{Hom}_{\mathcal{O}_{X_{\bullet}}}^{\bullet}(\mathbb{I}, \mathbb{J}).$$

Proof. Use [15, Theorem V.3.1].

32.12 Definition. Let the notation be as in (32.5), and $\mathbb{I}$ a fixed dualizing complex of $X_{\bullet}$. For any object $f_{\bullet} : Y_{\bullet} \rightarrow X_{\bullet}$ of $\mathcal{F}/X_{\bullet}$, we define *the* dualizing complex of $Y_{\bullet}$ (or better, of $f_{\bullet}$) to be $f_{\bullet}^!\mathbb{I}$. If $Y_{\bullet}$ is connected and Y_i is non-empty for some $i \in \text{ob}(I)$, then we define the *canonical sheaf* $\omega_{Y_{\bullet}}$ of $Y_{\bullet}$ (or better, $f_{\bullet}$) to be $H^s(f_{\bullet}^!\mathbb{I})$, where s is the smallest i such that $H^i(f_{\bullet}^!\mathbb{I}) \neq 0$. If $Y_{\bullet}$ is disconnected, then we define $\omega_{Y_{\bullet}}$ componentwise.

Let R be a noetherian ring, $S := \text{Spec } R$, and G a flat R -group scheme separated of finite type.

32.13 Lemma. *$G \rightarrow S$ is a (flat) local complete intersection morphism. That is, (it is flat and) all fibers are locally complete intersection.*

Proof. We may assume that $S = \text{Spec } k$, with k a field. Then by [3, Theorem 1], we may assume that k is algebraically closed.

First assume that the characteristic is $p > 0$. Then there is some $r \gg 0$ such that the image of the Frobenius map $F^r : G \rightarrow G^{(r)}$ is reduced (or equivalently, k -smooth) and agrees with $G_{\text{red}}^{(r)}$. Note that the induced

morphism $G \rightarrow G_{\text{red}}^{(r)}$ is flat, since the flat locus is a G -stable open subset of G [17, Lemma 2.1.10], and the morphism is flat at the generic point.

As the group G_{red} acts on G transitively, it suffices to show that G is locally a complete intersection at the unit element e . So by [3, Theorem 2], it suffices to show that the r th Frobenius kernel G_r is a complete intersection. As G_r is finite connected, this is well known [41, (14.4)].

Now consider the case that G is of characteristic zero. We are to prove that G is k -smooth. Take a finitely generated $\mathbb{Z}$ -subalgebra R of k such that G is defined. We may take R so that G_R is R -flat. Set $H := (G_R)_{\text{red}}$. We may take R so that H is also R -flat. Then H is a closed subgroup scheme over R , since $H \times_R H$ is reduced. As a reduced group scheme over a field of characteristic zero is smooth, we may localize R if necessary, and we may assume that H is R -smooth.

Let $\mathcal{J}$ be the defining ideal sheaf of H in G_R . There exists some $s \geq 0$ such that $\mathcal{J}^{s+1} = 0$. Note that $\mathcal{G} := \bigoplus_{i=0}^s \mathcal{J}^i / \mathcal{J}^{i+1}$ is a coherent $(H, \mathcal{O}_H)$ -module. Applying Corollary 26.15 to the case that $Y = \text{Spec } R$ and $X_{\bullet} = B_H^M(H)$, the coherent $(H, \mathcal{O}_H)$ -module $\mathcal{G}$ is of the form $f^*(V)$, where V is a finite R -module, and $f : H \rightarrow \text{Spec } R$ is the structure map. Replacing R if necessary, we may assume that $V \cong R^u$. Now we want to prove that $u = 1$ so that $H = G_R$, which implies G is k -smooth.

There exists some prime number $p > u$ and a maximal ideal $\mathfrak{m}$ of R such that $R/\mathfrak{m}$ is a finite field of characteristic p . Let κ be the algebraic closure of $R/\mathfrak{m}$, and consider the base change $(\bar{?}) := ? \otimes_R \kappa$. Note that $\bar{\mathcal{G}} = \bigoplus_i \bar{\mathcal{J}}^i / \bar{\mathcal{J}}^{i+1}$, since R is $\mathbb{Z}$ -flat and H and G_R are R -flat. Let $\mathcal{I}^{[p^r]}$ denote the defining ideal of the r th Frobenius kernel of $\bar{G}_R$. By [41, (14.4)] again, $\dim_k(\mathcal{O}_{\bar{G}_R} / \mathcal{I}^{[p^r]})_e$ is a power of p , say $p^{v(r)}$. Similarly, the k -dimension of the coordinate ring $(\mathcal{O}_{\bar{G}_R} / (\bar{\mathcal{J}} + \mathcal{I}^{[p^r]}))_e$ of the r th Frobenius kernel of $\bar{H}$ is a power of p , say $p^{w(r)}$. Note that

$$p^{w(r)} \leq p^{v(r)} \leq \dim_k(\mathcal{O}_{\bar{G}_R} / \mathcal{I}^{[p^r]} \otimes_{\mathcal{O}_{\bar{G}_R}} \bar{\mathcal{G}})_e = p^{w(r)} u < p^{w(r)+1}.$$

Hence $w(r) \leq v(r) < w(r) + 1$, and we have $\bar{\mathcal{J}}_e \subset \mathcal{I}_e^{[p^r]}$ for any r . By Krull's intersection theorem, $\bar{\mathcal{J}}_e \subset \bigcap_r (\mathcal{I}_e)^{p^r} = 0$. This shows that $\bar{G}_R$ is reduced at e , which shows that $\bar{G}_R$ is κ -smooth everywhere. So the nilpotent ideal $\bar{\mathcal{J}}$ must be zero, and this shows $u = 1$. $\square$

Now assume that R is a Gorenstein local ring of dimension d .

32.14 Lemma. *$B_G(S)$ is Gorenstein of finite Krull dimension. In particular, $\mathcal{O}_S[d]$ is an equivariant dualizing complex of X (i.e., $\mathcal{O}_{B_G(S)}$ is a dualizing*

complex of $B_G(S)$).

Proof. As $S = \operatorname{Spec} R$ is Gorenstein by assumption and G is Gorenstein over S by Lemma 32.13, the assertions are trivial. $\square$

32.15 Definition. We choose and fix the equivariant dualizing complex $\mathcal{O}_S[d]$. For an object $X \in \mathcal{A}_G$, the corresponding equivariant dualizing complex is called the equivariant dualizing complex of X . The corresponding equivariant canonical sheaf is called the equivariant canonical sheaf of X , and is denoted by ω_X .

32.16 Lemma. *Let $X \in \mathcal{A}_G$, and assume that G acts on X trivially. Then the dualizing complex $\mathbb{I}_X := f^!(\mathcal{O}_S[d])$ has G -trivial cohomology groups, where $f : X \rightarrow S$ is the structure map. In particular, ω_X is G -trivial.*

Proof. By Proposition 14.14,

$$f^!(\mathcal{O}_S[d]) \cong f^!((\mathcal{O}_{\tilde{B}_G^M(X)})_{B_G^M(X)}) \cong (?)_{B_G^M(X)}(\tilde{B}_G^M(f)^!(\mathcal{O}_{\tilde{B}_G^M(X)})).$$

By Corollary 18.4, $\tilde{B}_G^M(f)^!(\mathcal{O}_{\tilde{B}_G^M(X)})$ has coherent cohomology groups. Hence, $f^!(\mathcal{O}_S[d])$ has G -trivial cohomology groups. $\square$

33 A generalization of Watanabe's theorem

33.1 Lemma. *Let R be a noetherian commutative ring, and G a finite group which acts on R . Set $A = R^G$, and assume that $\operatorname{Spec} A$ is connected. Then G permutes the connected components of R transitively.*

Proof. Since $\operatorname{Spec} R$ is a noetherian space, $\operatorname{Spec} R$ has only finitely many connected components, say $X_1, \dots, X_n$. Then $R = R_1 \times \dots \times R_n$, and each R_i is of the form Re_i , where e_i is a primitive idempotent. Note that $E := \{e_1, \dots, e_n\}$ is the set of primitive idempotents of R , and G acts on E . Let E_1 be an orbit of this action. Then $e = \sum_{e_i \in E_1} e_i$ is in A . As A does not have any nontrivial idempotent, $e = 1$. This shows that G acts on E transitively, and we are done. $\square$

33.2 Lemma. *Let R be a noetherian commutative ring, and G a finite group which acts on R . Set $A = R^G$, and assume that the inclusion $A \hookrightarrow R$ is finite. If $\mathfrak{p} \in \operatorname{Spec} A$, then G acts transitively on the set of primes of R lying over $\mathfrak{p}$.*

Proof. Note that A is noetherian by Eakin-Nagata theorem [25, Theorem 3.7]. Let A' be the $\mathfrak{p}A_{\mathfrak{p}}$ -adic completion of $A_{\mathfrak{p}}$, and set $R' := A' \otimes_A R$. As A' is A -flat, $A' = (R')^G$. It suffices to prove that G acts transitively on the maximal ideals of R' . But R' is the direct product $\prod_i R'_i$ of complete local rings R'_i . Consider the corresponding primitive idempotents. Since A' is a local ring, G permutes these idempotents transitively by Lemma 33.1. It is obvious that this action induces a transitive action on the maximal ideals of R' . $\square$

(33.3) Let k be a field, and G a finite k -group scheme. Let $S = \operatorname{Spec} R$ be an affine k -scheme of finite type with a left G -action. It gives a k -algebra automorphism action of G on R . Let $A := R^G$ be the ring of invariants.

33.4 Proposition. *Assume that G is linearly reductive (i.e., any G -module is semisimple). Then the following hold.*

- 1 *If R satisfies Serre's (S_r) condition, then the A -module R satisfies (S_r) , and A satisfies (S_r) .*
- 2 *If R is Cohen-Macaulay, then R is a maximal Cohen-Macaulay A -module, and A is also a Cohen-Macaulay ring.*
- 3 *If R is normal, then so is A .*
- 4 *If R is Cohen-Macaulay, then $\omega_R^G \cong \omega_A$ as A -modules.*
- 5 *Assume that R is Gorenstein and $\omega_R \cong R$ as (G, R) -modules. Then $A = R^G$ is Gorenstein and $\omega_A \cong A$.*

Proof. Note that the associated morphism $\pi : S = \operatorname{Spec} R \rightarrow \operatorname{Spec} A$ is finite surjective.

To prove the proposition, we may assume that $\operatorname{Spec} A$ is connected.

Set $\bar{G} := G \otimes_k \bar{k}$, and $\bar{R} = R \otimes_k \bar{k}$, where $\bar{k}$ is the algebraic closure of k . Let G_0 be the identity component (or the Frobenius kernel for sufficiently high Frobenius maps, if the characteristic is nonzero) of $\bar{G}$, which is a normal subgroup scheme of $\bar{G}$. Note that $\operatorname{Spec} \bar{R} \rightarrow \operatorname{Spec} \bar{R}^{G_0}$ is finite and is a homeomorphism, since G_0 is trivial if the characteristic is zero, and $\bar{R}^{G_0}$ contains some sufficiently high Frobenius power of $\bar{R}$, if the characteristic is positive. On the other hand, the finite group $\bar{G}(\bar{k}) = (\bar{G}/G_0)(\bar{k})$ acts on $\bar{R}^{G_0}$, and the ring of invariants under this action is $A \otimes_k \bar{k}$. By Lemma 33.2, for any prime ideal $\mathfrak{p}$ of $A \otimes_k \bar{k}$, $\bar{G}(\bar{k})$ acts transitively on the set of prime ideals

of $\bar{R}$ (or $\bar{R}^{G_0}$) lying over $\mathfrak{p}$. It follows that for any prime ideal $\mathfrak{p}$ of A and a prime ideal $\mathfrak{P}$ of R lying over $\mathfrak{p}$, we have $\text{ht } \mathfrak{p} = \text{ht } \mathfrak{P}$.

Let M be the sum of all non-trivial simple G -submodules of R . As G is linearly reductive, R is the direct sum of M and A as a G -module. It is easy to see that $R = M \oplus A$ is a direct sum decomposition as a (G, A) -module.

1 Since A is a direct summand of R as an A -module, it suffices to prove that the A -module R satisfies the (S_r) -condition. Let $\mathfrak{p} \in \text{Spec } A$ and assume that $\text{depth}_{A_{\mathfrak{p}}} R_{\mathfrak{p}} < r$. Note that $\text{depth}_{A_{\mathfrak{p}}} R_{\mathfrak{p}} = \inf_{\mathfrak{P}} \text{depth } R_{\mathfrak{P}}$, where $\mathfrak{P}$ runs through the prime ideals lying over $\mathfrak{p}$. So there exists some $\mathfrak{P}$ such that $\text{depth } R_{\mathfrak{P}} \leq \text{depth}_{A_{\mathfrak{p}}} R_{\mathfrak{p}} < r$. As R satisfies Serre's (S_r) -condition, we have that $R_{\mathfrak{P}}$ is Cohen-Macaulay. So

$$\text{ht } \mathfrak{p} = \text{ht } \mathfrak{P} = \text{depth } R_{\mathfrak{P}} \leq \text{depth}_{A_{\mathfrak{p}}} R_{\mathfrak{p}} \leq \text{depth } A_{\mathfrak{p}} \leq \text{ht } \mathfrak{p},$$

and all $\leq$ must be $=$. In particular, $R_{\mathfrak{p}}$ is a maximal Cohen-Macaulay $A_{\mathfrak{p}}$ -module. This shows that the A -module R satisfies Serre's (S_r) -condition.

2 is obvious by **1**.

3 Consider the case that k is of characteristic zero. If so, then we may assume that k is algebraically closed, G is a finite group, and $\text{Spec } A$ is connected. The action of G on R is extended to an action on the total quotient ring $Q(R)$. We have $A = R \cap Q(R)^G$. On the other hand, any nonzero element of A is a nonzerodivisor of R . In fact, if $0 \neq a \in A$ is a zerodivisor of R , then there exists some primitive idempotent e_0 of R such that $e_0 a = 0$. Let $e_0, e_1, \dots, e_r$ be the primitive idempotents of R . As G acts transitively on these, there exists some $g_i \in G$ such that $e_i = g_i e_0$ for each i . Then $a = 1a = (e_0 + \dots + e_r)a = (g_0 + \dots + g_r)(e_0 a) = 0$, which is a contradiction. So $Q(A) \subset Q(R)^G$ (in fact $=$ holds), and A is integrally closed, as can be seen easily.

So assume that k is of characteristic $p > 0$. We need to show that A satisfies the (R_1) -condition. Let $\mathfrak{p}$ be a height one prime ideal of A . As $R_{\mathfrak{p}}$ is a one-dimensional (semilocal) regular ring, the direct summand subring $A_{\mathfrak{p}}$ is strongly F -regular, see [18]. Namely, $A_{\mathfrak{p}}$ is a DVR. It follows that A satisfies the (R_1) -condition.

4 Note that $\pi : S = \text{Spec } R \rightarrow \text{Spec } A$ is a finite G -morphism. Set $d = \dim R = \dim A$. As A is Cohen-Macaulay and $\text{Spec } A$ is connected, A is equidimensional of dimension d . So $\text{ht } \mathfrak{m} = d$ for all maximal ideals of A . The same is true of R , and hence R is also equidimensional. So $\omega_R[d]$ and $\omega_A[d]$ are the equivariant dualizing complexes of R and A , respectively.

In particular, we have $\pi^! \omega_A \cong \omega_R$. By Lemma 32.16, ω_A is G -trivial. By Theorem 30.5, we have

$$\begin{aligned} \omega_R^G &\cong R(?)^G R\pi_* R \underline{\mathrm{Hom}}_{\mathcal{O}_{\mathrm{Spec} R}}(\mathcal{O}_{\mathrm{Spec} R}, \pi^! \omega_A) \\ &\cong R(?)^G R \underline{\mathrm{Hom}}_{\mathcal{O}_{\mathrm{Spec} A}}(R\pi_* \mathcal{O}_{\mathrm{Spec} R}, \omega_A). \end{aligned}$$

As π is affine, $R\pi_* \mathcal{O}_{\mathrm{Spec} R} = R$. As R is a maximal Cohen-Macaulay A -module and ω_A is a finitely generated A -module which is of finite injective dimension, we have that $\mathrm{Ext}_A^i(R, \omega_A) = 0$ ($i > 0$). As G is linearly reductive, we have that

$$R(?)^G R \underline{\mathrm{Hom}}_{\mathcal{O}_{\mathrm{Spec} A}}(R\pi_* \mathcal{O}_{\mathrm{Spec} R}, \omega_A) \cong \mathrm{Hom}_A(R, \omega_A)^G = \mathrm{Hom}_{G,A}(R, \omega_A).$$

As G is linearly reductive, there is a canonical direct sum decomposition $R \cong R^G \oplus U_R$, where U_R is the sum of all non-trivial simple G -submodules of R . As ω_A is G -trivial, $\mathrm{Hom}_G(U_R, \omega_A) = 0$. In particular, $\mathrm{Hom}_{G,A}(U_R, \omega_A) = 0$.

On the other hand, we have that

$$\mathrm{Hom}_A(R^G, \omega_A)^G = \mathrm{Hom}_A(A, \omega_A)^G = \omega_A^G = \omega_A.$$

Hence $\omega_R^G \cong \omega_A$.

5 follows from **2** and **4** immediately. $\square$

33.5 Corollary. *Let k be a field, G a linearly reductive finite k -group scheme, and V a G -module. Assume that the representation $G \rightarrow GL(V)$ factors through $SL(V)$. Then the ring of invariants $A := (\mathrm{Sym} V)^G$ is Gorenstein, and $\omega_A \cong A$.*

Proof. Set $R := \mathrm{Sym} V$. As R is k -smooth, we have that $\omega_R \cong \bigwedge^n \Omega_{R/k} \cong R \otimes \bigwedge^n V$, where $n = \dim_k V$. By assumption, $\bigwedge^n V \cong k$, and we have that $\omega_R \cong R$, as (G, R) -modules. By the proposition, A is Gorenstein and $\omega_A \cong A$. $\square$

References

- [1] L. Alonso Tarrío, A. Jeremías López, and M. J. Souto Salorio, Localization in categories of complexes and unbounded resolutions, *Canad. J. Math.* **52** (2000), 225–247.

- [2] M. Artin, “Grothendieck Topology,” Mimeographed notes, Harvard University (1962).
- [3] L. L. Avramov, Flat morphisms of complete intersections, *Dokl. Akad. Nauk SSSR* **225** (1975), *Soviet Math. Dokl.* **16** (1975), 1413–1417.
- [4] L. L. Avramov and H.-B. Foxby, Locally Gorenstein homomorphisms, *Amer. J. Math.* **114** (1992), 1007–1047.
- [5] J. Bernstein and V. Lunts, *Equivariant sheaves and functors*, *Lect. Notes Math.* **1578**, Springer Verlag (1994).
- [6] M. Bökstedt and A. Neeman, Homotopy limits in triangulated categories, *Compositio Math.* **86** (1993), 209–234.
- [7] E. M. Friedlander, “Etale homotopy of simplicial schemes,” Princeton (1982).
- [8] P. Gabriel, Des categories abeliennes, *Bull. Math. Soc. France* **90** (1962), 323–448.
- [9] P. Gabriel, Construction de préschémas quotient, in “Schémas en Groupes I (SGA3),” *Lect. Notes Math.* **151**, Springer Verlag (1970), pp. 251–286.
- [10] A. Grothendieck, Sur quelques points d’algèbre homologique, *Tôhoku Math. J.* **9** (1957), 119–221.
- [11] A. Grothendieck, “Eléments de Géométrie Algébrique I,” IHES Publ. Math. **4** (1960).
- [12] A. Grothendieck, “Eléments de Géométrie Algébrique III,” IHES Publ. Math. **11** (1961), **17** (1963).
- [13] A. Grothendieck et J.-L. Verdier, Prefaisceaux, in “Théorie des Topos et Cohomologie Etale des Schémas, SGA 4,” *Lect. Notes Math.* **269**, Springer Verlag (1972), pp. 1–217.
- [14] A. Grothendieck et J.-L. Verdier, Topos, in “Théorie des Topos et Cohomologie Etale des Schémas, SGA 4,” *Lect. Notes Math.* **269**, Springer Verlag (1972), pp. 299–519.

- [15] R. Hartshorne, “Residues and Duality,” *Lect. Notes Math.* **20**, Springer Verlag, (1966).
- [16] R. Hartshorne, “Algebraic Geometry,” *Graduate Texts in Math.* **52**, Springer Verlag (1977).
- [17] M. Hashimoto, “Auslander-Buchweitz Approximations of Equivariant Modules,” *London Mathematical Society Lecture Note Series* **282**, Cambridge (2000).
- [18] M. Hochster and C. Huneke, Tight closure and strong F -regularity, *Mém. Soc. Math. France* (N.S.) **38** (1989), 119–133.
- [19] L. Illusie, Généralités sur les conditions de finitude dans les catégories dérivées, etc., in “Théorie des Intersections et Théorème de Riemann-Roch (SGA 6),” *Lect. Notes Math.* **225**, Springer Verlag, pp. 78–273.
- [20] J. C. Jantzen, *Representations of algebraic groups*, Academic Press (1987).
- [21] G. R. Kempf, Some elementary proofs of basic theorems in the cohomology of quasi-coherent sheaves, *Rocky Mountain J. Math.* **10** (1980), 637–645.
- [22] G. Lewis, Coherence for a closed functor, in “Coherence in Categories,” *Lect. Notes Math.* **281**, Springer Verlag (1972), pp. 148–195.
- [23] J. Lipman, “Notes on Derived Categories and Derived Functors,” preprint.
- [24] S. Mac Lane, “Categories for the Working Mathematician,” Springer Verlag (1971).
- [25] H. Matsumura, “Commutative Ring Theory,” First paperback edition, Cambridge (1989).
- [26] J. S. Milne, “Étale cohomology,” Princeton (1980).
- [27] D. Mumford, J. Fogarty and F. Kirwan, *Geometric Invariant Theory*, third edition, Springer (1994).

- [28] J. P. Murre, “Lectures on an introduction to Grothendieck’s theory of the fundamental group,” Tata Institute, Bombay (1967).
- [29] M. Nagata, A generalization of the imbedding problem of an abstract variety in a complete variety, *J. Math. Kyoto Univ.* **3** (1963), 89–102.
- [30] A. Neeman, The Grothendieck duality theorem via Bousfield’s techniques and Brown representability, *J. Amer. Math. Soc.* **9** (1996), 205–236.
- [31] A. Neeman, “Triangulated Categories,” Princeton (2001).
- [32] M. Raynaud and L. Gruson, Critères de platitude et de projectivité, *Invent. Math.* **13** (1971), 1–89.
- [33] B. Saint-Donat, An appendix to P. Deligne, Cohomologie à supports propres, in “Théorie des Topos et Cohomologie Etale des Schémas, SGA 4 (Tome 3),” *Lect. Notes Math.* **305**, Springer Verlag (1973), pp. 462–480.
- [34] N. Spaltenstein, Resolutions of unbounded complexes, *Compositio Math.* **65** (1988), 121–154.
- [35] H. Sumihiro, Equivariant completion II, *J. Math. Kyoto Univ.* **15** (1975), 573–605.
- [36] J.-L. Verdier, Base change for twisted inverse images of coherent sheaves, in “Algebraic Geometry (Internat. Colloq.),” Tata Inst. Fund. Res., Bombay (1968), pp. 393–408.
- [37] J.-L. Verdier, Topologies et faisceaux, in “Théorie des Topos et Cohomologie Etale des Schémas, SGA 4,” *Lect. Notes Math.* **269**, Springer Verlag (1972), pp. 219–263.
- [38] J.-L. Verdier, Fonctorialité des catégories de faisceaux, in “Théorie des Topos et Cohomologie Etale des Schémas, SGA 4,” *Lect. Notes Math.* **269**, Springer Verlag (1972), pp. 265–297.
- [39] J.-L. Verdier, Catégories dérivées, quelques résultats (etat 0), in “Cohomologie étale, SGA 4 $\frac{1}{2}$,” *Lect. Notes Math.* **569**, Springer Verlag (1977), pp. 262–311.

- [40] K.-i. Watanabe, Certain invariant subrings are Gorenstein I, *Osaka J. Math.* **11** (1974), 1–8.
- [41] W. C. Waterhouse, “Introduction to Affine Group Schemes,” *Graduate Texts in Math.* **66**, Springer Verlag (1979).